

hep-th/9711102
IFT-UAM/CSIC 97-4

Fields, Strings and Branes[†]

César Gómez and Rafael Hernández

*Instituto de Matemáticas y Física Fundamental, CSIC
Serrano 123, 28006 Madrid, Spain*

and

*Instituto de Física Teórica, C-XVI, Universidad Autónoma de Madrid
Cantoblanco, 28049 Madrid, Spain*

[†] CIME (Summer International Center of Mathematics) lectures given by C.G. at Cetraro, Italy, 1997.

Contents

1	Chapter I	6
1.1	Dirac Monopole.	6
1.2	The ‘t Hooft-Polyakov Monopole.	8
1.3	Instantons.	10
1.4	Dyon Effect.	13
1.5	Yang-Mills Theory on T^4	15
1.5.1	The Toron Vortex.	17
1.5.2	‘t Hooft’s Toron Configurations.	19
1.6	Instanton Effective Vertex.	21
1.7	Three Dimensional Instantons.	23
1.7.1	Callias Index Theorem.	24
1.7.2	The Dual Photon as Goldstone Boson.	26
1.8	$N=1$ Supersymmetric Gauge Theories.	27
1.9	Instanton Generated Superpotentials in Three Dimensional $N=2$	31
1.9.1	A Toron Computation.	32
2	Chapter II	33
2.1	Moduli of Vacua.	34
2.2	$N=4$ Three Dimensional Yang-Mills.	35
2.3	Atiyah-Hitchin Spaces.	43
2.4	Kodaira’s Classification of Elliptic Fibrations.	45
2.5	The Moduli Space of the Four Dimensional $N=2$ Supersymmetric Yang-Mills Theory. The Seiberg-Witten Solution.	49
2.6	Effective Superpotentials.	54
3	Chapter III	62
3.1	Bosonic String.	63
3.1.1	Classical Theory.	63
3.1.2	Background Fields.	65
3.1.3	World Sheet Symmetries.	66
3.1.4	Toroidal Compactifications.	66
3.1.5	σ -Model $K3$ Geometry. A First Look at Quantum Cohomology.	68
3.1.6	Elliptically Fibered $K3$ and Mirror Symmetry.	74
3.1.7	The Open Bosonic String.	76

3.1.8	D-Branes.	76
3.1.9	Chan-Paton Factors and Wilson Lines.	78
3.2	Superstring Theories.	80
3.2.1	Toroidal Compactification of Type IIA and Type IIB Theories. <i>U</i> - duality.	82
3.2.2	Heterotic String.	87
3.2.3	Heterotic Compactifications to Four Dimensions.	91
4	Chapter IV	94
4.1	M-Theory Compactifications.	94
4.2	M-Theory Instantons.	97
4.3	D-Brane Configurations in Flat Space.	100
4.4	D-Brane Description of Seiberg-Witten Solution.	110
4.4.1	M-Theory and Strong Coupling.	117
4.5	Brane Description of $N = 1$ Four Dimensional Field Theories.	121
4.5.1	Rotation of Branes.	125
4.5.2	QCD Strings and Scales.	126
4.5.3	$N = 2$ Models with Vanishing Beta Function.	130
4.6	M-Theory and String Theory.	133
4.7	Local Models for Elliptic Fibrations.	138
4.8	Singularities of Type $\hat{D}_4$: $\mathbf{Z}_2$ Orbifolds.	140
4.9	Singularities of Type $\hat{A}_{n-1}$	141
4.10	Singularities of Type $\hat{D}_{n+4}$	143
4.11	Decompactification and Affinization.	144
4.12	M-Theory Instantons and Holomorphic Euler Characteristic.	145
4.13	θ -Parameter and Gaugino Condensates.	147
4.14	Domain Walls and Intersections.	151
A	M(atrix) Theory.	155
A.1	The Holographic Principle.	155
A.2	Toroidal Compactifications.	159
A.3	M(atrix) Theory and Quantum Directions.	161

What is your aim in philosophy?
To show the fly the way out of the fly-bottle.

Wittgenstein. *Philosophical Investigations*, 309.

Introduction

The great challenge of high energy theoretical physics is finding a consistent theory of quantum gravity. For the time being, string theory is the best candidate at hand. Many physicists think that the solution to quantum gravity will have little, if any, practical implications in our daily way of doing physics; others, more optimistic, or simply with a less practical approach to science, hope that the forthcoming theory of quantum gravity will provide a new way of thinking of quantum physics. At present, string theory is an easy target for the criticisms of pragmatics, as no experimental evidence is yet available; however, but it is also a rich and deep conceptual construction where new ways of solving longstanding theoretical problems in quantum field theory are starting to emerge. Until now, progress in string theory is mostly “internal”, a way to evolve very similar to the one underlying evolution in pure mathematics. This is not necessarily a symptom of decadence in what is traditionally considered as an experimental science, but maybe the only possible way to improve physical intuition in the quantum realm.

Until very recently, most of the work in string theory was restricted to perturbation theory. Different string theories are, from this perturbative point of view, defined by two dimensional field theories, satisfying a certain set of constraints such as conformal and modular invariance. Different orders in the string perturbative expansion are obtained by working out these two dimensional conformal field theories on Riemann surfaces of different genus, and string amplitudes become good measures on the moduli space of these surfaces. This set of rules constitutes what we now call the “world-sheet” approach to string theory. From this perturbative point of view, we can think of many different string theories, as many as two dimensional conformal field theories, with an appropriate value of the central extension, which is determined by the generic constraint that amplitudes

should define good measures on the moduli of Riemann surfaces. Among these conformal field theories, of special interest are the ones possessing a spacetime interpretation, which means that we can interpret them as describing the dynamics of strings moving in a definite target spacetime. Different string theories will then be defined as different types of strings moving in the same spacetime. Using this definition, we find, for instance, four different types of closed superstring theories (type IIA, type IIB, $E_8 \times E_8$ heterotic and $SO(32)$ heterotic) and one open superstring. However, this image of string theory has been enormously modified in the last few years, due to the clear emergence of duality symmetries. These symmetries, of two different species, perturbative and non perturbative, relate through equivalence relations a string theory on a particular spacetime to a string theory on some different spacetime. When this equivalence is perturbative, it can be proved in the genus expansion, which in practice means a general type of Montonen-Olive duality for the two dimensional conformal field theory. These duality symmetries are usually referred to as T -duality. A more ambitious type of duality relation between string theories is known as S -duality, where the equivalence is pretended to be non perturbative, and where a transformation from strongly to weakly coupled string theory is involved. Obviously, the first thing needed in order to address non perturbative duality symmetries is searching for a definition of string theory beyond perturbation theory, i. e., beyond the worldsheet approach; it is in this direction where the most ambitious program of research in string theory is focussing.

An important step in this direction comes of course from the discovery of D-branes. These new objects, which appear as necessary ingredients for extending T -duality to open strings, are sources for the Ramond fields in string theory, a part of the string spectrum not coupling, at the worldsheet level, to the string, and that are therefore not entering the allowed set of backgrounds used in the definition of the two dimensional conformal field theory. Thus, adding this backgrounds is already going beyond the worldsheet point of view and, therefore, constitutes an open window for the desired non perturbative definition of string theory.

Maybe the simplest way to address the problem of how a non perturbative definition of string theory will look like is wondering about the strong coupled behaviour of strings. This question becomes specially neat if the string theory chosen is the closed string of type IIA, where the string coupling constant can be related to the metric of eleven dimensional supergravity, so that the strongly coupled string theory can be understood as a new eleven dimensional theory, M-theory. When thinking about the relation between D-branes and M-theory or, more precisely, trying to understand the way D-branes dynamics should

be used in order to understand the eleven dimensional dynamics describing the strong coupling regime of string theory, a good answer comes again from the mysterious, for a while, relation between type IIA strings and eleven dimensional supergravity: the Kaluza-Klein modes in ten dimensions are the D-0brane sources for the Ramond $U(1)$ field. What makes this, superficially ordinary Kaluza-Klein modes, very special objects is its nature of D-branes. In fact, D-branes are sources for strings, powerful enough to provide the whole string spectrum.

A very appealing way to think of these D-0branes comes recently under the name of M(atr)ix theory. The philosophical ground for M(atr)ix theory goes back to the holographic principle, based on black hole bounds on quantum information packing in space. From this point of view, the hologram of eleven dimensional M-theory is a ten dimensional theory for the peculiar set of ten dimensional degrees of freedom in terms of which we can codify all eleven dimensional physics. M(atr)ix theory is the conjecture that D-0brane dynamics, which is a very special type of matrix quantum mechanics, is the correct hologram of the unknown eleven dimensional M-theory. We do not know the non perturbative region of string theory, but it seems we have already its healthy radiography.

These lectures were originally addressed to mathematics audience. The content covered along them is of course only a very small part of the huge amount of material growing around string theory on these days, and needless to say that it reflects the personal point of view of the authors. References are certainly not exhaustive, so that we apologize for this in advance.

Last, but not least, C. G. would like to thank the organizers and participants of the CIME school for suggestions and interesting questions, most of them yet unanswered in the text.

1 Chapter I

1.1 Dirac Monopole.

Maxwell's equations in the absence of matter,

$$\begin{aligned}\nabla \vec{E} &= 0, & \nabla \vec{B} &= 0, \\ \nabla \times \vec{B} - \frac{\partial \vec{E}}{\partial t} &= 0, & \nabla \times \vec{E} + \frac{\partial \vec{B}}{\partial t} &= 0,\end{aligned}\tag{1.1}$$

are invariant under the duality transformation

$$\begin{aligned}\vec{E} &\longrightarrow -\vec{B}, \\ \vec{B} &\longrightarrow \vec{E},\end{aligned}\tag{1.2}$$

or, equivalently,

$$\begin{aligned}F^{\mu\nu} &\longrightarrow {}^*F^{\mu\nu}, \\ {}^*F^{\mu\nu} &\longrightarrow -F^{\mu\nu},\end{aligned}\tag{1.3}$$

with ${}^*F^{\mu\nu} \equiv \tilde{F}^{\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\rho\sigma}F_{\rho\sigma}$ the Hodge dual of $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$. In the presence of both electric and magnetic matter, Maxwell's equations become

$$\begin{aligned}\partial_\nu F^{\mu\nu} &= -j^\mu \\ \partial_\nu {}^*F^{\mu\nu} &= -k^\mu,\end{aligned}\tag{1.4}$$

and (1.2) must be generalized with a transformation law for the currents,

$$\begin{aligned}F^{\mu\nu} &\rightarrow {}^*F^{\mu\nu} & j^\mu &\rightarrow k^\mu \\ {}^*F^{\mu\nu} &\rightarrow -F^{\mu\nu} & k^\mu &\rightarrow -j^\mu.\end{aligned}\tag{1.5}$$

As is clear from the definition of $F^{\mu\nu}$, the existence of magnetic sources (monopoles) [1] requires dealing with singular vector potentials. The appropriate mathematical language for describing these vector potentials is that of fiber bundles [2].

To start with, we will consider $U(1)$ bundles on the two sphere S^2 . Denoting $H^\pm$ the two hemispheres, with $H^+ \cap H^- = S^1$, the $U(1)$ bundle is defined by

$$g_\pm = e^{i\psi_\pm}\tag{1.6}$$

$U(1)$ valued functions on the two hemispheres and such that on the S^1 equator

$$e^{i\psi_+} = e^{in\varphi} e^{i\psi_-},\tag{1.7}$$

with φ the equatorial angle, and n some integer number characterizing the $U(1)$ bundle. Notice that n defines the winding number of the map

$$e^{in\varphi} : S^1 \longrightarrow U(1), \quad (1.8)$$

classified under the first homotopy group

$$\Pi_1(U(1)) \simeq \Pi_1(S^1) \simeq \mathbf{Z}. \quad (1.9)$$

Using the $U(1)$ valued functions $g_{\pm}$, we can define pure gauge connections, $A_{\mu}^{\pm}$, on $H^{\pm}$ as follows:

$$A_{\mu}^{\pm} = g_{\pm}^{-1} \partial_{\mu} g_{\pm}. \quad (1.10)$$

From (1.7) we easily get, on the equator,

$$A^+ = A^- + n\varphi, \quad (1.11)$$

and, through Stokes theorem, we get

$$\int_{S^2} F = \frac{1}{2\pi} \left[\int_{H^+} dA^+ + \int_{H^-} dA^- \right] = \frac{1}{2\pi} \int_{S^1} A^+ - A^- = n \quad (1.12)$$

identifying the winding number n with the magnetic charge of the monopole.

In quantum mechanics, the presence of a magnetic charge implies a quantization rule for the electric charge. In fact, as we require that the Schrödinger wave function, for an electric field in a monopole background, be single valued, we get

$$\exp \frac{ie}{2\pi\hbar} \oint_{\Gamma} A = 1, \quad (1.13)$$

with Γ a non contractible loop. In the presence of a magnetic charge $m \equiv \frac{1}{2\pi} \oint_{\Gamma} A$ we get Dirac's quantization rule [1],

$$em = nh. \quad (1.14)$$

Notice that the quantization rule (1.14) is equivalent to the definition (1.12) of the magnetic charge as a winding number or, more precisely, as minus the first Chern class of a $U(1)$ principal bundle on S^2 . In fact, the single valuedness of the Schrödinger wave function is equivalent to condition (1.7), where we have required n to be integer for the transition function, in order to get a manifold. The gauge connection used in (1.12) was implicitly defined as eA , with A standing for the physical gauge configuration appearing in the Schrödinger equation. From now on, we will use units with $\hbar = 1$.

The main problem with Dirac monopoles is that they are not part of the spectrum of standard QED. In order to use the idea of duality as a dynamical symmetry, we need to search for more general gauge theories, containing in the spectrum magnetically charged particles [3, 4, 5].

1.2 The ‘t Hooft-Polyakov Monopole.

Let us consider the Georgi-Glashow model [6] for $SU(2)$,

$$\mathcal{L} = -\frac{1}{4}F_a^{\mu\nu}F_{a\mu\nu} + \frac{1}{2}\mathcal{D}^\mu\phi \cdot \mathcal{D}_\mu\phi - V(\phi), \quad a = 1, 2, 3, \quad (1.15)$$

with the Higgs field in the adjoint representation, $\mathcal{D}^\mu\phi_a \equiv \partial^\mu\phi_a - g\epsilon_{abc}A_b^\mu\phi_c$ the covariant derivative, and $V(\phi)$ the Higgs potential,

$$V(\phi) = \frac{1}{4}\lambda(\phi^2 - a^2)^2. \quad (1.16)$$

with $\lambda > 0$ and a arbitrary constants.

A classical vacuum configuration is given by

$$\phi_a = a\delta_{a3}, \quad A_\mu^a = 0. \quad (1.17)$$

We can now define the vacuum manifold $\mathcal{V}$ as

$$\mathcal{V} = \{\phi, V(\phi) = 0\}, \quad (1.18)$$

which in this case is a 2-sphere of radius equal a . A necessary condition for a finite energy configuration is that at infinity, the Higgs field ϕ takes values in the vacuum manifold $\mathcal{V}$,

$$\phi : S_\infty^2 \longrightarrow \mathcal{V} \quad (1.19)$$

and that $\mathcal{D}_\mu\phi|_{S_\infty^2} = 0$. Maps of the type (1.19) are classified by the second homotopy group, $\Pi_2(\mathcal{V})$, which for the Georgi-Glashow model (with $\mathcal{V} = S^2$) is non trivial, and equal to the set of integer numbers. These maps are characterized by their winding number,

$$N = \frac{1}{4\pi a^3} \int_{S_\infty^2} dS^i \frac{1}{2} \epsilon_{ijk} \phi \cdot (\partial^j \phi \wedge \partial^k \phi). \quad (1.20)$$

Once we impose the finite energy condition $\mathcal{D}_\mu\phi|_{S_\infty^2} = 0$, the gauge field at infinity is given by

$$A^\mu = \frac{1}{a^2 g} \phi \wedge \partial^\mu \phi + \frac{1}{a} \phi f^\mu, \quad (1.21)$$

where f_μ is an arbitrary function. The corresponding stress tensor is given by

$$F_a^{\mu\nu} = \frac{1}{a} \phi_a F^{\mu\nu} = \frac{1}{a} \phi_a \left(\frac{1}{a^2 g} \phi \cdot (\partial^\mu \phi \wedge \partial^\nu \phi) + \partial^\mu f^\nu - \partial^\nu f^\mu \right), \quad (1.22)$$

which implies that the magnetic charge

$$m = -\frac{1}{2ga^3} \int_{S_\infty^2} \epsilon_{ijk} \phi \cdot (\partial^j \phi \wedge \partial^k \phi) dS^i, \quad (1.23)$$

for a finite energy configuration is given in terms of the winding number (1.20) as [7]

$$m = -\frac{4\pi N}{g}. \quad (1.24)$$

In order to combine (1.24) with Dirac's quantization rule we should define the $U(1)$ electric charge. The $U(1)$ photon field is defined by

$$A^\mu = (A^\mu \cdot \phi) \frac{1}{a}. \quad (1.25)$$

Thus, the electric charge of a field of isotopic spin j is given by

$$e = g j. \quad (1.26)$$

From (1.24) and (1.26) we recover, for $j = \frac{1}{2}$, Dirac's quantization rule.

For a generic Higgs model, with gauge group G spontaneously broken to H , the vacuum manifold $\mathcal{V}$ is given by

$$\mathcal{V} = G/H, \quad (1.27)$$

with

$$\Pi_2(G/H) \simeq \Pi_1(H)_G, \quad (1.28)$$

where $\Pi_1(H)_G$ is the set of paths in H that can be contracted to a point in G , which again contains Dirac's condition in the form (1.9).

The mass of the monopole is given by

$$M = \int d^3x \frac{1}{2} [(E_a^i)^2 + (B_a^i)^2 + (D^0 \phi_a)^2 + (D^i \phi_a)^2] + V(\phi). \quad (1.29)$$

For a static monopole, the mass becomes

$$M = \int d^3x \frac{1}{2} [(B_a^i)^2 + (D^i \phi_a)^2] + V(\phi); \quad (1.30)$$

then, in the Prasad-Sommerfeld [8] limit $\lambda = 0$ (see equation (1.16)), we get

$$M = \int d^3x \frac{1}{2} [(B_a^i + D^i \phi_a)^2 - 2B_a^i D^i \phi_a], \quad (1.31)$$

which implies the Bogomolny [9] bound $M \geq am$. The Bogomolny bound is saturated if $B_a^k = D^k \phi_a$, which are known as the Bogomolny equations.

1.3 Instantons.

Let us now consider pure $SU(N)$ Yang-Mills theory,

$$\mathcal{L} = -\frac{1}{4}F^{a\mu\nu}F_{\mu\nu}^a. \quad (1.32)$$

In euclidean spacetime $\mathbf{R}^4$, the region at infinity can be identified with the 3-sphere S^3 . A necessary condition for finite euclidean action of configurations is

$$F_{\mu\nu}^a|_{S_\infty^3} = 0, \quad (1.33)$$

or, equivalently, that the gauge configuration A^μ at infinity is a pure gauge,

$$A^\mu|_{S_\infty^3} = g(x)^{-1}\partial^\mu g(x). \quad (1.34)$$

Hence, finite euclidean action configurations are associated with maps

$$g : S^3 \rightarrow SU(N), \quad (1.35)$$

which are topologically classified in terms of the third homotopy group,

$$\Pi_3(SU(N)) \simeq \mathbf{Z}. \quad (1.36)$$

The winding number of the map g defined by (1.35) is given by

$$n = \frac{1}{24\pi^2} \int_{S^3} d^3x \epsilon_{ijk} \text{tr}[g^{-1}\nabla_i g(x)g^{-1}\nabla_j g(x)g^{-1}\nabla_k g(x)]. \quad (1.37)$$

As for the Dirac monopole construction, we can use the map g in order to define $SU(N)$ bundles on S^4 . In this case, g defines the transition function on the equator. So, for the simplest group, $SU(2)$, we will have different bundles, depending on the value of n ; in particular, for $n = 1$, we obtain the Hopf bundle

$$S^7 \longrightarrow S^4. \quad (1.38)$$

Interpreting S^4 as the compactification of euclidean space $\mathbf{R}^4$, we can define a gauge configuration on S^4 such that on the equator, which now has the topology of S^3 , we have

$$A_\mu^+ = gA_\mu^-g^{-1} + g^{-1}\partial_\mu g, \quad (1.39)$$

with A^+ and A^- the gauge configurations on the two hemispheres. Using now the relation

$$\text{tr}(F_{\mu\nu}\tilde{F}^{\mu\nu}) = d\text{tr}(F \wedge A - \frac{1}{3}A \wedge A \wedge A), \quad (1.40)$$

we get

$$-\frac{1}{8\pi^2} \int_{S^4} \text{tr}(F_{\mu\nu} \tilde{F}^{\mu\nu}) = \frac{1}{24\pi^2} \int_{S^3} \epsilon_{ijk} \text{tr}[g^{-1} \partial_i g g^{-1} \partial_j g g^{-1} \partial_k g] = n, \quad (1.41)$$

which is the generalization to S^4 of the relation we have derived above between the magnetic charge of the monopole and the winding number of the transition function defining the $U(1)$ bundle on S^2 . The topological charge defined by (1.41) is a bound for the total euclidean action. In fact,

$$\frac{1}{4} \int F^{a\mu\nu} F_{\mu\nu}^a \equiv \frac{1}{2} \int \text{tr}(F^{\mu\nu} F_{\mu\nu}) \geq \left| \frac{1}{2} \int \text{tr}(F^{\mu\nu} \tilde{F}_{\mu\nu}) \right|. \quad (1.42)$$

The instanton configuration will be defined by the gauge field saturating the bound (1.42),

$$F_{\mu\nu} = \tilde{F}_{\mu\nu}, \quad (1.43)$$

and with topological charge equal one. Bianchi identity, $DF = 0$, together with the field equations, implies $D\tilde{F} = 0$; in fact, the self duality condition (1.43) can be related to the Bogomolny equation. If we start with euclidean Yang-Mills, and reduce dimensionally to three dimensions through the definition $A_4 \equiv \phi$, we get the three dimensional Yang-Mills-Higgs lagrangian. Then, the self duality relation (1.43) becomes the Bogomolny equation.

A solution to (1.43) for $SU(2)$ was discovered by Belavin et al [10]. Including the explicit dependence on the bare coupling constant g ,

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + g[A_\mu, A_\nu], \quad (1.44)$$

the BPST solution to (1.43) is given by

$$\begin{aligned} A_\mu^a &= -\frac{2i}{g} \frac{\eta_{a\mu\nu} x^\nu}{x^2 + \rho^2}, \\ F_{\mu\nu}^a &= \frac{4i}{g} \frac{\eta_{a\mu\nu} \rho^2}{(x^2 + \rho^2)^2}, \end{aligned} \quad (1.45)$$

with $\eta_{a\mu\nu}$ satisfying $\eta_{a\mu\nu} = \eta_{aij} = \epsilon_{aij}$, $\eta_{ai0} = \delta_{ai}$, $\eta_{a\mu\nu} = -\eta_{a\mu\nu}$, and $\bar{\eta}_{a\mu\nu} = (-1)^{\delta_{\mu 0} + \delta_{\nu 0}} \eta_{a\mu\nu}$, where a, i, j take values 1, 2, 3.

The value of the action for this configuration is

$$S = \frac{8\pi^2}{g^2}, \quad (1.46)$$

with Pontryagin number

$$\frac{g^2}{32\pi^2} \int F_{\mu\nu}^a \tilde{F}^{a\mu\nu} d^4x = 1. \quad (1.47)$$

Notice that the instanton solution (1.45) depends on a free parameter ρ , that can be interpreted as the classical size of the configuration. In particular, we can consider the gauge zero modes of the instanton solution, i. e., small self dual fluctuations around the instanton solution. From (1.45), it is clear that the action is invariant under changes of the size ρ , and under translations $x^\mu \rightarrow x^\mu + a^\mu$. This means that we will have five independent gauge zero modes. The number of gauge zero modes is called, in the mathematical literature, the dimension of the moduli space of self dual solutions. This number can be computed [11, 12, 13] using index theorems [14]; the result for $SU(N)$ instantons on S^4 is

$$\dim \text{Instanton Moduli} = 4nk - n^2 + 1, \quad (1.48)$$

with k the Pontryagin number of the instanton¹. For $k = 1$ and $n = 2$ we recover the five zero modes corresponding to translations and dilatations of the solution (1.45)². The generalization of equation (1.48) to instantons on a manifold $\mathcal{M}$ is

$$\dim = 4nk - \frac{1}{2}(N^2 - 1)[\chi - \tau], \quad (1.49)$$

with χ and τ the Euler number and the signature of the manifold $\mathcal{M}$.

In order to get a clear physical interpretation of instantons, it is convenient to work in the $A^0 = 0$ temporal gauge [15, 16, 17]. If we compactify $\mathbf{R}^3$ to S^3 by imposing the boundary condition

$$A_i(\vec{r})|_{|\vec{r}| \rightarrow \infty} \rightarrow 0, \quad (1.50)$$

the vacuum configurations in this gauge are pure gauge configurations, $A_\mu = g^{-1}\partial_\mu g$, with g a map from S^3 into the gauge group $SU(N)$. We can now define different vacuum states $|n\rangle$, characterized by the winding number of the corresponding map g . In the temporal gauge, an instanton configuration of Pontryagin number equal one satisfies the following boundary conditions:

$$\begin{aligned} A_i(t = -\infty) &= 0, \\ A_i(t = +\infty) &= g_1^{-1}\partial_i g_1, \end{aligned} \quad (1.51)$$

with g_1 a map from S^3 into $SU(N)$, of winding number equal one. We can now interpret the instanton configuration (1.51) as defining a tunnelling process between the $|0\rangle$ and $|1\rangle$ vacua.

¹ k must satisfy the irreducibility condition $k \geq \frac{n}{2}$. This condition must hold if we require the gauge configuration to be irreducible, i. e., that the connection can not be obtained by embedding the connection of a smaller group.

²Observe that the total number of gauge zero modes is 4, and that $n^2 - 1$ are simply gauge rotations of the instanton configuration.

Moreover, the vacuum states $|n\rangle$ are not invariant under gauge transformations with non vanishing winding number. A vacuum state invariant under all gauge transformations would be defined by the coherent state

$$|\theta\rangle = \sum_n e^{in\theta} |n\rangle, \quad (1.52)$$

with θ a free parameter taking values in the interval $[0, 2\pi]$. Under gauge transformations of winding number m , the vacuum states $|n\rangle$ transform as

$$\mathcal{U}(g_m)|n\rangle = |n+m\rangle, \quad (1.53)$$

and therefore the θ -vacua will transform as

$$\mathcal{U}(g_m)|\theta\rangle = e^{im\theta}|\theta\rangle, \quad (1.54)$$

which means invariance in the projective sense, i. e., on the Hilbert space of rays.

The generating functional now becomes

$$\langle\theta|\theta\rangle = \sum_n \langle 0|n\rangle e^{in\theta} = \int dA \exp - \left(i \int \mathcal{L}(A) \right), \quad (1.55)$$

with the Yang-Mills lagrangian

$$\mathcal{L} = -\frac{1}{4} F^{a\mu\nu} F_{\mu\nu}^a + \frac{\theta g^2}{32\pi^2} F^{a\mu\nu} \tilde{F}_{\mu\nu}^a. \quad (1.56)$$

The θ -topological term in (1.56) breaks explicitly the CP invariance of the lagrangian. Notice that if we consider the euclidean functional integral

$$\int dA \exp - \left(\int -\frac{1}{4} F^{a\mu\nu} F_{\mu\nu}^a + \frac{i\theta g^2}{32\pi^2} F \tilde{F} \right) d^4x, \quad (1.57)$$

the instanton euclidean action becomes

$$S = \frac{8\pi^2}{g^2} + i\theta. \quad (1.58)$$

1.4 Dyon Effect.

Let us now add the topological θ -term of (1.56) to the Georgi-Glashow model (1.15). At this level, we are simply considering the θ -angle as an extra coupling constant, multiplying the topological density $F\tilde{F}$. In order to define the $U(1)$ electric charge, we can simply

apply Noether's theorem for a gauge transformation in the unbroken $U(1)$ direction [18]. An infinitesimal gauge transformation in the ϕ direction would be defined by

$$\begin{aligned}\delta A_\mu &= \frac{1}{ag} \mathcal{D}_\mu \phi, \\ \delta \phi &= 0.\end{aligned}\tag{1.59}$$

The corresponding Noether charge,

$$N = \frac{\delta \mathcal{L}}{\delta \partial_0 A} \cdot \delta A + \frac{\delta \mathcal{L}}{\delta \partial_0 \phi} \cdot \phi,\tag{1.60}$$

will be given, after the θ -term is included, by

$$N = \frac{1}{ag} \int d^3x \partial_i (\phi \cdot F_{0i}) + \frac{\theta g}{8\pi^2 a} \int d^3x \partial_i (\phi \frac{1}{2} \epsilon_{ijk} F_{jk})\tag{1.61}$$

or, in terms of the electric charge, as

$$N = \frac{e}{g} + \frac{\theta g}{8\pi^2} m.\tag{1.62}$$

Notice from (1.61) that the θ -term only contributes to N in the background of the monopole field. If we now require invariance under a $U(1)$ rotation, we get

$$e^{2\pi i N} = e^{2\pi i (\frac{e}{g} + \frac{\theta g}{8\pi^2} m)} = 1,\tag{1.63}$$

and the electric charge becomes equal to [18]

$$e = ng - \frac{\theta g^2}{8\pi^2} m,\tag{1.64}$$

which implies that a magnetic monopole of charge m becomes a dyon with electric charge $-\frac{\theta g^2}{8\pi^2} m$.

We can reach the same result, (1.63), without including a θ term in the lagrangian, if we require, for the monopole state,

$$e^{2\pi i N} |m\rangle = e^{i\theta} |m\rangle,\tag{1.65}$$

for $N = \frac{e}{g}$. Equation (1.65) implies that the monopole state transforms under $e^{2\pi i N}$ as the θ vacua with respect to gauge transformations of non vanishing winding number. However, $e^{2\pi i N}$ can be continously connected with the identity which, in physical terms, means that the induced electric charge of the monopole is independent of instantons, and is not suppressed by a tunnelling factor of the order of $\exp -\frac{8\pi^2}{g^2}$ [18, 19].

1.5 Yang-Mills Theory on T^4 .

We will consider now $SU(N)$ pure Yang-Mills on a 4-box [20], with sides of length a_0, a_1, a_2, a_3 . Let us impose periodic boundary conditions for gauge invariant quantities,

$$\begin{aligned} A^\mu(x_0 + a_0, x_1, x_2, x_3) &= \Omega_0 A^\mu(x_0, x_1, x_2, x_3), \\ A^\mu(x_0, x_1 + a_1, x_2, x_3) &= \Omega_1 A^\mu(x_0, x_1, x_2, x_3), \\ A^\mu(x_0, x_1, x_2 + a_2, x_3) &= \Omega_2 A^\mu(x_0, x_1, x_2, x_3), \\ A^\mu(x_0, x_1, x_2, x_3 + a_3) &= \Omega_3 A^\mu(x_0, x_1, x_2, x_3), \end{aligned} \quad (1.66)$$

where

$$\Omega_\rho A^\mu \equiv \Omega_\rho A^\mu \Omega_\rho^{-1} + \Omega_\rho^{-1} \partial^\mu \Omega_\rho. \quad (1.67)$$

As the gauge field transforms in the adjoint representation, we can allow the existence of $\mathbf{Z}(N)$ twists,

$$\Omega_\mu \Omega_\nu = \Omega_\nu \Omega_\mu e^{2\pi i n_{\mu\nu}/N}, \quad (1.68)$$

and therefore we can characterize different configurations in T^4 by the topological numbers $n_{\mu\nu}$. Three of these numbers, n_{12}, n_{13} and n_{23} , can be interpreted as magnetic fluxes in the 3, 2 and 1 directions, respectively. In order to characterize these magnetic fluxes, we introduce the numbers

$$m_i = \epsilon_{ijk} n_{jk}, \quad (1.69)$$

These magnetic fluxes carry $\mathbf{Z}(N)$ charge, and their topological stability is due to the fact that

$$\Pi_1(SU(N)/\mathbf{Z}(N)) \simeq \mathbf{Z}(N). \quad (1.70)$$

In order to characterize the physical Hilbert space of the theory, let us again work in the temporal gauge $A^0 = 0$. For the three dimensional box T^3 , we impose twisted boundary conditions, corresponding to magnetic flow $\vec{m} = (m_1, m_2, m_3)$. The residual gauge symmetry is defined by the set of gauge transformations preserving these boundary conditions. We may distinguish the following different types of gauge transformations:

- i) Periodic gauge transformations, which as usual are characterized by their winding number in $\Pi_3(SU(N)) \simeq \mathbf{Z}$.
- ii) Gauge transformations periodic, up to elements in the center:

$$\begin{aligned} \Omega(x_1 + a_1, x_2, x_3) &= \Omega(x_1, x_2, x_3) e^{2\pi i k_1/N}, \\ \Omega(x_1, x_2 + a_2, x_3) &= \Omega(x_1, x_2, x_3) e^{2\pi i k_2/N}, \\ \Omega(x_1, x_2, x_3 + a_3) &= \Omega(x_1, x_2, x_3) e^{2\pi i k_3/N}. \end{aligned} \quad (1.71)$$

These transformations are characterized by the vector

$$\vec{k} = (k_1, k_2, k_3), \quad (1.72)$$

and will be denoted by $\Omega_{\vec{k}}(\vec{x})$. Among this type of transformations we can extract an extra classification:

- ii-1) Those such that $(\Omega_{\vec{k}}(\vec{x}))^N$ is periodic, with vanishing Pontryagin number.
- ii-2) Those such that $(\Omega_{\vec{k}}(\vec{x}))^N$ is periodic, with non vanishing Pontryagin number.

In the temporal gauge, we can represent the transformations in ii-2) in terms of unitary operators. Let $|\Psi\rangle$ be a state in the Hilbert space $\mathcal{H}(\vec{m})$; then, we get

$$\Omega_{\vec{k}}(\vec{x})|\Psi\rangle = e^{2\pi i \frac{\vec{e} \cdot \vec{k}}{N}} e^{i\theta \frac{\vec{k} \cdot \vec{m}}{N}} |\Psi\rangle, \quad (1.73)$$

where $\vec{e}$ and θ are free parameters. Notice that the second term in (1.73) is equivalent, for $\mathbf{Z}(N)$ magnetic vortices, to the Witten dyon effect described in the previous section. In fact, we can write (1.73) in terms of an effective $\vec{e}_{eff}$,

$$\vec{e}_{eff} = \vec{e} + \frac{\theta \vec{m}}{2\pi}. \quad (1.74)$$

Moreover, as $\theta \rightarrow \theta + 2\pi$, we change $\vec{e}_{eff} \rightarrow \vec{e}_{eff} + \vec{m}$. On the other hand, the Pontryagin number of a gauge field configuration with twisted boundary conditions, determined by a set $n_{\mu\nu}$, is given by [21]

$$\frac{g^2}{16\pi^2} \int \text{tr}(F^{\mu\nu} \tilde{F}_{\mu\nu}) d^4x = k - \frac{n}{N}, \quad (1.75)$$

where $n \equiv \frac{1}{4}n_{\mu\nu}n_{\mu\nu}$. A simple way to understand the origin of the fractional piece in the above expression is noticing that, for instance, a twist n_{12} corresponds to magnetic flux in the 3-direction, with value $\frac{2\pi n_{12}}{N}$, which can be formally described by $F_{12} \sim \frac{2\pi n_{12}}{Na_1 a_2}$, and a twist n_{03} , which corresponds to an electric field in the 3-direction, is described by $F_{03} \sim \frac{2\pi n_{03}}{Na_0 a_3}$. Using now the integral representation of the Pontryagin number we easily get the fractional piece, with the right dependence on the twist coefficients (see section 1.5.2). Moreover, $(\Omega_{\vec{k}}(\vec{x}))^N$ acting on the state $|\Psi\rangle$ produces

$$(\Omega_{\vec{k}}(\vec{x}))^N |\Psi\rangle = e^{i\theta \vec{k} \cdot \vec{m}} |\Psi\rangle, \quad (1.76)$$

which means that $\vec{k} \cdot \vec{m}$ is the Pontryagin number of the periodic gauge configuration $(\Omega_{\vec{k}}(\vec{x}))^N$. For a generic gauge configuration with Pontryagin number n we will get, as usual,

$$\Omega(\vec{x}; n) |\Psi\rangle = e^{in\theta} |\Psi\rangle. \quad (1.77)$$

Using (1.71), it is easy to see that the $\vec{k}$'s characterizing the residual gauge transformations are nothing else but the n_{0i} twists. The physical interpretation of the parameter $\vec{e}$ introduced in (1.73), in the very same way as the θ -term, is that of an electric flux. In fact, we can define the Wilson loop

$$A(C) = \frac{1}{N} \text{tr} \exp \int_C ig A(\xi) d\xi, \quad (1.78)$$

with C a path in the 3-direction. Under $\Omega_{\vec{k}}(\vec{x})$, $A(C)$ transforms as

$$A(C) \rightarrow e^{2\pi i k_3/N} A(C); \quad (1.79)$$

therefore, we get

$$\Omega_{\vec{k}}(\vec{x}) A(C) |\Psi\rangle = e^{2\pi i \frac{\vec{e} \cdot \vec{k}}{N}} A(C) |\Psi\rangle, \quad (1.80)$$

which means that $A(C)$ creates a unit of electric flux in the 3-direction.

1.5.1 The Toron Vortex.

We will now consider a vacuum configuration with non vanishing magnetic flux. It may a priori come as a surprise that we can have magnetic flux for a classical vacuum configuration. What we need, in order to achieve this goal, is to find two constant matrices in the gauge group, such that [21]

$$PQ = QP\mathcal{Z} \quad (1.81)$$

with $\mathcal{Z}$ a non trivial element in the center of the group. If such matrices exist, we can use them to define twisted boundary conditions in two directions in the box. The trivial configuration $A = 0$ automatically satisfies these boundary conditions, and we will get a classical vacuum with a non vanishing magnetic flux, characterized by the center element $\mathcal{Z}$ in (1.81). For the gauge group $SU(N)$ those matrices exist; they are

$$P = \begin{pmatrix} 0 & 1 & & \\ & 0 & 1 & \\ & & & 1 \\ 1 & & & 0 \end{pmatrix},$$

$$Q = \begin{pmatrix} 1 & & & \\ & e^{2\pi i/N} & & \\ & & \ddots & \\ & & & e^{2\pi i(N-1)/N} \end{pmatrix} \cdot e^{\pi i(1-N)/N}, \quad (1.82)$$

satisfying $PQ = QPe^{2\pi i/N}$. If we impose twisted boundary conditions,

$$\begin{aligned} A(x_1 + a_1, x_2, x_3) &= PA(x_1, x_2, x_3)P^{-1}, \\ A(x_1, x_2 + a_2, x_3) &= QA(x_1, x_2, x_3)Q^{-1}, \\ A(x_1, x_2, x_3 + a_3) &= A(x_1, x_2, x_3), \end{aligned} \quad (1.83)$$

in the temporal gauge $A^0 = 0$, then the classical vacuum $A = 0$ is in the sector with non vanishing magnetic flux, $m_3 = 1$.

Classical vacuum configurations, $A_i(\vec{x}) = g^{-1}(\vec{x})\partial_i g(\vec{x})$, satisfying (1.83), would be defined by gauge transformations $g(\vec{x})$ satisfying

$$\begin{aligned} g(x_1 + a_1, x_2, x_3) &= Pg(x_1, x_2, x_3)P^{-1}e^{2\pi i k_1/N}, \\ g(x_1, x_2 + a_2, x_3) &= Qg(x_1, x_2, x_3)Q^{-1}e^{2\pi i k_2/N}, \\ g(x_1, x_2, x_3 + a_3) &= g(x_1, x_2, x_3)e^{2\pi i k_3/N}, \end{aligned} \quad (1.84)$$

for generic (k_1, k_2, k_3) . Now, any gauge transformation satisfying (1.84) can be written as

$$g = T_1^{k_1} T_2^{k_2} T_3^{k_3} \tilde{g}, \quad (1.85)$$

with

$$T_1 = Q, \quad T_2 = P^{-1}, \quad (1.86)$$

and $\tilde{g}$ satisfying (1.84), with $k_1 = k_2 = k_3 = 0$. Acting on the vacuum $|A_i = 0\rangle$, we get, from (1.86),

$$\begin{aligned} T_1|A_i = 0\rangle &= |A_i = 0\rangle, \\ T_2|A_i = 0\rangle &= |A_i = 0\rangle, \end{aligned} \quad (1.87)$$

which implies, using (1.73), that the different vacua have $e_1 = e_2 = 0$. On the other hand, we get, acting with T_3 ,

$$T_3^{k_3}|A_i = 0\rangle \equiv |A_i = 0; k_3\rangle, \quad (1.88)$$

and, therefore, we get N different vacua defined by

$$|e_3\rangle \equiv \frac{1}{N} \sum_{k_3} e^{2\pi i \frac{k_3 e_3}{N}} |A_i = 0; k_3\rangle, \quad (1.89)$$

with $e_3 = 0, \dots, N-1$. Acting now with $T_3^{k_3}$ on $|e_3\rangle$, we get

$$T_3^{k_3}|e_3\rangle = e^{2\pi i \frac{k_3 e_3}{N}} e^{i\theta \frac{k_3 m_3}{N}} |e_3\rangle, \quad (1.90)$$

from which we observe that

$$T_3^N |e_3 \rangle = e^{i\theta} |e_3 \rangle, \quad (1.91)$$

i. e., T_3^N is periodic, with winding number equal one. Notice that in the definition of $|e_3 \rangle$ we have included the θ -parameter and the magnetic flux $m_3 = 1$, associated with the boundary conditions (1.83).

From the previous discussion we learn two basic things: first, that we can get zero energy states, with both electric and magnetic flux, provided both fluxes are parallel; secondly, that the number of vacuum states with twisted boundary conditions (1.83) is equal to N . In fact, what has been computed above is the well known Witten index, $\text{tr} (-1)^F$ [22].

1.5.2 ‘t Hooft’s Toron Configurations.

We will now try to find configurations on T^4 with fractional Pontryagin number, satisfying the equations of motion. Configurations of this type were initially discovered by ‘t Hooft for $SU(N)$ [21]. In order to describe this configurations, we first choose a subgroup $SU(k) \times SU(l) \times U(1)$ of $SU(N)$, with $k + l = N$. Let ω be the matrix corresponding to the $U(1)$ generators of $SU(k) \times SU(l) \times U(1)$,

$$\omega = 2\pi \begin{pmatrix} l & & & \\ & l & & \\ & & -k & \\ & & & -k \end{pmatrix}, \quad (1.92)$$

with $\text{tr} \omega = 0$. The toron configuration is defined by

$$A_\mu(x) = -\omega \sum_\lambda \frac{\alpha_{\mu\lambda} x_\lambda}{a_\lambda a_\mu}, \quad (1.93)$$

with

$$\alpha_{\mu\nu} - \alpha_{\nu\mu} = \frac{n_{\mu\nu}^{(2)}}{Nl} - \frac{n_{\mu\nu}^{(1)}}{Nk}, \quad (1.94)$$

and

$$n_{\mu\nu} = n_{\mu\nu}^{(1)} + n_{\mu\nu}^{(2)}. \quad (1.95)$$

The stress tensor for configuration (1.93) is given by

$$F_{\mu\nu} = \omega \frac{\alpha_{\mu\nu} - \alpha_{\nu\mu}}{a_\mu a_\nu}. \quad (1.96)$$

If we consider the simplest case, $n_{12} = n_{12}^{(1)} = 1$, and $n_{30} = n_{30}^{(2)} = 1$, we will be led to

$$\begin{aligned} F_{12} &= +\omega \frac{1}{Nka_1a_2}, \\ F_{30} &= -\omega \frac{1}{Nla_3a_4}, \end{aligned} \quad (1.97)$$

and therefore

$$\frac{g^2}{16\pi^2} \int \text{tr}(F_{\mu\nu} \tilde{F}^{\mu\nu}) = -\frac{1}{N}. \quad (1.98)$$

If we now impose the self duality condition, we get

$$\frac{a_1a_2}{a_3a_4} = \frac{l}{k} = \frac{N-k}{k}, \quad (1.99)$$

which constrains the relative sizes of the box.

The gauge zero modes for the toron configuration (1.93) can be derived from the general relation (1.49), with $k = \tau = 0$ for T^4 . Thus, for Pontryagin number equal $\frac{1}{N}$, we only get four translational zero modes for gauge group $SU(N)$. In this sense, we can think of the toron as having a size equal to the size of the box.

The toron of Pontryagin number equal $\frac{1}{N}$ can be interpreted, as we did for the instanton, as a tunnelling process between states $|m_3 = 1, \vec{k} >$ and $|m_3 = 1, \vec{k} + (0, 0, 1) >$.

Let us fix a concrete distribution of electric and magnetic fluxes, characterized by $\vec{e}$ and $\vec{m}$. The functional integral for this background is given by [21]

$$< \vec{e}, \vec{m} | \vec{e}, \vec{m} > = \sum_{\vec{k}} e^{2\pi i \frac{\vec{k} \cdot \vec{e}}{N}} W(\vec{k}, \vec{m}), \quad (1.100)$$

where

$$W(\vec{k}, \vec{m}) = \int [dA]_{\vec{k}, \vec{m}} \exp - \int \mathcal{L}(A), \quad (1.101)$$

with the integral in (1.101) over gauge field configurations satisfying the twisted boundary conditions defined by the twists $(\vec{k}, \vec{m})$. We can consider the particular case $\vec{m} = (0, 0, 1)$ to define the effective action for the toron configuration,

$$S = \frac{8\pi^2}{g^2N} + \frac{2\pi i e_3}{N}. \quad (1.102)$$

A possible generalization is obtained when using configurations with Pontryagin number equal $\frac{1}{N}$, but with $\vec{k} = (k_1, k_2, 1)$. In this case, the action (1.102) becomes

$$S = \frac{8\pi^2}{g^2N} + \frac{2\pi i (\vec{k} \cdot \vec{e})}{N}. \quad (1.103)$$

It must be noticed that we have not included in (1.102) the effect of θ , which contributes to the action with a factor $\frac{i\theta}{N}$.

1.6 Instanton Effective Vertex.

Next, we will consider the effect of instantons on fermions [15, 11]. For the time being, we will work on compactified euclidean spacetime, S^4 . The Dirac matrices satisfy

$$\{\gamma^\mu, \gamma^\nu\} = -2\delta^{\mu\nu}, \quad (1.104)$$

and the chiral operator γ_5 ,

$$\gamma_5 \equiv \gamma^0 \gamma^1 \gamma^2 \gamma^3 = \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{pmatrix}. \quad (1.105)$$

The space of Dirac fermions splits into two spaces of opposite chirality,

$$\gamma_5 \psi_\pm = \pm \psi_\pm. \quad (1.106)$$

Let us work with massless Dirac fermions coupled to an instanton gauge configuration. We consider normalized solutions to Dirac's equation,

$$\gamma^\mu \mathcal{D}_\mu(A) \psi = 0. \quad (1.107)$$

As a consequence of the index theorem, the number ν_+ of solutions to (1.107) with positive chirality, minus the number of solutions with negative chirality, ν_- , is given by

$$\nu_+ - \nu_- = \frac{g^2 N_f}{32\pi^2} \int F_{\mu\nu}^a \tilde{F}^{\mu\nu a} d^4x, \quad (1.108)$$

i. e., by the topological charge of the instanton gauge configuration. Thus, the change of chirality induced by an instanton configuration is given by

$$\Delta Q_5 = 2N_f k, \quad (1.109)$$

with k the Pontryagin number, and N_f the number of different massless Dirac fermions, transforming in the fundamental representation of the gauge group. We can generalize equation (1.108) to work with instanton configurations on a generic four dimensional euclidean manifold $\mathcal{M}$. The index theorem then becomes

$$\nu_+ - \nu_- = \frac{N}{24 \cdot 8\pi^2} \int_{\mathcal{M}} \text{tr}(R \wedge R) - \frac{g^2 N_f}{32\pi^2} \int_{\mathcal{M}} F_{\mu\nu}^a \tilde{F}^{\mu\nu a} d^4x, \quad (1.110)$$

where again we consider fermions in the fundamental representation of $SU(N)$. Equation (1.109) implies that instanton configurations induce effective vertices, with change of

chirality given by (1.109). In order to compute these effective vertices, we will use a semiclassical approximation to the generating functional,

$$Z(J, \bar{J}) = \int [dA][d\bar{\psi}][d\psi] \exp - \int \mathcal{L}(A, \bar{\psi}, \psi) + J\bar{\psi} + \psi\bar{J}, \quad (1.111)$$

around the instanton configuration. Let us first perform the gaussian integration of fermions in (1.111):

$$Z(J, \bar{J}) = \int [dA] \det' \mathcal{D}(A) \exp \int \bar{J}(x) G(x, y; A) J(y) dx dy \cdot \exp - \int \mathcal{L}(A) \cdot \prod_{n(A)} \int \bar{\psi}_0^{(n)}(x) J(x) d^4x \int \bar{J}(y) \psi_0^{(n)}(y) d^4y, \quad (1.112)$$

where $\psi_0^{(n)}$ are the fermionic zero modes for the configuration A , $\det' \mathcal{D}(A)$ is the regularized determinant, and $G(x, y; A)$ is the regularized Green's function,

$$\mathcal{D}(A) G(x, y; A) = -\delta(x - y) + \sum_n \psi_0^n(x) \psi_0^n(y). \quad (1.113)$$

In semiclassical approximation around the instanton, we get

$$Z(J, \bar{J}) = \int [dQ] \det' \mathcal{D}(A_{inst}) \exp - \frac{8\pi^2}{g^2} \cdot \exp \int \bar{J}(x) G(x, y; A) J(y) d^4x d^4y \cdot \exp \int \mathcal{L}_0''(A_{inst}) Q^2 \cdot \prod_{i=1}^m \int \bar{\psi}_0^i(x) J(x) d^4x \int \bar{J}(y) \psi_0^i(y) d^4y, \quad (1.114)$$

where

$$\mathcal{L}_0''(A_{inst}) = \left(\frac{\delta^2 \mathcal{L}_0}{\delta A \delta A} \right)_{A=A_{inst}}, \quad (1.115)$$

for $\mathcal{L}_0 = -\frac{1}{4} F^{a\mu\nu} F_{\mu\nu}^a$, and Q the small fluctuation. It is clear from (1.115) that the only non vanishing amplitudes are those with

$$\left(\frac{\delta^{2m} Z(J, \bar{J})}{\delta J(x_1) \delta \bar{J}(x_1) \dots \delta J(x_m) \delta \bar{J}(x_m)} \right)_{J=\bar{J}=0}, \quad (1.116)$$

for $m = \nu^+ + \nu^-$. In order to perform the integration over Q , we need to consider the gauge zero modes. Each gauge zero mode contributes with a factor $\frac{1}{g}$. So, as we have $4N$ zero modes, we get

$$\begin{aligned} & \langle \psi(x_1) \bar{\psi}(x_1) \dots \psi(x_m) \bar{\psi}(x_m) \rangle = \\ & C \int \left(\frac{1}{g} \right)^{4N} \frac{1}{\rho^5} \rho^{3N_f} \exp - \frac{8\pi^2}{g^2(\mu)} [\mu]^{\beta_1} \prod_{i=1}^{N_f} (\bar{\psi}_0^i \psi_0^i) d^4z d\rho, \end{aligned} \quad (1.117)$$

where β_1 is the coefficient of the β -function, in such a way that the result (1.117) is independent of the renormalization point μ . It must be stressed that $d^4z d\rho \rho^{-5}$ is a translation and dilatation invariant measure. The factor ρ^{3N_f} comes from the fermionic zero modes³,

$$\psi_0^i = \frac{\rho^{3/2}}{(x^2 + \rho^2)^{3/2}} \left(\frac{2}{\pi^2} \right)^{1/2} \omega. \quad (1.118)$$

(A chiral symmetry breaking condensate is obtained in the $N_f = 1$ case). The proportionality factor C in (1.117) comes from the determinants for fermions, gauge bosons and Faddeev-Popov ghosts.

The previous computation was carried out for $\theta = 0$. The effect of including θ is simply

$$< \psi \bar{\psi}(x_1) \dots \psi \bar{\psi}(x_m) >_{\theta} = < \dots >_{\theta=0} \cdot e^{i\theta} \quad (1.119)$$

It is important to stress that the integration over the instanton size in (1.117) is infrared divergent; thus, in order to get finite instanton contributions, we should cut off the integration size, something that can be implemented if we work with a Higgs model. The so defined instantons are known as constrained instantons [11].

1.7 Three Dimensional Instantons.

An instanton in three dimensions is a finite euclidean action configuration. This necessarily implies, in order to have topological stability, that the second homotopy group of the vacuum manifold is different from zero. This can not be realized for pure gauge theories, as $\Pi_2(SU(N)) \simeq 0$, so we will consider a Higgs model with spontaneous symmetry breaking from the G gauge group to a subgroup H , such that $\Pi_2(G/H) \neq 0$. Think of $G = SU(N)$ and $H = U(1)^{N-1}$, then $\Pi_2 = \mathbf{Z}^{N-1}$. Thus, we see that three dimensional instantons are nothing but 't Hooft-Polyakov monopoles (see table).

Dimension	Energy Density	Energy	Action
1 + 1		Π_0	Π_1
2 + 1	Π_0	Π_1	Π_2
3 + 1	Π_1	Π_2	Π_3
Name	Vortex	Monopole	Instanton

The first thing to be noticed in three dimensions is that the dual to the photon is a scalar field,

$$H_\mu = *F_{\rho\sigma} \equiv \frac{1}{2} \epsilon_{\mu\rho\sigma} F^{\rho\sigma},$$

³In fact, ρ^{3N_f} is the factor that appears in the fermionic Berezin measure for the fermionic zero modes.

$$H_\mu = \partial_\mu \chi. \quad (1.120)$$

In the weak coupling regime, we can describe the dilute gas of instantons and anti-instantons as a Coulomb gas. The partition function is given by [23]

$$Z = \sum_n \int \prod_{i=1}^{n_\pm} \frac{dx_i^+ dx_i^-}{n^+! n^-!} [\exp - S_0]^{n_+ + n_-} \exp - \frac{1}{2} \left(\frac{4\pi}{g} \right)^2 \int \rho(x) \left(-\frac{1}{\partial^2} \right) \rho(y) d^3x d^3y, \quad (1.121)$$

with $n^+ + n^- = n$, S_0 the instanton action, and ρ the instanton density,

$$\rho(x) = \sum_i \delta(x - x_i^+) - \sum_i \delta(x - x_i^-). \quad (1.122)$$

The Coulomb interaction term admits the following gaussian representation, in terms of the dual photon [23]:

$$\exp - \frac{1}{2} \left(\frac{4\pi}{g} \right)^2 \int d^3x d^3y \rho(x) - \frac{1}{\partial^2} \rho(y) = \int [d\chi] \exp - \int \frac{1}{2} (\partial\chi)^2 + \frac{4\pi i \chi \rho}{g}. \quad (1.123)$$

When we sum up the instanton and anti-instanton contributions, we get the effective lagrangian for χ ,

$$\mathcal{L}_{eff}(\chi) = \frac{1}{2} (\partial\chi)^2 + e^{-S_0} \cos \frac{4\pi\chi}{e}, \quad (1.124)$$

which implies a mass for the dual photon χ equal to e^{-S_0} . That χ is the dual photon becomes clear from the $\chi - \rho$ coupling in (1.123), between χ and the magnetic density ρ . The generation of a mass for the dual photon in a dilute gas of instantons is a nice example of confinement in the sense of dual Higgs phenomena.

The inclusion of massless fermions will drastically change the physical picture. In particular, as will be shown, the photon will become a massless Goldstone boson [24]. This will be due to the existence of effective fermionic vertices induced by the three dimensional instanton, of similar type to the ones studied in previous section. In order to analyze instanton induced effective interactions in three dimensions, we should first consider the problem of fermionic zero modes in the background of a monopole.

1.7.1 Callias Index Theorem.

Consider Dirac matrices in euclidean three dimensional spacetime,

$$\gamma^i \gamma^j + \gamma^j \gamma^i = 2\delta^{ij}. \quad (1.125)$$

We can get a representation of (1.125) using constant 2×2 matrices. In general, for euclidean space of dimension n , the corresponding γ^i are constant $2^{(n-1)/2}$ matrices.

Now, we define the Dirac operator,

$$L = i\gamma^i \partial_i + \gamma^i A_i + i\Phi(x), \quad (1.126)$$

with $A_i = gT^a A_i^a$, and $\Phi(x) = \phi^a(x)T^a$, for T^a the generators of the gauge group in some particular representation. We can now consider a Dirac fermion in Minkowski $3+1$ spacetime. This is a four component spinor,

$$\psi = \begin{pmatrix} \psi_+ \\ \psi_- \end{pmatrix}. \quad (1.127)$$

Then, Dirac's equation in $3+1$ dimensions becomes

$$\begin{pmatrix} \mathbf{0} & L \\ L^+ & \mathbf{0} \end{pmatrix} \begin{pmatrix} \psi_+ \\ \psi_- \end{pmatrix} = E \begin{pmatrix} \psi_+ \\ \psi_- \end{pmatrix}, \quad (1.128)$$

for fermion fields $\psi(x, t) = \psi(x)e^{iEt}$, and where L^+ is the adjoint of L . If we consider solutions to (1.128) with $E = 0$, we get

$$\begin{aligned} L\psi_- &= 0, \\ L^+\psi_+ &= 0, \end{aligned} \quad (1.129)$$

i. e., ψ_- and ψ_+ are zero modes of the euclidean Dirac equation in three dimensions, defined by (1.126).

Now, we can define the index

$$I(L) = k_- - k_+, \quad (1.130)$$

where k_- and k_+ are, respectively, the dimensions of $\text{Ker}(L)$ and $\text{Ker}(L^+)$. By generalizing the Atiyah-Singer index theorem, Callias [25] got the following formula for $I(L)$:

$$I(L) = \frac{1}{2\left(\frac{n-1}{2}\right)} \left(\frac{i}{8\pi}\right)^{\frac{n-1}{2}} \int_{S_\infty^{n-1}} \text{tr}[U(x)(dU(x))^{n-1}], \quad (1.131)$$

with n the dimension of euclidean spacetime, and

$$U(x) \equiv |\Phi(x)|^{-1}\Phi(x). \quad (1.132)$$

In our case, $n = 3$. In terms of the magnetic charge of the monopole, (1.20),

$$N = \frac{1}{8\pi} \int \epsilon_{ijk} \phi^i \partial \phi^j \partial \phi^k, \quad (1.133)$$

where we have normalized $a = 1$ in equation (1.20), and using (1.127) for Φ we get, for $SU(2)$,

$$I(L) = 2N, \quad (1.134)$$

for fermions in the adjoint representation. Notice that in odd dimensions, the index is zero for compact spaces. The contribution in (1.131) appears because we are working in a non compact space, with special boundary conditions at infinity, which are the ones defining the monopole configuration. We can also consider the more general case of massive fermions replacing (1.127) by

$$\Phi = \phi^a T^a + m. \quad (1.135)$$

In this case, we get, from (1.131),

$$I(L) = (j(j+1) - \{m\}(\{m\}+1))N, \quad (1.136)$$

with $\{m\}$ the largest eigenvalue of $\phi^a T^a$ smaller than m or, if there is no such eigenvalue, the smallest minus one. Thus, for massless fermions in the fundamental representation we have $\{m\} = -\frac{1}{2}$, and $I(L) = N$. It is important to observe that by changing the bare mass, the index also changes (we are using the normalization $a = 1$). Thus, for $m > \frac{1}{2}$, and fermions in the fundamental representation, we get $I(L) = 0$.

1.7.2 The Dual Photon as Goldstone Boson.

We will consider the $SU(2)$ lagrangian

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^2 + \frac{1}{2}(\mathcal{D}_\mu \phi)^2 + V(\phi) + \psi_+(i\mathcal{D} + g\phi)\psi_-, \quad (1.137)$$

where we have used notation (1.127), and the Dirac operator (1.126). Lagrangian (1.137) is invariant under the $U(1)$ transformation

$$\begin{aligned} \psi_- &\rightarrow e^{i\theta} \psi_-, \\ \psi_+ &\rightarrow e^{i\theta} \psi_+. \end{aligned} \quad (1.138)$$

We will assume that the $\psi_\pm$ transform in the adjoint representation of $SU(2)$. Using (1.134), the induced instanton couple ψ_- fermions to $\psi_-^T \gamma_0$, through an instanton, while

ψ_+ is coupled to $\psi_+^T \gamma$ in the anti-instanton case (the number $\psi_+(\psi_-)$ of zero modes for spherically symmetric monopoles in the instanton (anti-instanton) configuration is zero, and the two zero modes are $\psi_\pm$ and $\psi_\pm^T \gamma_0$). These vertices induce effective mass terms for fermions with mass $\mathcal{O}(e^{-S_0})^4$.

Now, we should include the Coulomb interaction between instantons; then, the effective lagrangian becomes

$$\mathcal{L} = \frac{1}{2}(\partial\chi)^2 + m\psi_-^T \gamma_0 e^{\frac{4\pi i\chi}{g^2}} \psi_- + m\psi_+ \gamma_0 e^{-\frac{4\pi i\chi}{g^2}} \psi_+ + \dots, \quad (1.139)$$

so that now the old vertices coupling $\psi_\pm$ to $\psi_\pm^T \gamma_0$ become vertices where the instanton or anti-instanton couple $\psi_\pm$ and $\psi_\pm^T \gamma_0$ to the dual photon χ^5 . From (1.139) it is now clear that χ becomes a Goldstone boson for the $U(1)$ symmetry [24] (1.138). In fact, $\mathcal{L}$ is invariant under (1.138) if

$$\chi \rightarrow \chi + \frac{g^2 \theta}{2\pi}. \quad (1.140)$$

Notice that now χ is massless, and that no potential for χ is generated by instanton effects. It is also important to stress that the symmetry (1.138) is not anomalous in $2+1$ dimensions, which explains, from a different point of view, the Goldstone boson nature of χ .

1.8 $N=1$ Supersymmetric Gauge Theories.

As a first example, we will consider the $N=1^6$ extension of pure Yang-Mills theory. This model is defined in terms of a vector superfield, containing the gluon and the gluino. The gluino will be represented by a real Majorana spinor, transforming in the adjoint representation. The lagrangian is given by

$$\mathcal{L} = -\frac{1}{4}F^{a\mu\nu}F_{\mu\nu}^a + \frac{1}{2}i\bar{\lambda}^a \gamma_\mu \mathcal{D}^\mu(A)\lambda^a + \frac{\theta g^2}{32\pi^2}F^{a\mu\nu}F_{\mu\nu}^a. \quad (1.141)$$

As it can be easily checked, (1.141) is invariant under the supersymmetry transformations

$$\begin{aligned} \delta A_\mu^a &= i\bar{\alpha}\gamma_\mu\lambda^a, \\ \delta\lambda^a &= \frac{1}{4}[\gamma_\mu, \gamma_\nu]\alpha F^{a\mu\nu}, \\ \delta\bar{\lambda}^a &= -\bar{\alpha}\frac{1}{4}[\gamma_\mu, \gamma_\nu]F^{a\mu\nu}, \end{aligned} \quad (1.142)$$

⁴These mass terms clearly break the $U(1)$ symmetry (1.138).

⁵The effective lagrangian (1.139) will not be interpreted in the wilsonian sense, but simply as the generating functional of the effective vertices induced by instantons.

⁶For a complete reference on supersymmetry, see [26].

with α a Majorana spinor. Notice that, for λ^a in (1.141), we can use either real Majorana or complex Weyl spinors.

We will now study instanton effects for (1.141) [27, 28, 29, 30, 31, 32]. For $SU(N)$ gauge group, the total number of fermionic zero modes is

$$\#\text{zero modes} = 2Nk, \quad (1.143)$$

with k the Pontryagin number of the instanton. For $SU(2)$ and Dirac fermions in the isospin representation, of dimension $2j + 1$, the generalization of (1.108) is

$$\nu_+ - \nu_- = \frac{2}{3}(j + 1)(2j + 1)k, \quad (1.144)$$

from which we certainly get (1.143) for $j = 1$, using Majorana fermions.

The $2N$ zero modes for $k = 1$ decompose, relative to the $SU(2)$ subgroup where the instanton lies, into

$$\begin{aligned} & 4 \quad \text{triplets,} \\ & 2(N - 2) \quad \text{doublets.} \end{aligned} \quad (1.145)$$

The meaning of the 4 triplet zero modes is quite clear from supersymmetry. Namely, two of them are just the result of acting with the supersymmetric charges on the instanton configuration. For $N = 1$ we have four supersymmetric charges, two of which annihilate the instanton configuration. The two other triplets result from superconformal transformations on the instanton. In fact, lagrangian (1.141) is not only invariant under supersymmetry, but also under the superconformal group. Now, we can repeat the computation of section 1.6. The only non vanishing amplitudes will be of the type

$$< \lambda\lambda(x_1) \cdots \lambda\lambda(x_N) > . \quad (1.146)$$

Imposing the instanton measure on collective coordinates to be translation and dilatation invariant, we get

$$\int \frac{d^4z d\rho \rho^{2N}}{\rho^5}, \quad (1.147)$$

where the factor ρ^{2N} comes from the $2N$ fermionic zero modes, that scale as $\frac{1}{\rho^2}$ (see table). We must include the instanton action, and the renormalization point, μ ,

$$\mu^{4N - \frac{2N}{2}} \exp - \frac{8\pi^2}{g(\mu)^2}, \quad (1.148)$$

where the power of μ is given by +1 for each gauge zero mode, and $-\frac{1}{2}$ for each Majorana fermionic zero mode. Defining the scale,

$$\Lambda = \mu \exp - \int \frac{dg'}{\beta(g')}, \quad (1.149)$$

and using the β -function for $SU(N)$ supersymmetric Yang-Mills,

$$\beta(g') = -\frac{g'^3}{16\pi^2} 3N, \quad (1.150)$$

(1.148) becomes

$$\Lambda^{3N}, \quad (1.151)$$

with

$$\Lambda = \mu \exp -\frac{8\pi^2}{3Ng(\mu)^2}. \quad (1.152)$$

Combining all these pieces, we get

$$\begin{aligned} < \lambda\lambda(x_1) \cdots \lambda\lambda(x_N) > = \int \frac{d^4z d\rho \rho^{2N}}{\rho^5} \Lambda^{3N}. \\ \sum_{\text{permutations}} (-1)^P \text{tr}(\lambda_{i_1} \lambda_{i_2}(x_1)) \cdots \text{tr}(\lambda_{i_{2N-1}} \lambda_{i_{2N}}(x_N)). \end{aligned} \quad (1.153)$$

In order to now perform the integration over the collective coordinates, we need the expression for the zero modes given in the table⁷.

Supersymmetric triplet	$\sim \rho^2 (f(x))^2$
Superconformal triplet	$\sim \rho x (f(x))^2$
Doblets	$\rho (f(x))^{1/2}$

The fermionic zero modes in the above table are given for the instanton in the singular gauge,

$$A_\mu^{inst} = \frac{2}{g} \frac{\rho^2}{(x-z)^2 + \rho^2} \eta_{\mu\nu}^a (x-z)_\nu, \quad (1.154)$$

with z the instanton position. Using the expressions given above, we can perform the integration over z and ρ , to obtain the result

$$< \lambda\lambda(x_1) \cdots \lambda\lambda(x_N) > \sim \text{constant} \Lambda^{3N}, \quad (1.155)$$

⁷The function $f(x)$ is the instanton factor $f(x) = \frac{1}{(x-z)^2 + \rho^2}$.

which is a very amusing and, a priori, surprising result. The reason leading to (1.155) is that the integral (1.153) is saturated by instantons with size of the same order as the $|x_1 - x_N|$ distance. If we now use cluster decomposition in (1.155), we get

$$\langle \lambda\lambda \rangle \sim \text{constant} \Lambda^3 e^{2\pi i n/N}, \quad (1.156)$$

with $n = 0, \dots, N-1$. Notice that result (1.156) is not generated by instanton configurations, and that we get it assuming clustering or, equivalently, the existence of mass gap in the theory. This map gap should be interpreted as confinement.

A different approach for computing the $\langle \lambda\lambda \rangle$ condensate starts with massive supersymmetric QCD, and requires a decoupling limit, $m \rightarrow \infty$. So, for $SU(2)$ with one flavor of mass m we get, from the instanton computation

$$\langle \lambda\lambda(x_1)\lambda\lambda(x_2) \rangle \sim \text{constant} \Lambda^5 m, \quad (1.157)$$

with Λ the scale of the $N = 1$ QCD theory. Relying now upon clustering, we get

$$\langle \lambda\lambda \rangle \sim \text{constant} \Lambda^{5/2} m^{1/2} e^{2\pi i n/2}. \quad (1.158)$$

We can now take the $m \rightarrow \infty$ limit, and define the scale Λ of pure $N = 1$ supersymmetric Yang-Mills as

$$\Lambda^3 \sim \Lambda^{5/2} m^{1/2}. \quad (1.159)$$

The only difference with the previous computation is that now we perform cluster decomposition before defining the decoupling limit.

Until now we have considered $\langle \lambda\lambda \rangle$ condensates for vacuum angle θ equal zero. We will now show the dependence of the condensate on θ , through an argument given by Shifman and Vainshtein. For $SU(N)$ gauge group, the axial anomaly is given by

$$\partial j_\mu^5 = \frac{N}{16\pi^2} F \tilde{F}. \quad (1.160)$$

This means that under the chiral transformation

$$\lambda \rightarrow e^{i\alpha} \lambda, \quad (1.161)$$

the lagrangian changes as

$$\mathcal{L} \rightarrow \mathcal{L} + \frac{\alpha N}{16\pi^2} F \tilde{F}. \quad (1.162)$$

Thus, $\langle \lambda\lambda \rangle$ at a non zero value of θ is the same as $\langle \lambda'\lambda' \rangle_{\theta=0}$, with

$$\lambda' = e^{i\alpha} \lambda, \quad (1.163)$$

where now

$$2\pi\alpha = \theta. \quad (1.164)$$

Hence [33],

$$\langle \lambda\lambda \rangle_{\theta=0} = \langle \lambda'\lambda' \rangle_{\theta=0} = \langle \lambda\lambda \rangle_{\theta=0} e^{i\frac{\theta}{n}}. \quad (1.165)$$

1.9 Instanton Generated Superpotentials in Three Dimensional $N=2$.

To start with, we will consider dimensional reduction of lagrangian (1.141) to three dimensions. In this case, we arrive to the Higgs lagrangian in $2+1$ discussed in section 1.7. We can then define a complex Higgs field, with the real part given by the fourth component of A_μ in $3+1$, and the imaginary part by the photon field χ . If, as was the case in section 1.7, we consider $\langle \phi \rangle = 0$ for the real Higgs field, then we automatically break superconformal invariance, and for the $SU(2)$ case we will find only two fermionic zero modes in the instanton background ('t Hooft-Polyakov monopole). The action of the three dimensional instanton is

$$S_{\text{inst}} = \frac{4\pi\phi}{g^2}, \quad (1.166)$$

with ϕ standing for the vacuum expectation value of the Higgs field⁸. The effective lagrangian (1.139) becomes

$$\begin{aligned} \mathcal{L} = \frac{1}{2}((\partial\chi)^2 + (\partial\phi)^2) + \bar{\psi}i\partial\psi + me^{-4\pi\phi/g^2}\psi^T\gamma_0\psi e^{i4\pi\chi/g^2} + \\ me^{4\pi\phi/g^2}\bar{\psi}\gamma_0\bar{\psi}^T e^{-i4\pi\chi/g^2}, \end{aligned} \quad (1.167)$$

where we have included the kinetic term for both the real Higgs field ϕ , and the dual photon χ . In (1.167) we can define a complex Higgs field,

$$\Phi = \phi + i\chi, \quad (1.168)$$

in order to notice that the instanton is certainly generating a Yukawa coupling, which is nothing but the vertex coupling ψ fields to the dual photon χ . In order to write (1.167) as a supersymmetric lagrangian, we need to add a superpotential term of the type [24]

$$W(\Phi)_{\theta\theta} = \exp -\Phi + \text{hc}, \quad (1.169)$$

⁸Notice that the gauge coupling constant, in three dimensions, has length^{-1/2} dimensions.

which induces an effective potential for ϕ of the type

$$V(\phi) = \frac{\partial W}{\partial \Phi} \frac{\partial \bar{W}}{\partial \Phi^*} = \exp -\phi, \quad (1.170)$$

i. e., no potential, as expected for the dual photon field, χ . The minima for the potential (1.170) is at $\phi = \infty$ ⁹.

It is important to stress some aspects of the previous computation: first of all, the superpotential (1.169) is simply given by the instanton action, with the extra term $\frac{i4\pi\chi}{g^2}$ the analog of a topological θ term in four dimensions. Secondly, the fermions appearing in (1.167), the effective lagrangian, are the ones in the hypermultiplet of the $N=2$ theory. Finally, the superpotential for Φ is defined on a flat direction.

The generalization of the previous picture to the four dimensional case is certainly not straightforward, as in that case we have not flat directions, and the effective lagrangian can not be written in terms of chiral superfields containing the gluino, but the gluino-gluino pair.

1.9.1 A Toron Computation.

A direct way to obtain $\langle \lambda\lambda \rangle$ condensates in four dimensional $N=1$ Yang-Mills using self dual gauge configurations, with Pontryagin number $\frac{1}{N}$ ¹⁰ [34]. In subsection 1.5.2 we have described these configurations. The main point in using these torons is that the number of fermionic zero modes automatically reduces to two, which we can identify with the two triplets defined by supersymmetry transformations of one instanton configurations. We will perform the computation in a box, sending at the end its size to infinity. The size of the box is the size of the toron, but we will avoid the dilatation zero mode and the two triplet zero modes defined by superconformal transformations. The toron measure now becomes, simply,

$$\int d^4z \quad (1.171)$$

⁹The reader might be slightly surprised concerning potential (1.170) for the Higgs field. The crucial issue for the correct understanding of this potential requires noticing that the $N=2$ three dimensional theory has been obtained through dimensional reduction of $N=1$ four dimensional Yang-Mills, which contains a flat direction (in next chapter we will define these flat directions more precisely, as Coulomb branches of moduli of vacua).

¹⁰It should already be noticed that topological configurations directly contributing to $\langle \lambda\lambda \rangle$ are most probably the relevant configurations for confinement, as $\langle \lambda\lambda \rangle$ was derived through a cluster argument assuming the existence of a mass gap.

for the translation collective coordinate. Now, we have a power of μ , given by the four translation zero modes, and two fermionic zero modes,

$$\mu^3 \exp -\frac{8\pi^2}{g(\mu)^2 N}, \quad (1.172)$$

where we have included the toron action $\frac{8\pi^2}{g^2 N}$. Notice that (1.172) is simply Λ^3 . Now, we integrate z over the box of size L . The two fermionic zero modes are obtained by the supersymmetry transformation (1.142) over the toron configuration (1.93), which means that each fermionic zero mode behaves as $\frac{1}{L^2}$, and therefore no powers of L should be included in the measure. The final result is

$$< \lambda \lambda > \sim \text{constant} \Lambda^3 e^{2\pi i e/N}, \quad e = 0, 1, \dots, N-1 \quad (1.173)$$

in agreement with the cluster derivation. How should this result be interpreted? First of all, the expectation value (1.173) corresponds to the amplitude

$$\begin{aligned} & < \vec{e}, \vec{m} = (0, 0, 1) | \lambda \lambda | \vec{e}, \vec{m} = (0, 0, 1) > = \\ & < \vec{k} + (0, 0, 1), \vec{m} = (0, 0, 1) | \lambda \lambda | \vec{k}, \vec{m} = (0, 0, 1) > e^{2\pi i \frac{\vec{e} \cdot (0, 0, 1)}{N}}. \end{aligned} \quad (1.174)$$

Then, the e in (1.173) is e_3 , and the different values in (1.173) correspond to the set of N different vacua described in subsection 1.5.1.

Notice that a change $\theta \rightarrow \theta + 2\pi$ in equation (1.165) produces a change

$$< \lambda \lambda >_{\theta \rightarrow \theta + 2\pi} = < \lambda \lambda >_{\theta} e^{2\pi i/N}, \quad (1.175)$$

i. e., a $\mathbf{Z}(N)$ rotation. In other words, $\theta \rightarrow \theta + 2\pi$ exchanges the different vacua. Let us now try the same argument for (1.174). Using (1.74), we observe that

$$< \lambda \lambda > \sim \Lambda^3 e^{2\pi i e_{eff}/N} = \Lambda^3 e^{2\pi i e/N} e^{i\theta/N}, \quad (1.176)$$

in agreement with (1.165). So, under $\theta \rightarrow \theta + 2\pi$, we go, using (1.74), from $\vec{e}_{eff}$ to $\vec{e}_{eff} + \vec{m}$. Notice that for the toron computation we are using $\vec{m} = \mathbf{1}$.

2 Chapter II

2.1 Moduli of Vacua.

In this part of the lectures, we will consider gauge theories possessing potentials with flat directions. The existence of flat potentials will motivate the definition of moduli of vacua, which we will understand as the quotient manifold

$$\mathcal{M} = \mathcal{V}/\mathcal{G}, \quad (2.1)$$

obtained from the modding of the vacuum manifold $\mathcal{V}$ by gauge symmetries. In the first chapter, an example has already been discussed, namely three dimensional $N=2$ Yang-Mills, defined as dimensional reduction of $N=1$ Yang-Mills in four dimensions. Denoting by $\phi^a = A_4^a$ the fourth component of the gauge field, the dimensionally reduced lagrangian is

$$\mathcal{L} = -\frac{1}{4}F_{ij}^a F^{aij} + \frac{1}{2}\mathcal{D}_i\phi^a\mathcal{D}^i\phi^a + i\bar{\chi}^a\gamma_i\mathcal{D}^i\chi + if_{abc}\bar{\chi}^b\chi^c\phi^a. \quad (2.2)$$

This is the Yang-Mills-Higgs lagrangian in the Prasad-Sommerfeld limit $V(\phi) = 0$. At tree level, the vacuum expectation value for the field ϕ is undetermined; therefore, at the classical level we can define a moduli of (real) dimension one, parametrizing the different values of $\langle \phi \rangle$. As we already know, in addition to the scalar ϕ we have yet another scalar field, χ , the dual photon field. No potential can be defined for χ , neither classically nor quantum mechanically. If we took into account the action of the Weyl group, $\phi \rightarrow -\phi$, $\chi \rightarrow -\chi$, the classical moduli manifold should be

$$\mathbf{R} \times S^1/\mathbf{Z}_2. \quad (2.3)$$

The fields ϕ and χ can be combined into a complex scalar, $\Phi = \phi + i\chi$. As discussed in chapter I, instantons generate a superpotential of type $e^{-\Phi}$, which induces a potential for the ϕ fields with its minimum at ∞ . This potential lifts the classical degeneracy of vacua. The vacuum expectation value of χ still remains undetermined, but can be changed by just shifting the coefficient of the topological term. The physics of this first example is what we expect from physical grounds: quantum effects breaking the classical vacuum degeneracy. However, there are cases where the amount of supersymmetry prevents, a priori, the generation of superpotential terms; it is in these cases, where we should be able to define the most general concept of quantum moduli [35, 36], where quantum effects will modify the topology and geometry of the classical moduli manifold.

2.2 $N=4$ Three Dimensional Yang-Mills.

$N=4$ three dimensional Yang-Mills will be defined through dimensional reduction of $N=1$ six dimensional Yang-Mills [37]. The three real scalars ϕ_i^a , with $i = 1, 2, 3$, corresponding to the $\mu = 3, 4, 5$ vector components of A_μ^a , are in the adjoint representation of the gauge group, and will transform as a vector with respect to the $SO(3)_R$ group of rotations in the 3, 4, 5-directions. The fermions in the model will transform, with respect to the $SU(2)_R$ double cover, as doublets, i. e., as spin one half particles. If we now consider the $SU(2)_E$ rotation group of euclidean space, $\mathbf{R}^3$, then fermions transform again as doublets, while scalars, ϕ_i^a , are singlets. By dimensional reduction, we get the following potential for the ϕ_i :

$$V(\phi) = \frac{1}{4g^2} \sum_{i < j} \text{tr}[\phi_i, \phi_j]^2, \quad (2.4)$$

where we have used a six dimensional lagrangian $-\frac{1}{4g^2} F_{\mu\nu}^a F^{a\mu\nu}$. Obviously, the potential (2.4) possesses flat directions, obtained as those whose ϕ_i fields are in the Cartan subalgebra of the gauge group. For $SU(N)$, we will get

$$\phi_i = \begin{pmatrix} a_i^1 & & \\ & \ddots & \\ & & a_i^N \end{pmatrix}, \quad (2.5)$$

with $\sum_{j=1}^N a_i^j = 0$, so that $3(N-1)$ parameters characterize a point in the flat directions of (2.4). In the general case of a gauge group of rank r , $3r$ coordinates will be required. A vacuum expectation value like (2.5) breaks $SU(N)$ to $U(1)^{N-1}$. As each $U(1)$ has associated a dual photon field, χ_j , with $j = 1, \dots, N-1$, the classical moduli has a total dimension equal to $3r + r = 4r$. The simplest case of $SU(2)$, corresponds to a four dimensional moduli, which classically is

$$\mathcal{M} = (\mathbf{R}^3 \times S^1)/\mathbf{Z}_2. \quad (2.6)$$

We have quotiented by the $\mathbf{Z}_2$ Weyl action, and S^1 parametrizes the expectation value of the dual photon field. The group $SU(2)_R$ acts on the $\mathbf{R}^3$ piece of (2.6). In the same sense as for the $N=2$ theory considered in chapter I, the $N=4$ model possesses instanton solutions to the Bogomolny equations which are simply the dimensional reduction to three dimensional euclidean space of four dimensional self dual Yang-Mills equations.

These Bogomolny-Prasad-Sommerfeld instantons involve only one scalar field, ϕ_i , out of the three available (we will therefore choose one of them, say ϕ_3), and satisfy the equation

$$F = *\mathcal{D}\phi_3. \quad (2.7)$$

Once we choose a particular vacuum expectation value for the ϕ_i fields, we break $SU(2)_R$ to an $U(1)_R$ subgroup. In particular, we can choose $\phi_1 = \phi_2 = 0$, and ϕ_3 different from zero, with ϕ_3 the field used in the construction of the BPS instanton. The remaining $U(1)_R$ stands for rotations around the ϕ_3 direction.

Now, as discussed above, the BPS instanton can induce effective fermionic vertices. We first will consider the case of no extra hypermultiplets. In this case, we have four fermionic zero modes, corresponding to the four supersymmetry transformations that do not annihilate the instanton solution. In addition, we know, from the results in chapter I, that the effective vertex should behave like

$$\psi\psi\psi\psi \exp -(I + i\chi), \quad (2.8)$$

with χ the dual photon field, and I the classical instanton action, behaving like $\frac{\phi_3}{g^2}$. The term (2.7) breaks the $U(1)_R$ symmetry, as the ψ fermions transform under the $U(1)_R$ subgroup of $SU(2)_R$ like

$$\psi \rightarrow e^{i\theta/2}\psi. \quad (2.9)$$

However, the dual photon field plays the role of a Goldstone boson, to compensate this anomalous behaviour; χ transforms under $U(1)_R$ as

$$\chi \rightarrow \chi - 4 \left(\frac{\theta}{2} \right). \quad (2.10)$$

This is a quantum effect, that will have important consequences on the topology of the classical moduli. In fact, the $U(1)_R$ is not acting only on the $\mathbf{R}^3$ part of $\mathcal{M}$, but also (as expressed by (2.10)) on the S^1 piece.

The topological meaning of (2.10) can be better understood if we work [37] on the boundary of $\mathbf{R}^3 \times S^1$, namely $S^2 \times S^1$. In this region at infinity, (2.10) will define a non trivial S^1 bundle on S^2 . In order to see the way this is working, it is useful to use the spinorial notation. Henceforth, let u_1 and u_2 be two complex variables, satisfying

$$|u_1|^2 + |u_2|^2 = 1. \quad (2.11)$$

This defines the sphere S^3 . Parametrizing points in S^2 by a vector $\vec{n}$, defined as follows:

$$\vec{n} = \bar{u}\vec{\sigma}u, \quad (2.12)$$

with $\vec{\sigma}$ the Pauli matrices, and using (2.12), we can define a projection,

$$\Phi : S^3 \rightarrow S^2, \quad (2.13)$$

associating to u , in S^3 , a point $\vec{n}$ in S^2 . The fiber of the projection (2.13) is S^1 . In fact, u_α and $e^{i\theta}u_\alpha$ yield the same value of $\vec{n}$; therefore, we conclude

$$S^2 = S^3/U(1), \quad (2.14)$$

with the $U(1)$ action

$$u_\alpha \rightarrow e^{i\theta}u_\alpha. \quad (2.15)$$

A $U(1)$ rotation around the point $\vec{n}$ preserves $\vec{n}$, but changes u as

$$u_\alpha \rightarrow e^{i\theta/2}u_\alpha, \quad (2.16)$$

where now θ in (2.16) is the $U(1)_R$ rotation angle. Moreover, it also changes χ to $\chi - 4\left(\frac{\theta}{2}\right)$. So, the infinity of our quantum moduli looks like

$$(S^3 \times S^1)/U(1), \quad (2.17)$$

with the $U(1)$ action

$$u_\alpha \rightarrow e^{i\theta}u_\alpha, \quad \chi \rightarrow \chi - 4\theta. \quad (2.18)$$

The space (2.17) is the well known Lens space L_{-4} . Generically, L_s spaces can be defined through (2.17), with the $U(1)$ action defined by

$$\chi \rightarrow \chi + s\theta. \quad (2.19)$$

The spaces L_s can also be defined as

$$L_s = S^3/\mathbf{Z}_s, \quad (2.20)$$

with the $\mathbf{Z}_s$ action

$$\beta : u_\alpha \rightarrow e^{2\pi i/s}u_\alpha. \quad (2.21)$$

To finish the construction of the infinity of the quantum moduli, we need to include the Weyl action, $\mathbf{Z}_2$, so that we get

$$L_s/\mathbf{Z}_2 \quad (2.22)$$

or, equivalently, S^3/Γ_s , where Γ_s is the group generated by β given in (2.21), and the Weyl action

$$\alpha : (u_1, u_2) \rightarrow (\bar{u}_2, -u_1), \quad (2.23)$$

which reproduces $\vec{n} \rightarrow -\vec{n}$. The relations defining the group Γ_s are

$$\alpha^2 = \beta^s = 1, \quad \alpha\beta = \beta^{-1}\alpha. \quad (2.24)$$

Moreover, for $s = 2k$, Γ_s is the dihedral group D_{2k} .

Before entering into a more careful analysis of the moduli space for these $N = 4$ theories, let us come back to the discussion on the $SU(2)_R$ symmetry, and the physical origin of the parameter s . In order to do so, we will first compactify six dimensional super Yang-Mills down to four dimensions. The resulting $N = 2$ supersymmetric Yang-Mills theory has only two scalar fields, ϕ_1 and ϕ_2 . The rotation symmetry in the compactified $(4, 5)$ -plane becomes now a $U(1)$ symmetry for the ϕ_i fields, which is the well known $U(1)_R$ symmetry of $N = 2$ supersymmetric four dimensional Yang-Mills. As in the case for the $N = 4$ theory in three dimensions, instantons in four dimensions generate effective fermionic vertices which break this $U(1)_R$ symmetry. Following the steps of the instanton computations presented in chapter I, we easily discover a breakdown of $U(1)_R$ to $\mathbf{Z}_8$. In the case of $N = 2$ in four dimensions, the potential obtained from dimensional reduction is

$$V(\phi) = \frac{1}{4g^2} \text{tr}[\phi_1, \phi_2]^2, \quad (2.25)$$

with the flat direction, for $SU(2)$,

$$\phi_i = \begin{pmatrix} a_i & \\ & -a_i \end{pmatrix}. \quad (2.26)$$

The moduli coordinate is therefore $u = \text{tr}\phi^2$, where we now define ϕ as a complex field. The $U(1)_R$ transformation of ϕ is given by

$$\phi \rightarrow e^{2i\alpha}\phi, \quad (2.27)$$

so that u is $\mathbf{Z}_4$ invariant, and the $\mathbf{Z}_8$ symmetry acts as $\mathbf{Z}_2$ on the moduli space.

The difference with the three dimensional case is that now the instanton contribution does not contain any dual photon field, that can play the role of a Goldstone boson. From the three dimensional point of view, the $U(1)_R$ symmetry of the four dimensional theory is the rotation group acting on fields ϕ_1 and ϕ_2 , and can therefore be identified with the $U(1)_R$ part of $SU(2)_R$ fixing the ϕ_3 direction. We should wonder about the relationship between the effect observed in three dimensions, and the breakdown of $U(1)_R$ in four dimensions. A qualitative answer is simple to obtain. The breakdown of $U(1)_R$ in four dimensions can be studied in the weak coupling limit (corresponding to $u \rightarrow \infty$), since

the theory is asymptotically free, using instantons. Then, we get an effective vertex of the type

$$\langle \psi\psi\psi\psi \rangle_0 e^{i\theta} = \langle \psi\psi\psi\psi \rangle_\theta. \quad (2.28)$$

We have only four fermionic zero modes, since for $u \neq 0$ we break the superconformal invariance of the instanton. It is clear, from (2.28), that a $U(1)_R$ transformation is equivalent to the change

$$\theta \rightarrow \theta - 4\alpha, \quad (2.29)$$

with α the $U(1)_R$ parameter. Now, this change in θ is, in fact, the perfect analog of transformation rule (2.10), for the dual photon field. This should not come as a surprise; in fact, the four dimensional topological term

$$\frac{i\theta}{32\pi^2} F * F \quad (2.30)$$

produces, by dimensional reduction, the three dimensional topological term

$$\frac{i\theta}{32\pi^2} \epsilon_{ijk} F_{jk} \partial_i \phi_3. \quad (2.31)$$

This is precisely the type of coupling of the dual photon, in three dimensions, with the topological charge, and thus we again recover the result of section 1.7.

From the previous discussion, we can discover something else, specially interesting from a physical point of view. The transformation law of χ was derived counting instanton fermionic zero modes; however, the effect we are describing is a pure perturbative one loop effect, as is the $U(1)_R$ anomaly in four dimensions. Consider the wilsonian [38, 39] effective coupling constant for the $N=2$ theory, without hypermultiplets. Recall that in the wilsonian approach [40], the effective coupling constant is defined in terms of the scale we use to integrate out fluctuations with wave length smaller than that scale (this is the equivalent to the Kadanoff approach for lattice models). In a Higgs model, the natural scale is the vacuum expectation value of the Higgs field. Using the above notation, the wilsonian coupling constant in the four dimensional model is $\frac{1}{g(u)^2}$, with u the moduli parameter defined by $\text{tr}\phi^2$. Now, let us write the lagrangian as follows:

$$\mathcal{L} = \frac{1}{64\pi} \text{Im} \int \tau (F + i * F)^2, \quad (2.32)$$

with τ defined by

$$\tau \equiv \frac{i8\pi}{g^2} + \frac{\theta}{\pi}. \quad (2.33)$$

Using $F^2 = - * F^2$ we get, from (2.32), the standard lagrangian in Minkowski space,

$$\mathcal{L} = \int \frac{1}{4g^2} FF + \frac{\theta}{32\pi^2} F * F. \quad (2.34)$$

Now, we use the one loop effective beta function for the theory,

$$\frac{8\pi}{g(u)^2} = \frac{2}{\pi} \ln u + \mathcal{O}(1). \quad (2.35)$$

In general, if we add n hypermultiplets, we get, for the four dimensional theory,

$$\frac{8\pi}{g(u)^2} = \frac{4-n}{2\pi} \ln u + \mathcal{O}(1), \quad (2.36)$$

recovering the well known result for $N=2$ supersymmetric $SU(2)$ gauge theories in four dimensions, of finiteness of the theory when $n = 4$, and infrared freedom for $n > 4$. For $n < 4$ the theory is asymptotically free, so that the perturbative computation (2.36) is only valid at small distances, for u in the asymptotic infinity.

Now, let us perform a rotation on u ,

$$u \rightarrow e^{2\pi i} u \quad (2.37)$$

From (2.35) we get, for $n = 0$,

$$\frac{8\pi}{g(u)^2} \rightarrow \frac{8\pi}{g(u)^2} + 4i, \quad (2.38)$$

so, using (2.33), we get

$$\theta \rightarrow \theta - 4\pi, \quad (2.39)$$

in perfect agreement with equation (2.29). Thus, we observe that the s term (at least for the case without hypermultiplets) that we have discovered above using three dimensional instanton effects, is exactly given by the one loop effect of the four dimensional $N = 2$ theory. But what about higher order loop effects? As the argument we have presented is nothing but the non renormalization theorem [39] in supersymmetric theories, the $U(1)_R$ action on the wilsonian scale u forces the renormalization of the coupling constant to be consistent with the $U(1)$ anomalous behaviour of the lagrangian, which is determined by the Adler-Bardeen theorem [41] to be exact at one loop. What happens as we include hypermultiplets? First of all, and from the point of view of the three dimensional theory, the instanton effect will now be a vertex of type,

$$\psi\psi\psi\psi \prod \chi e^{-(I+i\chi)}, \quad (2.40)$$

with the $2N_f$ fermionic zero modes appearing as a consequence of Callias index theorem [25], (1.136), for $j = 1/2$ and $\{m\} = -1/2$. From (2.40), we get $s = -4 + 2N_f$, which means a dihedral group Γ_s of type D_{2N_f-4} or, equivalently, a Dynkin diagram of type D_{N_f} . Notice that in deriving this diagram we have already taken into account the Weyl action, $\mathbf{Z}_2$.

The connection between the dihedral group, characterizing the moduli of the three dimensional $N = 4$ theory, and the beta function for the four dimensional $N = 2$ theory can be put on more solid geometrical grounds. The idea is simple. Let us work with the $N = 2$ four dimensional theory on $\mathbf{R}^3 \times S^1$, instead of on euclidean space, $\mathbf{R}^4$. The massless fields, from the three dimensional point of view, contain the fourth component of the photon in four dimensions, and the standard dual photon χ in three dimensions. Requiring, as in a Kaluza-Klein compactification, all fields to be independent of x_4 , we still have residual gauge transformations of the type

$$A_4 \rightarrow A_4 + \partial_4 \Lambda(x_4), \quad (2.41)$$

with $\Lambda(x_4) = \frac{x_4 b}{\pi R}$, with b an angular variable, $b \in [0, 2\pi]$. This is equivalent to saying that we have non trivial Wilson lines in the S^1 direction, if the gauge field is in $U(1)$, as happens to be the case for a generic value of u ¹¹. Now, at each point u in the four dimensional $N = 2$ moduli, we have a two torus E_u , parametrized by the dual photon field χ , and the field b . This E_u is obtained from the S^1 associated with χ , and the S^1 associated with b . Its volume, in units defined by the three dimensional coupling constant, is of order $\frac{1}{R}$,

$$\text{Vol } E \sim \frac{1}{R}. \quad (2.42)$$

In fact, the volume is $\frac{1}{R} \cdot g_3^2$, where g_3^2 , the three dimensional coupling constant, is the size of the S^1 associated to the dual photon (notice that the coupling constant g_3^2 , in three dimensions, has units of inverse length). Equation (2.42) shows how, in the four dimensional limit, E_u goes to zero volume. Now, we have a picture of the theory in $\mathbf{R}^3 \times S_R^1$, if we keep R finite, namely that of an elliptic fibration over the u -plane, parametrizing the vacuum expectation values of the $N = 2$ four dimensional theory.

If we keep ourselves at one particular point u , the torus E_u should be the target space for the effective lagrangian for the fields b and χ . There is a simple way to derive this

¹¹We are not imposing, to the magnetic flux through the S^1 , to be topologically stable in the sense of $\Pi_1(U(1)) \simeq \mathbf{Z}$. The crucial point is that the value of b at this point of the game is completely undetermined, and in that sense is a moduli parameter.

lagrangian by means of a general procedure, called dualization, that we will now describe. To show the steps to follow, we will consider the four dimensional lagrangian (2.32). In order to add a dual photon field, let us say A_D^μ , we must couple A_D^μ to the monopole charge,

$$\epsilon^{0ijk}\partial_i F_{jk} = 4\pi\delta^{(3)}(x). \quad (2.43)$$

Thus, we add a term

$$\frac{1}{4}\int A_D^\mu \epsilon_{\mu\nu\rho\sigma}\partial^\nu F^{\rho\sigma} \equiv \frac{1}{4\pi}\int *F_D F. \quad (2.44)$$

Using the same notation as in (2.32), we get

$$\frac{1}{4\pi}\int *F_D F = \frac{1}{8\pi}\text{Re}\int (*F_D - iF_D)(F + i*F), \quad (2.45)$$

so that our lagrangian is

$$\mathcal{L} = \frac{1}{64\pi}\text{Im}\int \tau(F + i*F)^2 + \frac{1}{8\pi}\text{Re}\int (*F_D - iF_D)(F + i*F). \quad (2.46)$$

After gaussian integration, we finally get

$$\mathcal{L} = \frac{1}{64\pi}\text{Im}\int \left(\frac{-1}{\tau}\right) (*F_D - iF_D)^2, \quad (2.47)$$

i. e., lagrangian (2.32), with τ replaced by $\frac{-1}{\tau}$. The reader should take into account that these gaussian integrations are rather formal manipulations. Now, we use the same trick to get an effective lagrangian for the fields b and χ . Start with the four dimensional lagrangian,

$$\mathcal{L} = \int \frac{1}{4g^2} F F + \frac{i\theta}{32\pi^2} F * F, \quad (2.48)$$

where we now work in euclidean space. In three dimensions, using $d^4x = 2\pi R d^3x$, we get

$$\mathcal{L} = \int d^3x \frac{1}{\pi R g^2} |db|^2 + \frac{\pi R}{2g^2} F_{ij}^2 + \frac{i\theta}{8\pi^2} \epsilon^{ijk} F_{jk} \partial_k b. \quad (2.49)$$

Now, as we did before, we couple the dual photon field to the monopole charge,

$$\partial_i H^i = 4\pi\delta^{(3)}(x), \quad (2.50)$$

with $H^i \equiv \epsilon^{ijk} F_{jk}$, to get a term

$$\frac{i}{8\pi} \epsilon^{ijk} F_{jk} \partial_i \chi, \quad (2.51)$$

so that we can perform a gaussian integration,

$$\mathcal{L} = \int d^3x \frac{1}{\pi R g^2} |db|^2 + \frac{g^2}{\pi R (8\pi)^2} |d\chi - \frac{\theta}{\pi} db|^2. \quad (2.52)$$

What we get is precisely a target space for χ and b fields, which is the torus of moduli τ given by (2.33). Observe that the complex structure of the torus E_u is given in terms of the four dimensional coupling constant g [37], and the four dimensional θ -parameter, while its volume, (2.42), depends on the three dimensional coupling constant g_3 , that acts as unit. When we go to the four dimensional $R \rightarrow \infty$ limit, this volume becomes zero, but the complex structure remains the same. The fact that the complex structure of E_u is given by the four dimensional effective coupling will make more transparent the meaning of equation (2.36). In fact, the monodromy around $u = \infty$ for (2.36) is given by a matrix

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} -1 & -n+4 \\ 0 & -1 \end{pmatrix}, \quad (2.53)$$

with τ transforming as

$$\tau \rightarrow \frac{a\tau + b}{c\tau + d}, \quad (2.54)$$

so that for $n = 0$ we get transformation (2.39). Next, we will see that transformation (2.53) is precisely what we need, in order to match the dihedral group characterization of the $N=4$ three dimensional moduli space; however, in order to do that we need a few words on Atiyah-Hitchin spaces [42].

2.3 Atiyah-Hitchin Spaces.

Atiyah-Hitchin spaces appear in the study of moduli spaces for static multi-monopole configurations. Static solutions are defined by the BPS equations, (2.7), which are simply the dimensional reduction to $\mathbf{R}^3$ of euclidean self-dual equations for instantons. Next, we simply summarize some of the relevant results on Atiyah-Hitchin spaces for our problem (we refer the interested reader to the book by M. Atiyah and N. Hitchin, [42]). First of all, the Atiyah-Hitchin spaces are hyperkähler manifolds of dimension $4r$, on which a rotation $SO(3)$ is acting in a specific way. This is part of what we need to define the moduli space of $N=4$ three dimensional Yang-Mills theory for gauge group of rank r . In fact, in order to define $N=4$ supersymmetry on this space, interpreted as a σ -model target space of the low energy effective lagrangian, we have to require hyperkähler structure. Recall here that hyperkähler simply means that we have three different complex structures, I ,

J and K , and therefore three different Kähler forms, ω_i , ω_j and ω_k , which are closed. Following the notation used by Atiyah and Hitchin, we define N_k as the moduli space of a k monopole configuration. The dimension of N_k is $4k - 1$, so for $k = 1$ we get dimension 3, corresponding to the position of the monopole center. If we mod out by the translation of the center of mass, we get the space

$$M_k^0 = N_k / \mathbf{R}^3, \quad (2.55)$$

of dimension $4(k - 1)$. For two monopoles, we get $\dim M_2^0 = 4$. Now, the spaces M_k^0 are generically non simply connected,

$$\Pi_1(M_k^0) \simeq \mathbf{Z}_k, \quad (2.56)$$

so we can define its k -fold covering $\tilde{M}_k^0$. The known results, for $k = 2$, are the following: the spaces M_2^0 and $\tilde{M}_2^0$ are, at infinity, respectively of type $L_{-4}/\mathbf{Z}_2$, and $L_{-2}/\mathbf{Z}_2$, which strongly indicates that M_2^0 is a good candidate for the moduli of the $N_f = 0$ case, and $\tilde{M}_2^0$ is the adequate for the $N_f = 1$ case. Moreover, the spaces $\tilde{M}_2^0$ can be represented by a surface in $\mathbf{C}^3$, defined by

$$y^2 = x^2 v + 1. \quad (2.57)$$

The space $M_2^0 = \tilde{M}_2^0 / \mathbf{Z}_2$, can be obtained using variables $\chi = x^2$ and $y = x$, so that we get

$$y^2 = \chi^2 v + \chi. \quad (2.58)$$

The spaces $\tilde{M}_2^0$, defined by (2.55), can be interpreted as a limit of the family of spaces

$$y^2 = x^2 v + v^l, \quad (2.59)$$

where l should, in our case, be identified with $N_f - 1$. Surfaces (2.59) are well known in singularity theory; they give rise to the type of singularities obtained from $\mathbf{C}^3/\Gamma$, with Γ a discrete subgroup of $SO(3)$, and are classified according to the following table [43],

Γ	Name	Singularity
$\mathbf{Z}_n$	A_{n-1}	$v^n + xy = 0$
$\mathbf{D}_{2n}$	D_{n+2}	$vx^2 - v^{n+1} + y^2 = 0$
$\mathbf{T}_{12}$	E_6	$v^4 + x^3 + y^2 = 0$
$\mathbf{O}_{24}$	E_7	$v^3 + vx^3 + y^2 = 0$
$\mathbf{I}_{60}$	E_8	$v^5 + x^3 + y^2 = 0$

As can be seen from this table, the manifold (2.59) corresponds to a D_{n+2} singularity, with $n = N_f - 2$, and dihedral group D_{2N_f-4} , i. e., the group Γ we have discussed in the previous section.

It is important to stress that the type of singularities we are describing in the above table are the so called rational singularities [44]. The geometrical meaning of the associated Dynkin diagram is given by the resolution of the corresponding singularity as the intersection matrix of the irreducible components obtained by blowing up the singularity. In this interpretation, each node of the diagram corresponds to an irreducible component, which is a rational curve X_i , with self dual intersection $X_i.X_i = -2$, and each line to the intersection $X_i.X_j$ between different irreducible components.

In the previous section we have modelled the $N = 4$ moduli space as an elliptic fibration, with fiber E_u of volume $\frac{1}{R}$, and moduli τ given by the coupling constant of the four dimensional $N = 2$ gauge theory. Next, we will try to connect the dihedral group, characterizing the Atiyah-Hitchin space describing the $N = 4$ moduli, with the monodromy at infinity of the elliptic modulus of E_u . But before doing this, we will briefly review Kodaira's theory on elliptic singularities.

2.4 Kodaira's Classification of Elliptic Fibrations.

According to Kodaira's notation [45], we define an elliptic fibration V onto Δ , where Δ will be chosen as a compact Riemann surface. In general, we take Δ to be of genus equal zero. The elliptic fibration,

$$\Phi : V \rightarrow \Delta, \quad (2.60)$$

will be singular at some discrete set of points, a_ρ . The singular fibers, C_{a_ρ} , are given by

$$C_{a_\rho} = \sum n_s \Theta_{\rho s}, \quad (2.61)$$

with $\Theta_{\rho s}$ irreducible curves. According to Kodaira's theorem (see section 4.7 for more details), all possible types of singular curves are of the following types:

•

$$I_{n+1} : C_{a_\rho} = \Theta_0 + \Theta_1 + \cdots + \Theta_n, \quad n+1 \geq 3, \quad (2.62)$$

where Θ_i are non singular rational curves with intersections $(\Theta_0, \Theta_1) = (\Theta_1, \Theta_2) = \cdots = (\Theta_n, \Theta_0) = 1$.

The A_n affine Dynkin diagram can be associated to I_{n+1} . Different cases are

- i) I_0 , with $C_\rho = \Theta_0$ and Θ_0 elliptic and non singular.
- ii) I_1 , with $C_\rho = \Theta_0$ and Θ_0 a rational curve, with one ordinary double point.
- iii) I_2 , with $C_\rho = \Theta_0 + \Theta_1$ and Θ_0 and Θ_1 non singular rational points, with intersection $(\Theta_0, \Theta_1) = p_1 + p_2$, i. e., two points.

Notice that I_1 and I_2 correspond to diagrams A_0 and A_1 , respectively.

- Singularities of type I_{n-4}^* are characterized by

$$I_{n-4}^* : C_\rho = \Theta_0 + \Theta_1 + \Theta_2 + \Theta_3 + 2\Theta_4 + 2\Theta_5 + \cdots + 2\Theta_n, \quad (2.63)$$

with intersections $(\Theta_0, \Theta_4) = (\Theta_1, \Theta_4) = (\Theta_2, \Theta_4) = (\Theta_3, \Theta_4) = (\Theta_4, \Theta_5) = (\Theta_5, \Theta_6) = \cdots = 1$, these singularities correspond to the D_n Dynkin diagram.

- Singularities of type II^* , III^* and IV^* correspond to types E_6 , E_7 and E_8 .

In addition to these singularities, we have also the types

- II : $C_\rho = \Theta_0$, with Θ_0 a rational curve with a cusp.
- III : $C_\rho = \Theta_0 + \Theta_1$, with Θ_0 and Θ_1 non singular rational curves, with intersection $(\Theta_0, \Theta_1) = 2p$.
- IV : $C_\rho = \Theta_0 + \Theta_1 + \Theta_2$, with Θ_0 , Θ_1 and Θ_2 non singular rational curves, with intersections $(\Theta_0, \Theta_1) = (\Theta_1, \Theta_2) = (\Theta_2, \Theta_0) = p$.

In contrast to the singularities described in last section (the rational ones), these singularities are associated to affine Dynkin diagrams. Observe that for all these singularities we have

$$C.C = 0, \quad (2.64)$$

while in the rational case the corresponding maximal cycle satisfies

$$C.C = -2. \quad (2.65)$$

The origin for the affinization of the Dynkin diagram is the elliptic fibration structure. In fact, we can think of a rational singularity of ADE type in a surface, and get the affinization of the Dynkin diagram whenever there is a singular curve passing through the singularity. In the case of an elliptic fibration, this curve is the elliptic fiber itself. So, the extra node in the Dynkin diagram can be interpreted as the elliptic fiber. This can be seen

more clearly as we compute the Picard of the surface. In fact, for the elliptic fibration the contribution to the Picard comes from the fiber, the basis, and the contribution from each singularity. Now, in the contribution to Picard from each singularity, we should not count the extra node, since this has already been taken into account when we count the fiber as an element in the Picard.

The previous discussion is already telling us what happens when we go to the $R = 0$ limit, i. e., to the three dimensional $N = 4$ gauge theory. In this limit, the elliptic fiber E_u becomes of infinite volume, and therefore we can not consider it anymore as a compact torus, i. e., as an elliptic curve. Thus, in this limit the corresponding singularity should become rational, and the Dynkin diagram is not affine.

However, before entering that discussion, let us work out the monodromies for the elliptic fibrations of Kodaira's classification.

We will then define

$$\tau(u) = \frac{\int_{u \times \gamma_1} \varphi(u)}{\int_{u \times \gamma_2} \varphi(u)}, \quad (2.66)$$

with $\varphi(u)$ the holomorphic one form on C_u . From (2.66), it follows that $\tau(u)$ is a holomorphic function of u . Next, we define the elliptic modular function, $j(\tau(u))$, on the upper half plane,

$$j(\tau(u)) = \frac{1728g_2^3}{\Delta}, \quad (2.67)$$

where Δ is the discriminant $\Delta = g_2^3 - 27g_3^2$, with the modular functions

$$\begin{aligned} g_2 &= 60 \sum_{n_1, n_2 \in \mathbf{Z}} \frac{1}{(n_1 + n_2 \tau)^4}, \\ g_3 &= 140 \sum_{n_1, n_2 \in \mathbf{Z}} \frac{1}{(n_1 + n_2 \tau)^6}. \end{aligned} \quad (2.68)$$

Defining $\mathcal{F}(u) \equiv j(w(u))$ as a function of Δ , it turns out to be a meromorphic function. To each pole a_ρ , and each non contractible path γ_ρ in Δ (that is, an element in $\Pi_1(\Delta)$), we want to associate a monodromy matrix, A_ρ ,

$$A_\rho \tau \rightarrow \frac{a\tau + b}{c\tau + d}. \quad (2.69)$$

If a_ρ is a pole of $\mathcal{F}(u)$, of order b_ρ , then it can be proved that A_ρ is of type

$$A_\rho \tau \rightarrow \tau + b_\rho, \quad (2.70)$$

for some b_ρ . The matrix A_ρ , of finite order, $A_\rho^m = 1$, for some m , corresponds to singularities which can be removed. Moreover, if A_ρ is of infinite order, then it is always possible to find numbers p, q, r and s such that

$$\begin{pmatrix} s & -r \\ -q & p \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} p & r \\ q & s \end{pmatrix} = \begin{pmatrix} 1 & b_\rho \\ 0 & 1 \end{pmatrix}, \quad (2.71)$$

with $ps - qr = 1$. Next, we relate matrices A_ρ with the different types of singularities. The classification, according to Kodaira's work, is as shown in the table below.

Matrix	Type of singularity
$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$	I_1
$\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}$	I_b
$\begin{pmatrix} -1 & -b \\ 0 & -1 \end{pmatrix}$	I_b^*
$\begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}$	II
$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$	III
$\begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$	IV

Now, we can compare the monodromy (2.53) with the ones in the table. It corresponds to the one associated with a singularity of type I_b^* , with $b = n - 4$, i. e., a Dynkin diagram of type D_n . In the rational case, this corresponds to a dihedral group D_{2n-4} . In (2.53), n represents the number of flavors, so that we get the dihedral group of the corresponding Atiyah-Hitchin space.

Summarizing, we get that the dihedral group of $N = 4$ in three dimensions is the one associated with the type of elliptic singularity at infinity of the elliptic fibration defined by the $N = 2$ four dimensional theory. In other words, the picture we get is the following: in the $R \rightarrow 0$ three dimensional limit we have, at infinity, a rational singularity of type $\mathbf{C}^3/D_{2N_f-4}$. When we go to the $R \rightarrow \infty$ limit we get, at infinity, an elliptic singularity with Dynkin diagram D_{N_f} . Both types of singularities describe, respectively, one loop effects in three dimensional $N = 4$ and four dimensional $N = 2$.

2.5 The Moduli Space of the Four Dimensional $N=2$ Supersymmetric Yang-Mills Theory. The Seiberg-Witten Solution.

From our previous discussion, we have observed that the complex structure of the moduli space of three dimensional $N=4$ supersymmetric Yang-Mills theory is given by the elliptic fibration on the moduli space of the four dimensional $N=2$ theory, where the elliptic modulus is identified with the effective complexified coupling constant τ , as defined in (2.33). This result will in practice mean that the complete solution to the four dimensional $N=2$ theory can be directly read out from the complex structure of the Atiyah-Hitchin spaces (2.59), with $l = N_f - 1$. In previous sections, we have already done part of this job, comparing the monodromy of τ around $u = \infty$, i. e., in the asymptotic freedom regime, with the dihedral group characterizing the infinity of the three dimensional $N=4$ moduli space. In this section, we will briefly review the Seiberg-Witten solution [35, 36, 46, 47, 48, 49, 50, 51, 52, 53] for four dimensional $N=2$ Yang-Mills theory, and compare the result with the complex structure of Atiyah-Hitchin spaces. Recall that the Atiyah-Hitchin spaces are hyperkähler, and therefore possess three different complex structures. The complex structure determined by the four dimensional $N=2$ solution is one of these complex structures, namely the one where the Atiyah-Hitchin space becomes elliptically fibered. The analysis of Seiberg and Witten was originally based on the following argument: the moduli space parametrized by u should be compactified to a sphere (we will first of all consider the $N_f = 0$ case, for $SU(2)$ gauge group). According to Kodaira's notation, Δ is taken to be of genus equal zero. Next, the behaviour of τ at $u = \infty$ is directly obtained from the one loop beta function (see equation (2.36)); this leads to a monodromy around infinity of the type (2.53). Next, if $\tau(u)$ is a holomorphic function of u , which is clear from the elliptic fibration mathematical point of view (see equation (2.66)), and is a direct consequence of $N=2$ supersymmetry, then the real and imaginary parts are harmonic functions. As the coupling constant is the imaginary part of the complex structure $\tau(u)$, which is on physical grounds always positive, we are dealing with an elliptic fibration, so we already know all possible types of singularities. That some extra singularities should exist, in addition to the one at infinity, is clear from the harmonic properties of $\text{Im}\tau(u)$, and the fact that it is positive, but in principle we do not know how many of them we should expect, and of what type. The answer to this question can not, in principle, be derived from Kodaira's theory. In fact, all what we can obtain from Kodaira's approach, using the adjunction formula, is a relation between the canonical bundle K of the elliptic fibration, the K of the base space, which we can take as $\mathbb{P}^1$, and

the type of singularities,

$$K_V = \Pi^*(K_\Delta + \sum a_i P_i), \quad (2.72)$$

where the a_i , for each type of singularity, are given below.

Singularity	a_i
I_1	$1/12$
I_b^*	$1/2 + b/12$
I_b	$b/12$
II	$1/6$
III	$1/4$
IV	$1/3$
II^*	$5/6$
III^*	$3/4$
IV^*	$2/3$

However, (2.72) is not useful, at this point, since we do not know the V manifold, which is what we are looking for. We will therefore proceed according to physical arguments.

The singularities we are looking for are singularities in the strong coupling regime of the moduli space of the theory, so it is hopeless to try to use a naive perturbative analysis; instead, we can rely on a duality approach. In dual variables (see equation (2.47)), the effective coupling constant behaves like $\frac{-1}{\tau}$, i. e., we have performed an S transformation, with

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}. \quad (2.73)$$

Thinking of $\frac{-1}{\tau}$ as the effective magnetic coupling, τ^{mag} , we can reduce our analysis to looking for perturbative monodromies of type

$$\tau^{mag} \rightarrow \tau^{mag} + b. \quad (2.74)$$

Indeed, we know that any singularity of Kodaira's type is related to a monodromy of type (2.74), up to a unitary transformation, (see equation (2.71)).

Now, and on physical grounds, we can expect a transformation of the type (2.74) as the monodromy singularity for the effective coupling constant of an effective $U(1)$ theory, with b equal to the number of massless hypermultiplets. In fact, the beta function for the $U(1)$ $N=2$ theory, with n hypermultiplets, is given by

$$\tau^{mag}(u) = -\frac{ik}{2\pi} \ln(u), \quad (2.75)$$

with k the number of massless hypermultiplets. This yields the monodromy

$$\begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}, \quad (2.76)$$

or, in Kodaira's notation, a monodromy of type A_{k-1} . Notice that the difference in sign between the type D , and the type A monodromies, reflects that we are obtaining type A for infrared free theories, and type D (that is D_0 , D_1 , D_2 , and D_3) for asymptotically free theories (notice the sign in (2.75)) [54]. Now, we should wonder about the meaning of (2.75). Recall that our analysis relies upon the wilsonian coupling constant, so the meaning of u in (2.75) must be related to the scale in the $U(1)$ theory, i. e. the vacuum expectation value for the scalar field in the photon multiplet or, more properly, in the dual photon multiplet. This vacuum expectation value gives a mass to the hypermultiplets through the standard Yukawa coupling, so the singularity of (2.75) should be expected at $u = 0$, with u proportional to the mass of the hypermultiplet. Fortunately, we do know which hypermultiplet we should consider: the one defined by the monopole of the theory. In fact, we should rewrite (2.75) as

$$\tau^{mag}(u) = -\frac{ik}{2\pi} \ln(M(u)), \quad (2.77)$$

with $M(u)$ the mass of the monopole, and consider (2.77) perturbatively around the point u_0 , where

$$M(u_0) = 0. \quad (2.78)$$

Therefore, we conclude that a singularity of A_0 type will appear whenever the mass of the monopole equals zero. The nature of the point u_0 is quite clear from a physical point of view: the magnetic effective coupling constant is zero, as can be seen from (2.77), so that the dual electric coupling should become infinity. But the point where the coupling constant is infinity is by definition the scale Λ of the theory; then, $u_0 = \Lambda$.

Now, it remains to discover how many singularities of A_0 type are there. In principle, a single point where the monopole becomes massless should be expected (the $u_0 = \Lambda$ point); however, as mentioned in section 2.2, the $U(1)_R$ symmetry is acting on the moduli space as a $\mathbf{Z}_2$ transformation. Therefore, in order to implement this symmetry, an extra singularity of A_0 type must exist. The simplest solution for the $N_f = 0$ theory, with $SU(2)$ gauge group, corresponds to an elliptic fibration over $\mathbb{P}^1$, the compactified u -plane, with three singular points, of type

$$D_0; A_0, A_0, \quad (2.79)$$

with D_0 the singularity at infinity, and the two A_0 singularities at the points $\pm\Lambda$, with Λ the scale of the theory.

What about the inclusion of flavors? In this case, we know that D_0 in (2.79) is replaced by D_{N_f} . The case $N_f = 2$ should be clear, as D_2 is equivalent to two A_1 singularities and therefore, we should expect

$$D_2; A_1, A_1. \quad (2.80)$$

The singularities of A_1 type indicate that two hypermultiplets become massless. Another simple case is that with $N_f = 4$, where there is a trivial monodromy D_4 , which is now the monodromy around the origin. The two other cases of asymptotically free theories can be obtained through decoupling arguments, and taking into account the residual $U(1)_R$ symmetry. The results are [54]

$$\begin{aligned} D_1 & ; A_0, A_0, A_0, \\ D_3 & ; A_0, A_3. \end{aligned} \quad (2.81)$$

Now, with these elliptic fibrations, we should consider the complex structure. As we know from Kodaira's argument for the $N_f = 0$ case, the A_0 singularities correspond to a rational curve with a double singular point; as we know that this double singularity appears at $u = \pm\Lambda$, the simplest guess for the corresponding complex structure is, with $\Lambda = 1$,

$$y^2 = x^3 - x^2u + x. \quad (2.82)$$

The curve (2.83), for generic u , does not have singular points. Recall that for a curve defined by $f(x, y; u) = 0$, the singular points are those such that

$$\begin{aligned} F & = 0, \\ F_x = F_y & = 0, \end{aligned} \quad (2.83)$$

with F_x and F_y the derivatives with respect to x and y , respectively. The genus of the curve can be obtained using Riemann's theorem,

$$g = \frac{(n-1)(n-2)}{2} - \sum_p \frac{r_p(r_p-1)}{2}, \quad (2.84)$$

where the sum is over singular points, r_p is the order of the singularity, and n in (2.84) is the degree of the polynomial F , defining the curve. So, for generic u , we get, for (2.82), $g = 1$.

Now, for $u = \pm 2$, we have a singular point satisfying (2.83), namely

$$\begin{aligned} y &= 0, \\ x &= \frac{u}{2}. \end{aligned} \tag{2.85}$$

This is a double point and therefore, using (2.84), we get $g = 0$. From Kodaira's classification, we know that at this points we get two singularities of A_0 type. Notice also that at the origin, $u = 0$, we have the curve $y^2 = x^3 + x$, which is of genus one, since there are no singular points. Moreover, if we take $\Lambda = 0$, we get the curve

$$y^2 = x^3 - x^2u. \tag{2.86}$$

This curve has a double point at $x = y = 0$ for generic u . Using (2.84), we now get genus equal zero. Thus, the curve (2.82) satisfies all the properties derived above.

The curve (2.82) has a point at $x = y = \infty$. In order to compactify the curve, we must add the point at infinity. This can be done going to the projective curve

$$zy^2 = x^3 - zx^2u + z^2x. \tag{2.87}$$

The region at infinity of this curve is defined by $z = 0$. The curve, in the three dimensional $R \rightarrow 0$ limit, can be described by (2.82), but with $\text{Vol}(E_u) = \infty$. Next, we will see that this limit is equivalent to deleting the points at infinity of (2.87), i. e., the points with $z = 0$. In fact, for $z \neq 0$ we can define a new variable,

$$v = x - zu, \tag{2.88}$$

and write (2.87) as

$$zy^2 = x^2v + z^2x. \tag{2.89}$$

We can interpret (2.89) as defining a surface in the projective space $\mathbb{P}^3$, but (2.89) is in fact the Atiyah-Hitchin space in homogeneous coordinates. Thus, we conclude that the $R \rightarrow 0$ limit is equivalent to deleting the points at infinity of the curves E_u defined by (2.82).

We can see this phenomena in a different way as follows. The representation (2.82) of the Atiyah-Hitchin space is as an elliptic fibration, so that we have selected one complex structure. However, we can yet rotate in the space of complex structures, preserving the one selected by the elliptic fibration. This defines a $U(1)$ action. This $U(1)$ action must act on E_u ; however, this is impossible if E_u is a compact torus. But when we delete the

point at infinity, and pass to the projective curve (2.89), we have a well defined $U(1)$ action [37],

$$\begin{aligned} x &\rightarrow \lambda^2 x, \\ y &\rightarrow \lambda y, \\ v &\rightarrow \lambda^{-2} v. \end{aligned} \tag{2.90}$$

Only a $\mathbf{Z}_2$ subgroup of this action survives on u :

$$u \rightarrow \lambda^2 x - \lambda^{-2} v \equiv \hat{\lambda} u, \tag{2.91}$$

which means

$$\lambda^2 = \lambda^{-2} = \hat{\lambda}, \tag{2.92}$$

i. e., $\lambda^4 = 1$ or $\hat{\lambda}^2 = 1$. This $\mathbf{Z}_2$ action moves $u \rightarrow -u$, and is the only part of the $U(1)$ action surviving when we work in the four dimensional limit. More simply, at $z = 0$, i. e., at infinity, in the projective sense, $v = x$, and we get $\lambda^2 = \lambda^{-2}$, and the $\mathbf{Z}_4$ symmetry of (2.82) becomes

$$\begin{aligned} y &\rightarrow iy, \\ x &\rightarrow -x, \\ u &\rightarrow -u. \end{aligned} \tag{2.93}$$

Notice also the relation between Λ and the breaking of $U(1)$. In fact, for $\Lambda = 0$ we have

$$y^2 = x^3 - x^2 u, \tag{2.94}$$

which is invariant under

$$\begin{aligned} y &\rightarrow \lambda^3 y, \\ x &\rightarrow \lambda^2 x, \\ u &\rightarrow \lambda^2 u. \end{aligned} \tag{2.95}$$

2.6 Effective Superpotentials.

Maybe the most spectacular result derived from the Seiberg-Witten solution to $N = 2$ supersymmetric theories is the first dynamical proof of electric confinement. In order to properly understand this proof, we need first to go through the recent history of confinement. The simplest physical picture of confinement is that of dual BCS superconductivity

theory [23, 55, 56]. In that picture, a confining vacua is to be represented as the dual of the standard superconducting vacua, which is characterized by the condensation of Cooper pairs. In ordinary superconductivity we find, under the name of Meisner effect, the mechanism for magnetic confinement. In a superconducting vacua, a monopole-antimonopole pair creates a magnetic flux tube that confines them. The relativistic Landau-Ginzburg description of superconductivity was first introduced by Nielsen and Olesen [57], where vortices in the Higgs phase are interpreted as Meisner magnetic flux tubes. The order parameter of the phase is the standard vacuum expectation value of the Higgs field; in this model, simply a scalar coupled to the $U(1)$ electric-magnetic field. The confined monopoles would be $U(1)$ Dirac monopoles, and the magnetic string is characterized by the Higgs mass of the photon. The dual version of this picture is in fact easy to imagine. We simply consider a dual photon, or dual $U(1)$ theory, now coupled to magnetic Higgs matter, a field representing the magnetic monopoles with magnetic $U(1)$ charge, and we look for a dual Higgs mechanism that, by a vacuum expectation value of the monopole field, will induce a Higgs mass for the dual magnetic photon. This mass gap will characterize the confinement phase. As the reader may realize, this whole picture of confinement is based on Higgs, or dual Higgs mechanisms for abelian gauge theories; however, in standard QCD, we expect confinement to be related to the very non abelian nature of the gauge groups. Indeed, only non abelian gauge theories are asymptotically free, and would possess the infrared slavery, or confinement, phenomena. Moreover, in a pure non abelian gauge theory, we do not have the right topology to define stable 't Hooft-Polyakov abelian monopoles, so the extension of the superconductivity picture to the $N=0$ pure Yang-Mills theory, or standard QCD, is far from being direct. Along the last two decades, with 't Hooft and Polyakov as leaders, some pictures for confinement have been suggested. Perhaps, the main steps in the story are

- i) 2 + 1 Polyakov quantum electrodynamics [23].
- ii) 't Hooft $\mathbf{Z}(N)$ duality relations [56].
- iii) 't Hooft twisted boundary conditions [20].
- iv) 't Hooft abelian projection gauge [58].

Concerning *i*), we have already described the relevant dynamics in chapter I. Let us therefore now consider the other points. Concerning *ii*), the general idea is dealing with the topology underlying pure $SU(N)$ Yang-Mills theory, namely

$$\Pi_1(SU(N)/U(1)) \simeq \mathbf{Z}(N). \quad (2.96)$$

This is the condition for the existence of magnetic $\mathbf{Z}(N)$ vortices. The 't Hooft loop $B(C)$ is the magnetic analog of the Wilson loop $A(C)$, and was defined for creating a $\mathbf{Z}(N)$ magnetic flux tube along the path C . The Wilson criteria for confinement, $A(C)$ going like the area, has now its dual in $B(C)$ behaving like the perimeter, reproducing again the picture that dual Higgs is equivalent to confinement. The duality relations established by 't Hooft reduce to

$$A(C)B(C') = e^{2\pi i \nu(C, C')/N} B(C')A(C), \quad (2.97)$$

where $\nu(C, C')$ is the link number between the loops C and C' . From (2.96), the different possible phases compatible with duality were obtained. A way to make more quantitative the previous picture was also introduced by 't Hooft, by means of twisted boundary conditions in a box. Some of the main ingredients were already introduced in chapter I, but we will come back to them later on. In what follows of this section we will mainly be interested in the abelian projection gauge.

The idea of the abelian projection gauge was originally that of defining a unitary gauge, i. e., a gauge absent of ghosts. The simplest way to do it is first reducing the theory to an abelian one, and then fixing the gauge, which is (in the abelian theory) a certainly easier task. Using a formal notation, if G is the non abelian gauge group, and L is its maximal abelian subgroup, then the non abelian part is simply given by G/L , so that we can take, as the degrees of freedom for the abelian gauge theory, the space $R/(G/L)$, where R generically represents the whole space of gauge configurations. Now, the theory defined by $R/(G/L)$, is an abelian theory, and we can fix the gauge, going finally to the unitary gauge, characterized by $R/G = L \backslash R/(G/L)$. Now, two questions arise, concerning the content of the intermediate abelian theory, $R/(G/L)$, and the more important point of how such a theory should be defined. In order to fix the non abelian part of the gauge group, i. e., the piece G/L , 't Hooft used the following trick [58]: let X be a field that we can think of as a functional of A , $X(A)$, or an extra field that will be decoupled at the end. For the time being, we simply think of X as a functional, $X(A)$. We will require $X(A)$ to transform under the adjoint representation, i. e.,

$$X(A) \rightarrow gX(A)g^{-1}. \quad (2.98)$$

Now, the gauge condition that fixes the non abelian part of the gauge group is

$$X(A) = \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_N \end{pmatrix}. \quad (2.99)$$

Indeed, if $X(A)$ is diagonal, the residual group is just the maximal abelian subgroup. Notice that $X(A)$ is playing a similar role to a Higgs field in the adjoint representation, and (2.99) is what we will interpret as a vacuum expectation value, breaking the G symmetry to its maximal abelian subgroup. As in the standard Higgs mechanism, now the degrees of freedom are the diagonal parts of the gauge field, $A_\mu^{(ij)}$, that transform as $U(1)$ charged particles. In addition, we have the N scalar fields λ , appearing in (2.99). Summarizing, the particle content we get in the maximal abelian gauge is

- i) $N - 1$ photons, $A^{(ii)}$.
- ii) $\frac{1}{2}N(N - 1)$ charged particles, $A^{(ij)}$.
- iii) N scalar fields, λ_i .

Notice that (2.99) does not require the λ_i to be constant; in fact, λ_i are fields depending on the spacetime position. Another important aspect of (2.99) is that, by means of this maximal abelian gauge we are not introducing, in principle, any form of potential for the λ_i fields, so that their expectation values are a priori undetermined. Concerning the previous spectrum, charged particles of type *ii*) can be considered formally massive, with the mass being proportional to $\lambda_i - \lambda_j$, as is the case in the standard Higgs mechanism.

The spectrum *i*), *ii*) and *iii*) is not complete. Extra spectrum, corresponding to singularities of the maximal abelian gauge, (2.99), is also allowed. These singularities correspond to points in spacetime, where $\lambda_i(x) = \lambda_{i+1}(x)$, i. e., where two eigenvalues coincide. We have imposed that $\lambda_i > \lambda_{i+1}$, i. e., the eigenvalues of (2.99) are ordered. These singularities are point-like in three dimensions, and $d - 3$ dimensional for spaces of dimension d . It is easy to see that these singularities of the gauge (2.99) are ‘t Hooft-Polyakov monopoles. Once we have this set of degrees of freedom to describe the non abelian theory, we may proceed to consider the phenomenon of confinement, following in essence the same philosophy as in abelian superconductors. ‘t Hooft’s rules of construction are:

- R1 Eliminate the electric charges. This means constructing an effective lagrangian, where the “massive” electric particles $A^{(ii)}$ have been integrated out inside loops.
- R2 Perform duality transformations on the effective lagrangian obtained upon the above integration of the electric charges, going to dual photons. These dual photons should interact with the charged monopoles by ordinary vertices, coupling the dual photon to two monopoles. The interaction between monopoles is certainly not reduced to the single exchange of dual photons; there is in practice a missing link connecting the dual photon-monopole vertices, and the effective lagrangian, and which is played by the λ -fields: the Γ_{eff} action depends also on the λ -fields, that have Yukawa coupling with the charged $A^{(ij)}$ particles, running inside the loop. As we dualize, we should also take into account duality on these fields λ_i . In fact, this should be the most relevant part of our story, as it is the potential interaction between monopoles and the dual λ_i fields what naturally leads to next rule.
- R3 The expectation value $\langle M \rangle$, for the theory obtained in R2, must be computed. In fact, this vacuum expectation value should be obtained after minimizing the theory with respect to the λ_i field values.

In spite of the beautiful physical structure underlying ‘t Hooft’s approach, this program is far from being of practical use in standard QCD or pure Yang-Mills theory. However, progress in lattice computations is being made at present.

After this introduction to ‘t Hooft’s abelian projection gauge, let us come back to the simpler example of $N=2$ pure Yang-Mills theory to find out the validity of the above rules. The careful reader will have already found some similarities in our discussion and the way the Seiberg-Witten solution for $N=2$ supersymmetric Yang-Mills has been presented. In fact, in the $N=2$ theory, the X field can simply be interpreted as the Higgs field in the adjoint, breaking $SU(2)$ to $U(1)$ on generic points of the moduli (for a group of higher rank, r , the breaking is down to $U(1)^r$). Moreover, we also have the spectrum of ‘t Hooft-Polyakov monopoles and, according to degrees of freedom, we are certainly quite close to the abelian projection picture; however, we should be careful at this point. In ‘t Hooft’s abelian projection, it was not assumed at any moment that we must be at a Higgs phase with well defined massive monopoles. The type of monopoles we find in the abelian projection gauge are not massive in the usual sense and, moreover, they have not finite size but are simply point like singularities.

Rule R1 is almost accomplished through the Seiberg-Witten solution [35, 36]. In fact, we can consider the effective lagrangian obtained from $\Gamma_{eff}(A_\mu^0, a)$, where A_μ^0 represents

the photon, and a is the scalar field in the $N=2$ hypermultiplet (notice that this effective lagrangian is constrained to be $N=2$ invariant). For each value of $u = \frac{1}{2} < \text{tr}\phi^2 >$, the vacuum expectation value of the field a , in the perturbative regime, is simply

$$a(u) = \sqrt{2u}. \quad (2.100)$$

The effective lagrangian contains only one loop logarithmic contributions (see equation (2.36)), and instanton effects. The instanton and multiinstanton contributions contribute each with four fermionic zero modes, as we kill the four zero modes associated with superconformal transformations. The expansion of the effective lagrangian in perturbative and non perturbative effects can be done in the weak coupling regime and, if we know how to perform the duality transformation, we can start obtaining non trivial information on the strong coupling regime. Let us formally denote through $\Gamma_{eff}(A^D, a_D)$ the dual effective lagrangian. In the dual perturbative regime, the effective lagrangian is an expansion in one loop terms, corresponding to light magnetic monopoles, and non perturbative higher order terms. From the moduli space point of view, the dual perturbative expansion should appear as a good description of the infrared region, i. e., for values of u such that the electric constant is large, which are points at the neighbourhood of $u \simeq \Lambda$, with Λ the dynamically generated scale. To complete the equivalent dual description, the equivalent to expression (2.100) for the dual variable a_D should be constructed; imposing $N=2$ supersymmetry, we obtain that the dual theory has a coupling

$$a_D M \tilde{M}, \quad (2.101)$$

of Yukawa type for monopoles. Then, a_D is the mass of the monopole, in the very same way as the mass of $W^\pm$ particles, in the standard Higgs mechanism, is given by a . We can now write a general formula for electrically and magnetically charged particles,

$$M(n_e, n_m) = |n_e a + n_m a_D|. \quad (2.102)$$

Here, we have only motivated equation (2.102) from physical arguments but, as we will see, the mathematical and supersymmetric meaning of (2.102) goes far beyond the scope of the simple argument we have used.

Coming back to our problem of discovering $a_D(u)$, a proper description will require some results on Kähler geometry. In fact, we know that the metric on the u moduli, is certainly Kähler with respect to the complex structure distinguished by the elliptic fibration representation of the $N=4$ three dimensional moduli space. If it has a Kähler

structure, the corresponding Kähler potential can be defined through

$$g_{u\bar{u}} = \text{Im} \left(\frac{\partial^2 K}{\partial u \partial \bar{u}} \right). \quad (2.103)$$

This Kähler potential can be read out from the effective $N=2$ low energy action. In fact, as a general statement, the metric on the moduli space is given by the quadratic terms of the effective low energy lagrangian. Now, for $N=2$ the lagrangian can be written in terms of the so called prepotential as follows:

$$\mathcal{L} = \int d^4\theta \mathcal{F}(\mathcal{A}), \quad (2.104)$$

where $\mathcal{A}$ is an $N=2$ superfield, which is holomorphic or, in supersymmetric language, depends only on chiral fields. The Kähler potential is derived from $\mathcal{F}$ as

$$K = \text{Im} \left(\frac{\partial \mathcal{F}}{\partial \mathcal{A}} \cdot \bar{\mathcal{A}} \right), \quad (2.105)$$

from which (2.103) becomes

$$g_{u\bar{u}} = \text{Im} \left(\frac{\partial a_D}{\partial u} \frac{\partial \bar{a}}{\partial \bar{u}} \right), \quad (2.106)$$

where we have defined

$$a_D \equiv \frac{\partial \mathcal{F}}{\partial a}, \quad (2.107)$$

in the sense of lower components. Using (2.104) and (2.105) we get, for the metric,

$$ds^2 = \text{Im} \frac{\partial^2 \mathcal{F}}{\partial a^2} da d\bar{a}, \quad (2.108)$$

and therefore we can identify $\frac{\partial^2 \mathcal{F}}{\partial a^2}$ with $\tau(u)$ or, equivalently,

$$\tau(u) = \frac{da_D}{da}. \quad (2.109)$$

Notice that equation (2.109) is perfectly consistent with what we expect for the definition of a_D , as it provides the mass of the monopole. In the perturbative regime, we know that it behaves like $\text{Im} \tau \cdot a \simeq \frac{a}{g^2}$. Therefore, (2.107) is the right generalization. Fortunately, thanks to (2.109) and relation (2.66), we get a definite representation of $a_D(u)$ in terms of the Seiberg-Witten elliptic fibration solution,

$$\frac{da_D}{du} = \oint_{\gamma} \varphi(z; u) dz, \quad (2.110)$$

where φ is the holomorphic differential of E_u , which is given by

$$\varphi(z; u) = \frac{dx}{y}. \quad (2.111)$$

Now that we have a candidate for $a_D(u)$, we can continue our analysis following 't Hooft's rules (in fact, $a_D(u)$ or, equivalently, how to define the dual scalar field λ_i^D , was a missing part in 't Hooft's program). Next, we want to work out the dynamics of the monopoles. Until now, we have used $N=2$ dynamics, so that the fields a and a_D are part of our original lagrangian, and not a gauge artifact, as in 't Hooft's abelian projection gauge. However, if we softly break $N=2$ to $N=1$ [35] adding a mass term for the scalar fields,

$$m \text{tr} \Phi^2, \quad (2.112)$$

then for large enough m the low energy theory is $N=1$, where the interpretation of the fields a and a_D should become closer and closer to the fields of the abelian projection. The soft breaking term (2.112) should reproduce 't Hooft's hidden dynamics governing the λ -fields. In fact, there is a simple procedure, discovered by Seiberg and Witten, to do that. The effect of (2.112) on the low energy description of the theory is to add a superfield $\mathcal{U}$, with lower component u , such that $\langle u \rangle = \langle \text{tr} \phi^2 \rangle$, with superpotential

$$W = m \mathcal{U}. \quad (2.113)$$

This extra term contains in fact the dynamics about a_D fields we are looking for, so we can write (2.113) as

$$W = m \mathcal{U}(a_D), \quad (2.114)$$

and interpret it as a lagrangian term for a_D . The monopole dynamics is then controlled by a superpotential of type

$$W = a_D M \tilde{M} + m \mathcal{U}(a_D), \quad (2.115)$$

where the first term is the $N=2$ Yukawa coupling. Now, in order to fulfill rule R3, we only need to minimize the superpotential (2.115). Clearly, we get two minima with monopole vacuum expectation value given by

$$\langle M \rangle = \pm \left(\frac{\partial \mathcal{U}}{\partial a_D} \right)^{1/2} m, \quad (2.116)$$

which is the desired proof of confinement. 't Hooft's program is then completed. In order to extend this approach to non supersymmetric theories, we can still use the the trick

of adding a mass term for the X field; however, because of the lack of holomorphy, no translation of such procedure in the form of (2.114) is possible.

Instead of using the relation for $\mathcal{U}(a_D)$, we can try to get a more direct geometrical interpretation of (1.103): let us work with the curve (2.82), and consider the points A and B with $y = 0$,

$$x^2 - xu + \Lambda^2 = 0. \quad (2.117)$$

Now, we can define the function

$$\mathcal{U}(x) = x + \frac{\Lambda^2}{x}. \quad (2.118)$$

The purpose of this function is giving a value of $\mathcal{U}$, such that x is one of the crossing points. Obviously, $\mathcal{U}(x)$ possesses two minima, at

$$x = \pm\Lambda, \quad (2.119)$$

and therefore the superpotential $m\mathcal{U}$ has two minima, at $\pm\Lambda_1$, with Λ_1 the scale of the $N = 1$ theory. Of course, the minima of $\mathcal{U}(x)$ take place when the two points A and B coincide, i. e., at the singular nodal curves. Now, we can use the following heuristic argument to find out what happens in the three dimensional $R \rightarrow 0$ limit. In projective coordinates, the region at infinity of (2.82) is

$$zy^2 = x^3 - zx^2u + \Lambda^2xz^2, \quad (2.120)$$

at $z = 0$. If we delete the infinity point, i. e., the intersection of the projective curve C defined by (2.120) and $H_\infty = \{(x, y, 0)\}$, and we then put $x^3 = 0$ in (2.120) we get, instead of (2.118) [37],

$$\mathcal{U}_{3D}(x) = \frac{\Lambda_{N=2}^2}{x}, \quad (2.121)$$

with $\Lambda_{N=2}^2$ the $N = 2$ three dimensional scale.

3 Chapter III

Taking into account the enormous amount of good reviews and books [59, 61, 60, 62] in string theory, we will reduce ourselves in this section to simply establishing some notation and motivating fundamental relations as mass formulas.

3.1 Bosonic String.

3.1.1 Classical Theory.

Let us start considering classical bosonic string theory in flat Minkowski spacetime. This physical system is characterized by the lagrangian

$$\mathcal{L} = -\frac{T}{2} \int d^2\sigma \sqrt{-h} h^{\alpha\beta} \partial_\alpha X \partial_\beta X, \quad (3.1)$$

where $h^{\alpha\beta}$ is the worldsheet metric. The equations of motion, with respect to $h^{\alpha\beta}$, imply that

$$T_{\alpha\beta} = -\frac{2}{T} \frac{1}{\sqrt{-h}} \frac{\delta S}{\delta h^{\alpha\beta}} = 0. \quad (3.2)$$

The parameter T in (3.1) has units of squared mass, and can be identified with the string tension,

$$T = \frac{1}{2\pi\alpha'}. \quad (3.3)$$

Using the Weyl invariance of (3.1), the gauge

$$h_{\alpha\beta} = \eta_{\alpha\beta} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \quad (3.4)$$

can be chosen. In this gauge, the equations of motion for (3.1) become

$$\square X = 0. \quad (3.5)$$

Defining light cone coordinates,

$$\begin{aligned} \sigma^- &= \tau - \sigma, \\ \sigma^+ &= \tau + \sigma, \end{aligned} \quad (3.6)$$

the generic solution to (3.5) can be written as

$$X^\mu = X_R^\mu(\sigma^-) + X_L^\mu(\sigma^+). \quad (3.7)$$

Now, we will introduce open and closed strings. We will first work out the case of the closed bosonic string; in this case, we impose periodic boundary conditions,

$$X^\mu(\tau, \sigma) = X^\mu(\tau, \sigma + \pi). \quad (3.8)$$

The solution to (3.5), compatible with these boundary conditions, becomes

$$\begin{aligned} X_R^\mu &= \frac{1}{2}x^\mu + \frac{1}{2}(2\alpha')p^\mu(\tau - \sigma) + i\sqrt{\frac{\alpha'}{2}} \sum_{n \neq 0} \frac{1}{n} \alpha_n^\mu e^{-2in(\tau - \sigma)}, \\ X_L^\mu &= \frac{1}{2}x^\mu + \frac{1}{2}(2\alpha')p^\mu(\tau + \sigma) + i\sqrt{\frac{\alpha'}{2}} \sum_{n \neq 0} \frac{1}{n} \tilde{\alpha}_n^\mu e^{-2in(\tau + \sigma)}. \end{aligned} \quad (3.9)$$

Using this Fourier decomposition we get, for the hamiltonian,

$$H = \frac{1}{2} \left[\sum_{-\infty}^{\infty} \alpha_{m-n} \alpha_n + \sum_{-\infty}^{\infty} \tilde{\alpha}_{m-n} \tilde{\alpha}_n \right], \quad (3.10)$$

where we have used the notation

$$\alpha_0^\mu = \sqrt{\frac{\alpha'}{2}} p^\mu. \quad (3.11)$$

Using now (3.2), we get the classical mass formula

$$M^2 = \frac{2}{\alpha'} \sum_{n=1}^{\infty} (\alpha_{-n} \alpha_n + \tilde{\alpha}_{-n} \tilde{\alpha}_n). \quad (3.12)$$

The constraint (3.2) also implies that the left and right contributions to (3.12) are equal. Using the standard quantization rules,

$$\begin{aligned} [\alpha_m^\mu, \tilde{\alpha}_n^\nu] &= 0, \\ [\alpha_m^\mu, \alpha_n^\nu] &= m \delta_{m+n} \eta^{\mu\nu}, \\ [\tilde{\alpha}_m^\mu, \tilde{\alpha}_n^\nu] &= m \delta_{m+n} \eta^{\mu\nu}, \\ [x^\mu, p^\nu] &= i \eta^{\mu\nu}, \end{aligned} \quad (3.13)$$

and taking into account the normal ordering factors we get, for $\alpha' = \frac{1}{2}$,

$$M^2 = -8a + 8 \sum_{n=1}^{\infty} \tilde{\alpha}_{-n} \tilde{\alpha}_n = -8a + 8 \sum_{n=1}^{\infty} \alpha_{-n} \alpha_n. \quad (3.14)$$

Two things are left free in deriving (3.14), the constant a , defining the zero point energy, and the number of dimensions of the target space. The classical way to fix these constants is imposing Lorenz invariance in the light cone gauge, where physical degrees of freedom are reduced to transversal oscillations. The result, for the closed bosonic string, is that a should equal one and the number of dimensions should be 26.

From (3.14), we can easily deduce the spectrum of massless states. First of all, we have a tachyon with no oscillator modes, and squared mass negative (-8) . The massless modes are of the type

$$\alpha_{-1}^\mu \alpha_{-1}^\mu |0\rangle. \quad (3.15)$$

To discover the meaning of these modes, we can see the way they transform under $SO(24)$ in the light cone gauge; then, we get three different types of particles: gravitons for the symmetric and traceless part, a dilaton for the trace part and, finally, the antisymmetric part.

3.1.2 Background Fields.

The simplest generalization of the worldsheet lagrangian (3.1) corresponds to including background fields. The obvious is the $G^{\mu\nu}$ metric of the target spacetime,

$$S_1 = -\frac{T}{2} \int d^2\sigma \sqrt{h} h_{\alpha\beta} G^{\mu\nu}(X) \partial_\alpha X_\mu \partial_\beta X_\nu. \quad (3.16)$$

However, not any background $G^{\mu\nu}$ is allowed, since we want to preserve Weyl invariance on the worldsheet. Scale invariance, for the two dimensional system defined by (3.16) is equivalent, from the quantum field theory point of view, to requiring a vanishing β -function. At one loop, the β -function for (3.16) is given by

$$\beta = -\frac{1}{2\pi} R, \quad (3.17)$$

for $\alpha' = \frac{1}{2}$, and with R the Ricci tensor of the target spacetime. Therefore, the first condition we require on allowed spacetime backgrounds is to be Ricci flat manifolds. We will allow the addition of extra manifolds to (3.16), namely the spectrum of massless particles of the bosonic closed string,

$$S = S_1 - \frac{T}{2} \int d^2\sigma \epsilon^{\alpha\beta} \partial_\alpha X^\mu \partial_\beta X^\nu B_{\mu\nu}(X) + \frac{1}{4} \int d^2\sigma \sqrt{h} \Phi(X) R^{(2)}, \quad (3.18)$$

where $R^{(2)}$ in (3.18) is the worldsheet curvature. α' does not appear in the last term due to dimensional reasons (the first two terms in (3.18) contain the X^μ field, which has length units).

Notice that for a constant dilaton field, the last term in (3.18) is simply

$$\chi \cdot \Phi, \quad (3.19)$$

with χ the Euler number; in terms of the genus, g , for a generic Riemann surface the Euler number is simply given by

$$\chi = 2 - 2g. \quad (3.20)$$

Thus, the powers of Φ in the partition function behave like $2-2g$. This topological number possesses a nice meaning in string theory: it is equal to the number of vertices joining three closed strings, needed to build up a Riemann surface of genus g . This naturally leads to a precise physical meaning of the dilaton background field: it is the string coupling constant,

$$g = e^\Phi. \quad (3.21)$$

Once the background fields in (3.18) have been added, the condition of Weyl invariance generalizes to vanishing β -funtions for G , B and Φ . At one loop, they are

$$\begin{aligned} R_{\mu\nu} + \frac{1}{4}H_\mu^{\lambda\rho}H_{\nu\lambda\rho} - 2D_\mu D_\nu \Phi &= 0, \\ D_\lambda H_{\mu\nu}^\lambda - 2(D_\lambda \Phi)H_{\mu\nu}^\lambda &= 0, \\ 4(D_\mu \Phi)^2 - 4D_\mu D^\mu \Phi + R + \frac{1}{12}H_{\mu\nu\rho}H^{\mu\nu\rho} + (D - 26) &= 0, \end{aligned} \quad (3.22)$$

where $H_{\mu\nu\rho} = \partial_\mu B_{\nu\rho} + \partial_\rho B_{\mu\nu} + \partial_\nu B_{\rho\mu}$.

3.1.3 World Sheet Symmetries.

Before ending this quick survey on the bosonic string, let us mention an aspect of world-sheet symmetries. Worldsheet parity acts exchanging left and right oscillators,

$$\Omega : \alpha_n^\mu \leftrightarrow \tilde{\alpha}_n^\mu. \quad (3.23)$$

Among massless states (3.15), only the symmetric part (the graviton) is invariant under this transformation. We can now reduce the Hilbert space to states invariant under Ω . The immediate effect of this on the worldsheet geometry is that a one loop surface can be defined in two ways: the opposite S^1 boundaries of a cylinder can be glued preserving orientation, to generate a torus, or up to an Ω trasnformation, giving rise to a Klein bottle.

3.1.4 Toroidal Compactifications.

A torus is a Ricci flat manifold that can be used as target spacetime. Let us consider the simplest case, $\mathbf{R}^{25} \times S^1$, where the compact S^1 dimension is taken to be of radius R .

Then, the coordinate x^{25} , living on this S^1 , must satisfy

$$x^{25} \equiv x^{25} + 2\pi nR. \quad (3.24)$$

If we now include the identification (3.24) in the mode expansion (3.9) we get, for the right and left momenta,

$$\begin{aligned} p_L &= \frac{m}{2R} - nR, \\ p_R &= \frac{m}{2R} + nR, \end{aligned} \quad (3.25)$$

while the mass formula becomes

$$M^2 = 4 \left(\frac{m}{2R} - nR \right)^2 + 8(N - 1) = 4 \left(\frac{m}{2R} + nR \right)^2 + 8(\bar{N} - 1), \quad (3.26)$$

with N and $\bar{N}$ the total level of left and right moving excitations, respectively. The first thing to be noticed, from (3.25), is the invariance under the transformation

$$\begin{aligned} T : R &\rightarrow \frac{1}{2R}, \\ m &\rightarrow n. \end{aligned} \quad (3.27)$$

A nice way to represent (3.25) is using a lattice of $(1, 1)$ type, which will be referred to as $\Gamma^{1,1}$. This is an even lattice, as can be observed from (3.25),

$$p_L^2 - p_R^2 = 2mn. \quad (3.28)$$

If Π is the spacelike 1-plane where p_L lives, then $p_R \in \Pi^\perp$. In fact, p_L forms a θ^{12} angle with the positive axis of the $\Gamma^{1,1}$ lattice, while p_R forms a negative angle, $-\theta$, and changes in R , which are simply changes in θ (or Lorentz rotations in the $\Gamma^{1,1}$ hyperbolic space), are changes in the target space preserving the $\beta = 0$ condition, and therefore are what can be called the *moduli of the σ -model* (3.16). Of course, no change arises in the spectrum upon rotations of the Π and $\Pi^\perp$ planes. We have now obtained a good characterization of the moduli space for the string σ -model on a simple S^1 torus. However, in addition to rotations in Π and $\Pi^\perp$, we should also take into account the symmetry (3.27), representing rotations of the $\Gamma^{1,1}$ lattice.

The previous discussion can be generalized to compactifications on higher dimensional tori, T^d (i. e., working in a background spacetime $\mathbf{R}^{26-d} \times T^d$). In this case, (p_L, p_R) will belong to a lattice $\Gamma^{d,d}$, and the moduli space will be given by [64]

$$O(d, d; \mathbf{Z}) \backslash O(d, d) / O(d) \times O(d), \quad (3.29)$$

¹² θ is the coordinate parametrizing the radius of the compact dimension.

where the $O(d, d; \mathbf{Z})$ piece generalizes the T -transformations (3.27) to T^d . From now we will call these transformations T -duality [65]. Notice also that the dimension of the moduli (3.29) is $d \cdot d$, which is the number of massless degrees of freedom that have been used to define the background fields of the σ -model (3.18). The manifold (3.29) is the first example of moduli of a σ -model we find; these moduli spaces will be compared, in next section, to the $K3$ moduli described.

3.1.5 σ -Model $K3$ Geometry. A First Look at Quantum Cohomology.

The concept of moduli space introduced in previous paragraph, for the σ -model (3.18), when the target space is a T^d torus, leading to manifold (3.29), can be generalized to more complicated spacetime geometries satisfying the constraints derived from conformal invariance, namely Ricci flat manifolds. This is a physical way to approach the theory of moduli spaces where, instead of working out the cohomology of the manifold, a string is forced to move on it, which allows to wonder about the moduli of the so defined conformal field theory. In order to properly use this approach, let us first review some facts about $K3$ geometry.

Let us first recall the relation between supersymmetry and the number of complex structures. Let us think of a σ -model, with target space $\mathcal{M}$. Now, we want this σ -model to be invariant under some supersymmetry transformations. It turns out that in order to make the σ -model, whose bosonic part is given by

$$\eta^{\mu\nu} g_{ij}(\phi(x)) \partial_\mu \phi^i \partial_\nu \phi^j, \quad (3.30)$$

with η the metric on spacetime, and g the metric on the target, invariant under $N = 2$ supersymmetry we have to require the manifold to be Kähler and, in order to be $N = 4$, to be hyperkähler.

Let us now enter the description of the $K3$ manifold [66, 67, 68]. To characterize topologically $K3$, we will first obtain its Hodge diamond. The first property of $K3$ is that the canonical class,

$$K \equiv -c_1(T), \quad (3.31)$$

with $c_1(T)$ the first Chern class of the tangent bundle, T , is zero,

$$K = 0. \quad (3.32)$$

Equation (3.32) implies that there exists a holomorphic 2-form Ω , everywhere non vanishing. Using the fact that only constant holomorphic functions are globally defined, we

easily derive, from (3.32), that

$$\dim H^{2,0} = h^{2,0} = 1. \quad (3.33)$$

In fact, if there are two different 2-forms Ω_1 and Ω_2 , then Ω_1/Ω_2 will be holomorphic and globally defined, and therefore constant.

The second important property characterizing $K3$ is

$$\Pi_1 = 0, \quad (3.34)$$

so that

$$h^{1,0} = h^{0,1} = 0, \quad (3.35)$$

as $b_1 = h^{1,0} = h^{0,1} = 0$, because of (3.34).

The Euler number can be now derived using Noether-Riemann theorem, and property (3.32), and it turns out to be 24. Using now the decomposition of the Euler number as an alternating sum of Betti numbers, we can complete the Hodge diamond,

$$24 = b_0 - b_1 + b_2 - b_3 + b_4 = 1 - 0 + b_2 - 0 + 1, \quad (3.36)$$

which implies that

$$\dim H^2 = 22, \quad (3.37)$$

and therefore, from (3.33), we get

$$\dim H^{1,1} = h^{1,1} = 20, \quad (3.38)$$

leading to the Hodge diamond

$$\begin{array}{ccccc} & & 1 & & \\ & 0 & & 0 & \\ 1 & & 20 & & 1 \\ & 0 & & 0 & \\ & & 1 & & \end{array} \quad (3.39)$$

Using Hirzebruch's pairing, we can give an inner product to the 22 dimensional space H^2 . In homology terms, we have

$$\alpha_1 \cdot \alpha_2 = \#(\alpha_1 \cap \alpha_2), \quad (3.40)$$

with $\alpha_1, \alpha_2 \in H^2(X, \mathbb{Z})$, and $\#(\alpha_1 \cap \alpha_2)$ the number of oriented intersections. From the signature complex,

$$\tau = \int_X \frac{1}{3}(c_1^2 - 2c_2) = -\frac{2}{3} \int_X c_2 = -\frac{2 \cdot 24}{3} = -16, \quad (3.41)$$

we know that $H^2(X, \mathbf{Z})$ is a lattice of signature $(3, 19)$. The lattice turns out to be self dual, i. e., there exists a basis α_i^* such that

$$\alpha_i \cdot \alpha_j^* = \delta_{ij}, \quad (3.42)$$

and even,

$$\alpha \cdot \alpha \in 2\mathbf{Z}, \quad \forall \alpha \in H^2(X, \mathbf{Z}). \quad (3.43)$$

Fortunately, lattices with these characteristics are unique up to isometries. In fact, the $(3, 19)$ lattice can be represented as

$$E_8 \perp E_8 \perp \mathcal{U} \perp \mathcal{U} \perp \mathcal{U}, \quad (3.44)$$

with $\mathcal{U}$ the hyperbolic plane, with lattice $(1, 1)$, and E_8 the lattice of $(0, 8)$ signature, defined by the Cartan algebra of E_8 . The appearance of E_8 in $K3$ will be at the very core of future relations between $K3$ and string theory, mainly in connection with the heterotic string.

Next, we should separately characterize the complex structure and the metric of $K3$. Recall that this is exactly what we did in our study of the moduli of $N=4$ supersymmetric three dimensional Yang-Mills theories. Concerning the complex structure, the proper tool to be used is Torelli's theorem, that establishes that the complex structure of a $K3$ marked surface¹³ is completely determined by the periods of the holomorphic 2-form, Ω . Thus, the complex structure is fixed by

- i) The holomorphic form Ω .
- ii) A marking.

To characterize $\Omega \in H^{2,0}(X, \mathbf{C})$, we can write

$$\Omega = x + iy, \quad (3.45)$$

with x and y in $H^2(X, \mathbf{R})$, that we identify with the space $\mathbf{R}^{3,19}$. Now, we know that

$$\begin{aligned} \int_X \Omega \wedge \Omega &= 0, \\ \int_X \Omega \wedge \bar{\Omega} &> 0, \end{aligned} \quad (3.46)$$

¹³By a marked $K3$ surface we mean a specific map of $H^2(X, \mathbf{Z})$ into the lattice (3.44), that we will denote, from now on, $\Gamma_{3,19}$.

and we derive

$$\begin{aligned} x \cdot y &= 0, \\ x \cdot x &= y \cdot y. \end{aligned} \tag{3.47}$$

Therefore, associated with Ω , we define a plane of vectors $v = nx + my$ which, due to (3.46), is space-like, i. e.,

$$v \cdot v > 0. \tag{3.48}$$

The choice of (3.45) fixes an orientation of the two plane, that changes upon complex conjugation. Thus, the moduli space of complex structures of $K3$, will reduce to simply the space of oriented space-like 2-planes in $\mathbf{R}^{3,19}$. To describe this space, we can use a Grassmanian [67],

$$Gr = \frac{(O(3, 19))^+}{(O(2) \times O(1, 19))^+}, \tag{3.49}$$

where $(\)^+$ stands for the part of the group preserving orientation. If, instead of working with the particular marking we have been using, we change it, the result turns out to be an isometry of the $\Gamma^{3,19}$ lattice; let us refer to this group by $O(\Gamma^{3,19})$. The moduli then becomes

$$\mathcal{M}^C = Gr/O^+(\Gamma^{3,19}). \tag{3.50}$$

The group $O(\Gamma^{3,19})$ is the analog to the modular group, when we work out the moduli space of complex structures for a Riemann surface ($Sl(2, \mathbf{Z})$ for a torus).

Let us now make some comments on the distinguished complex structure we have used in the study of the moduli of the three dimensional $N = 4$ theories. This complex structure is such that the the elliptic curve is a $(1, 1)$ -form, and is characterized by the 2-form

$$\Omega = du \wedge \frac{dx}{y}, \tag{3.51}$$

with $\frac{dx}{y}$ the holomorphic differential on the elliptic fiber. However, before entering a more detailed discussion on this issue, let us consider the question of metrics. Once a complex structure has been introduced, we have a Hodge decomposition of H^2 , as

$$H^2 = H^{2,0} \oplus H^{1,1} \oplus H^{0,2}. \tag{3.52}$$

Thus, relative to a complex structure characterized by Ω , the Kähler form J in $H^{1,1}$ is orthogonal to Ω , and such that

$$\text{Vol} = \int_X J \wedge J > 0, \tag{3.53}$$

which means that J is represented by a space-like vector in $\mathbf{R}^{3,19}$ and, therefore, together with Ω , spans the whole three dimensional space-like subspace of $\mathbf{R}^{3,19}$. Yau's theorem now shows how the metric is completely determined by J and Ω , i. e., by a space-like 3-plane in $\mathbf{R}^{3,19}$. Thus, we are in a similar position to the characterization of the moduli space of complex structures, and we end up with a Grassmannian manifold of three space-like planes in $\mathbf{R}^{3,19}$,

$$Gr = O(3, 19)/O(3) \times O(19). \quad (3.54)$$

Now, we need to complete Gr with two extra ingredients. One is the volume of the manifold, that can change by dilatations, and the other is again the modular part, corresponding to isometries of $\Gamma^{3,19}$, so that finally we get

$$\mathcal{M}^M = O(\Gamma_{3,19}) \backslash Gr \times \mathbf{R}^+. \quad (3.55)$$

Hence, the moduli of the σ -model (3.18), defined on a $K3$ surface, will contain the moduli of Einstein metrics on $K3$ (see equations (3.54) and (3.55)). Now, the dimension of manifold (3.26) is 58. For the σ -model (3.18) we must also take into account the moduli of B -backgrounds. In the string action, what we have is the integral, $\int B$, over the worldsheet, which now becomes a 2-cycle of $K3$; thus, the moduli of B -backgrounds is given by the second Betti number of the $K3$ manifold, which is 22. Finally, the dilaton field Φ has to be taken into account in (3.18). As mentioned, if Φ is constant, as we will require, it counts the number of loops in the perturbation series, so we will not consider it as an extra moduli. More precisely, we will probe the $K3$ geometry working at tree level in string theory. Under these conditions, the σ moduli space is of dimension [69]

$$58 + 22 = 80, \quad (3.56)$$

and the natural guess is the manifold

$$\mathcal{M}^\sigma = O(4, 20)/O(4) \times O(20). \quad (3.57)$$

Naturally, this is not the final answer, as we have not divided yet by the equivalent to the T -duality transformations in the toroidal case, which are, for $K3$, isometries of the $H^2(X; \mathbf{Z})$ lattice, i. e.,

$$O(\Gamma^{3,19}). \quad (3.58)$$

However, the final answer is not the quotient of (3.57) by (3.58), as an important symmetry from the point of view of conformal field theory is yet being missed: mirror symmetry. In

order to get a geometrical understanding of mirror symmetry [70], we need first to define the Picard lattice.

Let us consider curves inside the $K3$ manifold. The Picard lattice is defined as

$$\text{Pic}(X) = H^{1,1}(X) \cap H^2(S, \mathbf{Z}), \quad (3.59)$$

which means curves (i. e., 2-cycles) holomorphically embedded in X . By definition (3.59), $\text{Pic}(X)$ defines a sublattice of $H^2(S; \mathbf{Z})$. This Picard lattice has signature $(18, t)$. Let us consider, as an example, an elliptic fibration where the base is a 2-cycle B , and F is the fiber. The Picard lattice defined by these two 2-cycles is given by

$$\begin{aligned} B \cdot B &= -2, \\ B \cdot F &= 1, \\ F \cdot F &= 0, \end{aligned} \quad (3.60)$$

which is a lattice of $(1, 1)$ type. Self intersections are given by the general expression

$$C \cdot C = 2(g - 1), \quad (3.61)$$

where g is the genus, so that for $g = 0$, the base space, we get -2 , and for the elliptic fiber, with $g = 1$, we get 0 for the intersection. The intersection between the base and the fiber, $B \cdot F$, reflects the nature of the fibration. Notice that expression (3.61) is consistent with the even nature of the lattice $\Gamma_{3,19}$. Now, from (3.59), it is clear that the number of curves we have in $\text{Pic}(X)$ depends on the complex structure. Taking this fact into account, we can ask ourselves about the moduli space of complex structures preserving a given Picard sublattice; for instance, we can be interested in the moduli space of elliptic fibrations preserving the structure of the fibration. As $\text{Pic}(X)$ are elements in $H^{1,1}(X)$, they should be orthogonal to Ω , so the moduli we are looking for will be defined in terms of the Grassmannian of space-like 2-planes in $\mathbf{R}^{2,19-t}$, i. e.,

$$Gr^P = O(2, 19 - t)/O(2) \times O(19 - t), \quad (3.62)$$

where we should again quotient by the corresponding modular group. This modular group will be given by isometries of the lattice Λ , called the transcendental lattice, and is simply defined as the orthogonal complement to the Picard lattice. Thus, Λ is of $\Gamma^{2,19-t}$ type, and the moduli preserving the Picard group is

$$\mathcal{M}^P = Gr^P/O(\Lambda). \quad (3.63)$$

As is clear from (3.62), the dimension of the moduli space of complex structures preserving the Picard group, reduces in an amount given by the value of t for the Picard lattice. At this point of the discussion, a question at the core of mirror symmetry comes naturally to our mind, concerning the possibility to define a manifold X^* whose Picard group is the transcendental lattice Λ of X [71]. In these terms, the answer is clearly negative, as the Picard lattice is of signature $(1, t)$, and Λ is of signature $(2, 19 - t)$, so that we need either passing from Λ to a $(1, t')$ lattice, or generalize the concept of Picard lattice, admitting lattices of signature $(2, t)$. It turns out that both approaches are equivalent, but the second has a more physical flavor; in order to get from Λ a Picard lattice, what we can do is to introduce an isotropic vector f in Λ , and define the new lattice through

$$f^\perp/f, \quad (3.64)$$

which is of $(1, 18 - t)$ type; now, the mirror manifold X^* is defined as the manifold possessing as Picard lattice the one defined by (3.64). The moduli space of the mirror manifold is therefore given by the equivalent to expression (3.62),

$$Gr^{*P} = O(2, t + 1)/O(2) \times O(t + 1). \quad (3.65)$$

Then, we observe that the dimension of the two moduli spaces sums up to 20, and that the dimension of the moduli space of the mirror manifold is exactly given by the rank $t + 1$ of the Picard of the original moduli space.

A different approach will consist in defining the so called quantum Picard lattice. Given a Picard lattice of signature $(1, t)$, we define its quantum analog as the lattice of signature $(2, t + 1)$, obtained after multiplying by the hyperbolic lattice $\Gamma^{1,1}$. So, the question of mirror will be that of given a manifold X , with transcendental lattice Λ , finding a manifold X^* such that its quantum Picard lattice is precisely Λ . Now, we observe that the quantum Picard lattices of X and X^* produce a lattice of signature $(4, 20)$. The automorphisms $O(\Gamma^{4,20})$ will result of composing the T -duality transformations and mirror symmetry. Coming back to (3.57), and including mirror symmetry, we get, as moduli space of the σ -model on $K3$,

$$O(4, 20; \mathbf{Z}) \backslash O(4, 20)/O(4) \times O(20). \quad (3.66)$$

This concludes our analysis of σ -models on $K3$.

3.1.6 Elliptically Fibered $K3$ and Mirror Symmetry.

We are now going to consider singularities in the $K3$ manifold. Let C be a rational curve in the $K3$ manifold; then, by equation (3.61), $C \cdot C = -2$. If the curve C is holomorphically

embedded it will be an element of the Picard lattice. Its volume is defined as

$$\text{Vol}(C) = J \cdot C, \quad (3.67)$$

with J the Kähler class. A singularity will appear whenever the volume of C goes zero, i. e., whenever the Kähler class J is orthogonal to C . Notice that this implies that C should be orthogonal to the whole 3-plane defined by Ω and J , as C is in fact $(1, 1)$, and therefore orthogonal to Ω .

Now, we can define the process of blowing up or down a curve C in X . In fact, a way to blow up is simply changing the moduli space of metrics J , until $J \cdot C$ becomes different from zero. The opposite is the blow down of the curve. The other way to get rid off the singularity is simply changing the complex structure in such a way that the curve is not in $H^{1,1}$, i. e., the curve does not exist anymore.

We can have different types of singularities, according to how many rational curves C_i are orthogonal to J . The type of singularity will be given by the lattice generated by these C_i curves. Again, these lattices would be characterized by Dynkin diagrams.

Let us now consider an elliptically fibered $K3$ manifold,

$$E \rightarrow X \rightarrow B. \quad (3.68)$$

Now, we can come back to Kodaira's analysis on elliptic fibrations, as presented in chapter II. Elliptic singularities of Kodaira type are characterized by the set of irreducible components X_i of the corresponding singularities. The Picard lattice for these elliptic fibrations contains the $\Gamma^{1,1}$ lattice generated by the fiber and the base, and the contribution of each singularity as given by the Shioda-Tate formula [71]. Defining the Picard number $\rho(X)$ as $1 + t$ for a Picard lattice of type $(1, t)$ we get

$$\rho(X) = 2 + \sum_{\nu} \sigma(F_{\nu}), \quad (3.69)$$

where the sum is over the set of singularities, and where σ is given by $\sigma(A_{n-1}) = n - 1$, $\sigma(D_{n+4}) = n + 4$, $\sigma(E_6) = 6$, $\sigma(E_7) = 7$, $\sigma(E_8) = 8$, $\sigma(IV) = 2$, $\sigma(III) = 1$, $\sigma(II) = 0$. Equation (3.69) is true provided the Mordell-Weyl group of sections is trivial.

As described in the previous section, the mirror map goes from a manifold X , with Picard lattice of type $(1, t)$, to X^* , with Picard lattice $(1, 18 - t)$ or, equivalently,

$$\rho(X) + \rho(X^*) = 20. \quad (3.70)$$

Through mirror, we can then pass from an elliptically fibered $K3$ surface, with Picard number $\rho(X) = 2$, which should for instance have all its singularities of type A_0 , to a $K3$

surface of Picard number $\rho(X^*) = 18$, which should have 16 singularities of A_1 type, or some other combination of singularities.

3.1.7 The Open Bosonic String.

Repeating previous comments on closed strings for the open case is straightforward. The only crucial point is deciding the type of boundary conditions to be imposed. From (3.1), we get boundary terms of the form

$$\frac{T}{2} \int \partial X^\mu \partial_n X_\mu, \quad (3.71)$$

with ∂_n the normal boundary derivative. In order to avoid momentum flow away from the string, it is natural to impose Neumann boundary conditions,

$$\partial_n X_\mu = 0. \quad (3.72)$$

Using these boundary conditions the mode expansion (3.9) becomes, for the open string,

$$X^\mu(\sigma, \tau) = x^\mu + 2\alpha' p^\mu \tau + i\sqrt{2\alpha'} \sum_{n \neq 0} \frac{1}{n} \alpha_n^\mu e^{-in\tau} \cos n\sigma, \quad (3.73)$$

and the quantum mass formula (3.14) is, for $\alpha' = \frac{1}{2}$,

$$M^2 = -2 + 2 \sum_{n=1}^{\infty} \alpha_{-n} \alpha_n. \quad (3.74)$$

Now, the first surprise arises when trying to generalize the T -duality symmetry, (3.27), to the open string case.

3.1.8 D-Branes.

By introducing the complex coordinate

$$z = \sigma^2 + i\sigma, \quad (3.75)$$

with $\sigma^2 \equiv i\tau$, (3.73) can be rewritten as

$$X^\mu(\sigma, \tau) = x^\mu - i\alpha' p^\mu \ln(z\bar{z}) + i\sqrt{\frac{\alpha'}{2}} \sum_{n \neq 0} \frac{1}{n} \alpha_n^\mu (z^{-n} + \bar{z}^{-n}). \quad (3.76)$$

Let us now consider the open string moving in $\mathbf{R}^{25} \times S^1$. Neumann boundary conditions in the compactified direction are

$$\partial_n X^{25} = 0. \quad (3.77)$$

Now, we will work out the way these boundary conditions modify under the $R \rightarrow \frac{1}{R}$ transformation [72]. To visualize the answer, we will consider the cylinder swept out by a time evolving closed string, both from the closed and open string pictures (in the open string picture the cylinder can be understood as an open string with both ends at the S^1 edges of the cylinder). In fact, from the open string point of view, the propagation of the string is at tree level, while the open string approach is a one loop effect. We will now assume that the S^1 boundary circles of the cylinder are in the 25 direction. Recalling then what happens in the closed string case, under change (3.27), the mode expansion (3.9) turns (3.27) equivalent to the change

$$\tilde{\alpha}_n^{25} \rightarrow -\tilde{\alpha}_n^{25}. \quad (3.78)$$

In the $n = 0$ case we get, from (3.11) and (3.25) (with $\alpha' = \frac{1}{2}$),

$$\alpha_0^{25} = \frac{m}{2R} - nR \rightarrow nR - \frac{m}{2R} = -\tilde{\alpha}_0^{25}. \quad (3.79)$$

What this means is that the theory in the dual circle of radius $\frac{1}{2R}$ is equivalent to a theory on a circle of radius R , but written in terms of a new space coordinate Y^{25} , defined from X^{25} by the change (3.78). Now, it easy to see that

$$\partial_\alpha Y^{25} = \epsilon_{\alpha\beta} \partial^\beta X^{25}. \quad (3.80)$$

Returning now to the cylinder image described above, let us consider boundary conditions in the open string picture. From the closed string approach, they will be represented as

$$\partial_\tau X^{25} = 0. \quad (3.81)$$

Now, after performing the duality transformation (3.27), equation (3.80) implies

$$\partial_\sigma Y^{25} = 0, \quad (3.82)$$

that, from the open string point of view, looks as Dirichlet boundary conditions, so that the extreme points of the open string do not move in time in the 25 direction. Summarizing, we observe that under $R \rightarrow \frac{\alpha'}{R}$, Neumann and Dirichlet boundary conditions for the open string are exchanged. Besides, the picture we get if the end points of the open string do not move in the 25 direction is that of D-brane hypersurfaces, with fixed 25 coordinate, where the open string should end.

For a better understanding of the dynamical nature of these D-brane hypersurfaces, and their physical meaning, the above approach must be generalized to include several D-brane hypersurfaces; the tool needed comes from the old fashioned primitive string theory, interpreted as a meson model: the Chan-Paton factors [73].

3.1.9 Chan-Paton Factors and Wilson Lines.

Chan-paton factors are simply defined encoding the end points of the open string with labels i, j , with $i, j = 1, \dots, N$. The corresponding string states will be defined as $|k; i, j\rangle$. Let us now define a set of $N \times N$ matrices, $\lambda_{N \times N}^a$, hermitian and unitary, which define the adjoint representation of $U(N)$. We can now define the open string state $|k; a\rangle$ as

$$|k; a\rangle = \sum_{i,j} \lambda_{i,j}^a |k; i, j\rangle. \quad (3.83)$$

The string states $|i, j\rangle$ can now be easily interpreted in the language of gauge theories. In order to do that, we will again use the abelian projection introduced in previous chapter. In the abelian projection gauge, states $|i, i\rangle$ correspond to $U(1)$ photons, while $|i, j\rangle$ states (non diagonal components of the gauge field) correspond to charged massive particles. The way they transform under the abelian $U(1)^N$ group is

$$|i, j\rangle \rightarrow e^{i(\alpha_j - \alpha_i)} |i, j\rangle, \quad (3.84)$$

for the abelian transformation

$$\begin{pmatrix} e^{i\alpha_1} & & \\ & \ddots & \\ & & e^{i\alpha_N} \end{pmatrix}. \quad (3.85)$$

As discussed in chapter II, to define an abelian projection gauge, a field X must be chosen to transform in the adjoint representation; then, the gauge is fixed through imposing X to be diagonal. A simple example of field X is a Wilson line. So, let us assume we are working in $\mathbf{R}^{25} \times S^1$, and define X as the Wilson line in the 25 compactified direction. Choosing X diagonal means taking A^{25} in the abelian group $U(1)^N$; a diagonal Wilson line is obtained from

$$A^{25} = \frac{1}{2\pi R} \begin{pmatrix} \theta_1 & & \\ & \ddots & \\ & & \theta_N \end{pmatrix}, \quad (3.86)$$

corresponding to a pure gauge

$$A^{25} = \partial_{25} \Lambda = \partial_{25} \frac{X^{25}}{2\pi R} \begin{pmatrix} \theta_1 & & \\ & \ddots & \\ & & \theta_N \end{pmatrix}. \quad (3.87)$$

Now, $\{\theta_1, \dots, \theta_N\}$ are the analogs to $\{\lambda_1, \dots, \lambda_N\}$, used in the standard abelian projection. The effect of the Wilson line (3.86) on a charged state $|i, j\rangle$ is transforming it in the way (3.84) defines, which in particular means that the p^{25} momentum of the $|i, j\rangle$ state becomes

$$p^{25} = \frac{n}{R} + \frac{\theta_j - \theta_i}{2\pi R}. \quad (3.88)$$

When moving from R to $R' = \frac{1}{2R}$, the momentum (3.88) turns into a winding,

$$2nR' + (\theta_j R' - \theta_i R') \frac{1}{\Gamma}. \quad (3.89)$$

The geometrical meaning of (3.89) is quite clear: the open string can wind around the dual circle of radius R' any number of times, but its end points are fixed, as expected after the $R \rightarrow R'$ duality transformation, to be in $\theta_j R'$ and $\theta_i R'$ positions. Thus, the picture we get is that of several D-brane hypersurfaces fixed in the dual circle to be at positions $\theta_1 R', \dots, \theta_N R'$, and the string states of type $|i, j\rangle$ are now living between the i^{th} and j^{th} D-brane hypersurface.

Using mass formula (3.26), and equation (3.88) for the momentum, we observe that only $\alpha_{-1}^\mu |i, i\rangle$ states can be massless (the $U(1)$ photons), and the mass of the $\alpha_{-1}^\mu |i, j\rangle$ states goes like $\left(\frac{(\theta_i - \theta_j)R'}{\Gamma}\right)^2$. Both of these states have the kinematical index μ in the uncompactified directions. We can also consider the massless Kaluza-Klein states, $\alpha_{-1}^{25} |i, i\rangle$, which can be interpreted as scalars living on the 24 dimensional space defined by the D-brane hypersurface. However, this spectrum is the abelian projected gauge spectrum for a $U(N)$ gauge theory, now defined on the D-brane hypersurface. Therefore, two complementary pictures arise,

- The distribution of D-branes represents a new type of background for string theory, where a $U(N)$ Wilson line has been introduced in the internal or compactified S^1 .
- The distribution of D-branes provides, for the massless spectrum, a geometrical representation of a gauge theory living on the worldvolume of the D-brane. Moreover, the spectrum is presented as the abelian projection spectrum.

Of course, this second approach only takes into account, as is usual in string theory, low energy degrees of freedom. Properly speaking, what we are doing is embedding the gauge theory into string theory in a new way.

To end this first contact with D-branes (for more details see, for instance, [60], and references therein) we should, at least qualitatively, answer the question posed above

on the dynamical nature of D-branes. The simplest answer will be obtained analyzing the gravitational interactions through the computation of the mass density, leading to the tension of the D-brane hypersurface. A graviton, which is a closed string state can couple a D-brane, defining an interaction vertex. The disc coupling the graviton to the D-brane can be interpreted in terms of open strings ending on its circle boundary. Without performing any computation, we already know something on the order of magnitude of the process: it is a process determined by the topology of a disc, with half the Euler number of a sphere, so the order in the string coupling constant, defined in (3.21), is $O(\frac{1}{g})$.

A more detailed discussion on D-branes needs the use of more general string theories (superstring theories), which is what we will discuss in next section.

3.2 Superstring Theories.

Superstrings correspond to the supersymmetric generalization of the σ -model (3.1). This is done adding the fermionic term

$$S_F = \int d^2\sigma i\bar{\psi}^\mu \rho^\alpha \partial_\alpha \psi_\mu, \quad (3.90)$$

where ψ^μ are spinors, relative to the worldsheet, and vectors with respect to the spacetime Lorentz group, $SO(1, D-1)$. Spinors in (3.90) are real Majorana spinors, and the Dirac matrices ρ^α , $\alpha = 0, 1$, are defined by

$$\begin{aligned} \rho^0 &= \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \\ \rho^1 &= \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \end{aligned} \quad (3.91)$$

satisfying

$$\{\rho^\alpha, \rho^\beta\} = -2\eta^{\alpha\beta}. \quad (3.92)$$

The supersymmetry transformations are defined by

$$\begin{aligned} \delta x^\mu &= \bar{\epsilon} \psi^\mu, \\ \delta \psi^\mu &= -i\rho^\alpha \partial_\alpha x^\mu \epsilon, \end{aligned} \quad (3.93)$$

with ϵ a constant anticommuting spinor. Defining the components

$$\psi^\mu = \begin{pmatrix} \psi_-^\mu \\ \psi_+^\mu \end{pmatrix}, \quad (3.94)$$

the fermionic lagrangian (3.90) can be written as

$$S_F = \int d^2\sigma (\psi_-^\mu \partial_+ \psi_-^\mu + \psi_+^\mu \partial_- \psi_+^\mu), \quad (3.95)$$

with $\partial_\pm \equiv \frac{1}{2}(\partial_\tau \pm \partial_\sigma)$. As was the case for the bosonic string, we need now to specify the boundary conditions for the fermion fields, both in the open and closed string case. For open strings, there are two possibilities:

$$\begin{aligned} \text{Ramond} & : \psi_+^\mu(\pi, \tau) = \psi_-^\mu(\pi, \tau), \\ \text{Neveu-Schwarz} & : \psi_+^\mu(\pi, \tau) = -\psi_-^\mu(\pi, \tau), \end{aligned} \quad (3.96)$$

which produce the mode expansions

$$\begin{aligned} \text{Ramond} & : \psi_\mp^\mu = \frac{1}{\sqrt{2}} \sum_z d_n^\mu e^{-in(\tau \mp \sigma)}, \\ \text{Neveu-Schwarz} & : \psi_\mp^\mu = \frac{1}{\sqrt{2}} \sum_{z+\frac{1}{2}} b_n^\mu e^{-in(\tau \mp \sigma)}. \end{aligned} \quad (3.97)$$

In the case of closed strings, we can impose either periodic or antiperiodic boundary conditions for the fermions, obtaining Ramond (R) or Neveu-Schwarz (NS) for both $\psi_\pm$ fields. After quantization we get, following similar steps to those in the bosonic case, that the critical dimension is 10, and that the mass formulas and normal ordering correlators are given by

$$M^2 = 2(N_L - \delta_L) = 2(N_R - \delta_R), \quad (3.98)$$

with $\delta = \frac{1}{2}$ in the NS sector, and $\delta = 0$ in the R sector. Using this formula, and the GSO projection, we easily get the massless spectrum. For the closed string we get

$$\begin{aligned} \text{NS-NS sector} & : b_{-1/2}^\mu b_{-1/2}^\nu |0\rangle, \\ \text{NS-R sector} & : b_{-1/2}^\mu |S\rangle, \\ \text{R-R sector} & : |S\rangle \otimes |S\rangle. \end{aligned} \quad (3.99)$$

The state $|S\rangle$ corresponds to the Ramond vacua (recall $\delta = 0$ in the Ramond sector).

The d_0^μ oscillators in (3.97) define a Clifford algebra,

$$\{d_0^\mu, d_0^\nu\} = \eta^{\mu\nu}, \quad (3.100)$$

and therefore the $|S\rangle$ vacua can be one of the two $\mathbf{8}_S, \mathbf{8}_{S'}$ spinorial representations of $SO(8)$. Depending on what is the spinorial representation chosen we get, from (3.99), two

different superstring theories. In the chiral case, we choose the same chirality for the two fermionic states in the NS-R and R-NS sectors. This will lead to two gravitinos of equal chirality. Moreover, in the R-R sector we get, for same chirality,

$$\mathbf{8}_S \times \mathbf{8}_S = \mathbf{1} \oplus \mathbf{28} \oplus \mathbf{35}_S, \quad (3.101)$$

corresponding to a scalar field being identified with the axion, an antisymmetric field, and a 4-form field. We will call this superstring theory type IIB. In case we choose different chiralities for the spinor representations associated with the Ramond vacua, what we get is type IIA superstring theory, which is also an $N = 2$ theory, but this time with two gravitinos of different chirality; now, the R-R sector contains

$$\mathbf{8}_S \otimes \mathbf{8}_{S'} = \mathbf{8}_V \oplus \mathbf{56}_V, \quad (3.102)$$

i. e., a vector field and a 3-form. These are the first two types of superstring theories that we will consider.

3.2.1 Toroidal Compactification of Type IIA and Type IIB Theories. *U*-duality.

Before considering different compactifications of superstring theories, we will first review some general results on the maximum number of allowed supersymmetry, depending on the spacetime dimension.

Spinors should be considered as representations of $SO(1, d - 1)$. Irreducible representations have dimension

$$2^{\lfloor \frac{d+1}{2} \rfloor - 1}, \quad (3.103)$$

where $\lfloor \rfloor$ stands for the integer part. Depending on the dimension, the larger spinor can be real, complex or quaternionic,

$$\begin{aligned} \mathbf{R}, & \text{ if } d = 1, 2, 3 \bmod 8, \\ \mathbf{C}, & \text{ if } d = 0 \bmod 4, \\ \mathbf{H}, & \text{ if } d = 5, 6, 7 \bmod 8. \end{aligned} \quad (3.104)$$

Using (3.103) and (3.104), we get the number of supersymmetries listed in the table below¹⁴.

¹⁴This table is constrained by the physical requirement that particles with spin > 2 do not appear.

Dimension	N	Irreducible Representation
11	1	$\mathbf{R}^{32}$
10	2	$\mathbf{R}^{16}$
9	2	$\mathbf{R}^{16}$
8	2	$\mathbf{C}^8$
7	2	$\mathbf{H}^8$
6	4	$\mathbf{H}^4$
5	4	$\mathbf{H}^4$
4	8	$\mathbf{C}^2$
3	16	$\mathbf{R}^2$

The maximum number of supersymmetries in three dimensions is then 16. From the table it is also clear that through standard Kaluza-Klein compactification, starting with six dimensional $N = 1$ supersymmetry leads to four dimensional $N = 2$, and three dimensional $N = 4$ supersymmetry. We can also notice that ten dimensional $N = 1$ leads to $N = 4$ supersymmetry in four dimensions.

It must be stressed that the counting of supersymmetries after dimensional reduction is slightly more subtle if we compactify on manifolds with non trivial topology. Here, the adequate concept is the holonomy of the internal manifold; let us therefore recall some facts on the concept of holonomy. Given a Riemannian manifold $\mathcal{M}$, the holonomy group $H_{\mathcal{M}}$ is defined as the set of transformations M_{γ} associated with paths γ in $\mathcal{M}$, defined by parallel transport of vectors in the tangent bundle. The connection used in this definition is the Levi-Civita connection. In general, for a vector bundle $E \rightarrow \mathcal{M}$, the holonomy group $H_{\mathcal{M}}$ is defined by the parallel transport of v in the fiber, with respect to the connection on E . The Ambrose-Singer theorem shows how the holonomy is generated by the curvature.

Manifolds can be classified according to its holonomy group. Therefore, we get [74]

- $H_{\mathcal{M}} = O(d)$, for real manifolds of dimension d .
- $H_{\mathcal{M}} = U(\frac{d}{2})$, for Kähler manifolds.
- $H_{\mathcal{M}} = SU(\frac{d}{2})$, for Ricci flat Kähler manifolds.
- $H_{\mathcal{M}} = Sp(\frac{d}{4})$, for hyperkähler manifolds¹⁵.

The answer to the question of what the role of holonomy is in the counting of the number of supersymmetries surviving after compactification is quite simple: let us suppose we are in dimension d , so that the spinors are in $SO(1, d - 1)$. Now, the theory is

¹⁵Notice that any hyperkähler manifold is always Ricci flat.

compactified on a manifold of dimension d_1 , down to $d_2 = d - d_1$. Supersymmetries in d_2 are associated with representations of $SO(1, d_2 - 1)$, so we need to decompose an irreducible representation of $SO(1, d - 1)$, into $SO(1, d_2 - 1) \times SO(d_1)$. Now, the holonomy group of the internal manifold $H_{\mathcal{M}_{d_1}}$ will be part of $SO(d_1)$. Good spinors in d_2 dimensions would be associated with singlets of the holonomy group of the internal manifold. Let us consider the simplest case, with $d_1 = 4$; then,

$$SO(4) = SU(2) \otimes SU(2) \quad (3.105)$$

and, if our manifold is Ricci flat and Kähler, the holonomy will be one of these $SU(2)$ factors. Therefore, we will need a singlet with respect to this $SU(2)$. As an example, let us consider the spinor in ten dimensions, with $N = 1$; as we can see from the above table, it is a **16**, that we can decompose with respect to $SO(1, 5) \times SU(2) \times SU(2)$ as

$$\mathbf{16} = (\mathbf{4}, \mathbf{2}, \mathbf{1}) \otimes (\mathbf{4}, \mathbf{1}, \mathbf{2}). \quad (3.106)$$

Therefore, we only get one surviving supersymmetry in six dimensions. This is a general result: if we compactify a ten dimensional theory on a manifold of dimension four, with $SU(2)$ holonomy, we will get a six dimensional theory with only one supersymmetry. However, if the compactification is on a torus with trivial holonomy, two supersymmetries are obtained (the maximum number of supersymmetries available).

As the first contact with type IIA string theory we will then consider its compactification on a d -dimensional torus, T^d . To start with, let us work in the particular case $d = 4$. From the above table, we learn that the number of supersymmetries in six dimensions is 4, as the holonomy of T^4 is trivial. If we do not take into account the R-R fields, the moduli of the string σ -model is exactly the one described in section 3.1,

$$O(4, 4; \mathbf{Z}) \backslash O(4, 4) / O(4) \times O(4), \quad (3.107)$$

with the T -duality $O(4, 4; \mathbf{Z})$ corresponding to changes of the type $R_i \rightarrow \frac{\alpha'}{R_i}$, for the four S^1 cycles composing the torus. The situation becomes different if we allow R-R background fields. In such a case, we should take into account the possibility of including Wilson lines for the A_μ field (the **8_V** in (3.102)), and also a background for the 3-form $A_{\mu\nu\rho}$ (the **56_V** of (3.102)). The number of Wilson lines is certainly 4, one for each non contractible loop in T^d , so we need to add 4 dimensions to the 16-dimensional space (3.107). Concerning an $A_{\mu\nu\rho}$ background, the corresponding moduli is determined by $H_3(T^4)$, which implies 4 extra parameters. Finally, the dimension equals

$$16 + 4 + 4 = 24. \quad (3.108)$$

Now, a new extra dimension coming from the dilaton field must be added. It is important here to stress this fact: in the approach in previous section to σ -model moduli space the dilaton moduli has not been considered. This corresponds to interpreting the dilaton as a string coupling constant, and allowing changes only in the string. Anyway, this differentiation is rather cumbersome. Adding the dilaton moduli to (3.108), we get a moduli space of dimension (3.25), that can be written as

$$O(5, 5; \mathbf{Z}) \backslash O(5, 5) / O(5) \times O(5). \quad (3.109)$$

The proposal of moduli (3.109) for type IIA on T^4 already contains a lot of novelties. First of all, the modular group $O(5, 5; \mathbf{Z})$ now acts on the dilaton and the resting Ramond fields. In fact, relative to the $O(4, 4; \mathbf{Z})$ T -duality of toroidal compactifications, we have now an extra symmetry which is S -duality [5, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86],

$$g \rightarrow \frac{1}{g}, \quad (3.110)$$

with g the string coupling constant. This new modular symmetry is called in the physics literature U -duality [78]. The phenomena found here resembles very much what arises from mirror symmetry in the analysis of $K3$. There, the “classical” modular group was $O(\Gamma^{3,19}; \mathbf{Z})$, and quantum mirror symmetry creates the enhancement to $O(\Gamma^{4,20}; \mathbf{Z})$ where, in addition to T -duality, we have mirror transformations. In the case of type IIA on T^4 , it is because we include the R-R backgrounds and the dilaton that the modular symmetry $O(4, 4; \mathbf{Z})$ is enhanced to the U -duality symmetry. In spite of the analogies, the physical meaning is different. To appreciate this, let us now consider type IIA on $K3$. The dilaton moduli can be added, but the R-R fields are not producing any new moduli. In fact, recall that $\Pi_1(K3) = 0$, and $H_3 = 0$, so the moduli of type IIA on $K3$ is simply

$$O(4, 20; \mathbf{Z}) \backslash O(4, 20) / O(4) \times O(20) \times \mathbf{R}, \quad (3.111)$$

with $\mathbf{R}$ parametrizing the dilaton, and the modular group not acting on it.

The way to interpret the moduli (3.109) goes under the name of M-theory. Before entering a more precise definition of M-theory, the basic idea is thinking of (3.109) simply as the moduli of a toroidal compactification on T^5 ; however, in order to obtain a six dimensional $N=4$ theory, we need to start with some theory living in 11 dimensions. The theory satisfying this is M-theory, a theory whose low energy supergravity description is well understood: it should be such that through standard Kaluza-Klein compactification it gives the field theory limit of type IIA strings; but this a theory known as eleven dimensional type IIA supergravity.

Once we have followed the construction of the type IIA string theory moduli on T^4 , let us consider the general case of compactification on T^d . The dimension of the moduli is

$$\dim = d^2 + 1 + d + \frac{d(d-1)(d-2)}{3}, \quad (3.112)$$

where d^2 is the NS-NS contribution, the 1 sumand comes from the dilaton, d from the Wilson lines, and $\frac{d(d-1)(d-2)}{3}$ from the 3-form $A_{\mu\nu\rho}$. The formula (3.112) has to be completed, for $d \geq 5$, by including dual scalars. For $d = 5$, the dual to the 3-form $A_{\mu\nu\rho}$ is a scalar. The result is

$$\begin{aligned} \frac{d(d-1)(d-2)(d-3)(d-4)}{5} & \text{ duals to } A_{\mu\nu\rho}, \\ \frac{d(d-1)\dots(d-6)}{7} & \text{ duals to } A_\mu. \end{aligned} \quad (3.113)$$

The moduli spaces, according to the value of the dimension of the compactification torus, are listed in the table below.

Dimension	Moduli
$d = 4$	$O(5, 5; \mathbf{Z}) \backslash O(5, 5) / O(5) \times O(5)$
$d = 5$	$E_{6,(6)}(\mathbf{Z}) \backslash E_{6,(6)} / Sp(4)$
$d = 6$	$E_{7,(\tau)}(\mathbf{Z}) \backslash E_{7,(\tau)} / SU(8)$
$d = 3$	$Sl(5, \mathbf{Z}) \backslash Sl(5) / SO(5)$
$d = 2$	$Sl(3, \mathbf{Z}) \times Sl(2, \mathbf{Z}) \backslash Sl(3) / SO(3) \quad Sl(2) / SO(2)$

For supergravity practitioners, the appearance of E_6 and E_7 in this table should not be a surprise.

Let us now see what happens in the type IIB case. The moduli on, for instance, T^4 , is again the 16 dimensional piece coming from the NS-NS sector; now, the R-R sector is determined by the cohomology groups H^0 , H^2 and H^4 (see equation (3.101)). From the Hodge diamond for T^4 ,

$$\begin{array}{ccccc} & & 1 & & \\ & 2 & & 2 & \\ 1 & & 4 & & 1 \\ & 2 & & 2 & \\ & & 1 & & \end{array} \quad (3.114)$$

we get 8 extra moduli, exactly the same number as in the type IIA case. This is a general result for any T^d compactification. The reason for this is that type IIA and type IIB string theories are, after toroidal compactification, related by T -duality. However, on a manifold as $K3$, with $\Pi_1 = 0$, the moduli for IIA and IIB are drastically different, as

can be derived from direct inspection of the $K3$ Hodge diamond (see equation (3.39)). Therefore, for type IIB we get, from the R-R sector, 1 coming from H^0 , 22 from H^2 , and 1 from H^4 , which sums up a total of 24 extra moduli to be added to the $58 + 22$ of the NS-NS sector. Then, including the dilaton,

$$\dim \text{IIB}(K3) = 22 + 58 + 24 + 1 = 105. \quad (3.115)$$

Therefore, the natural guess for the moduli is

$$O(5, 21; \mathbf{Z}) \backslash O(5, 21) / O(5) \times O(21). \quad (3.116)$$

Here, something quite surprising is taking place. As we can see from (3.111), when type IIA is compactified on $K3$, we do not find any appearance of U -duality or, in other words, S -duality. By contrast, in the type IIB case we find a modular group $O(5, 2; \mathbf{Z})$, that contains the dilaton and, therefore, the S -duality transformation. This is what can be called the S -duality of type IIB string theory [87], which can already be observed from equation (3.101). In fact, the R-R and NS-NS sectors both contain scalar fields and the antisymmetric tensor.

3.2.2 Heterotic String.

The idea of “heterosis”, one of the most beautiful and productive ideas in the recent history of string theory [88] was motivated by two basic facts. First of all, the need to find a natural way to define non abelian gauge theories in string theory, without entering the use of Chan-Paton factors, and, secondly, the sharpness of the gap in string theory between left and right moving degrees of freedom. Here, we will concentrate on some of the ideas leading to the construction of heterosis. In the toroidal compactification of the bosonic string on T^d , we have found that the momenta live in a $\Gamma^{d,d}$ lattice. This is also true for the NS sector of the superstring. The lattice $\Gamma^{d,d}$, where the momenta live, is even and self dual. Taking into account the independence between left and right sectors, we can think on the possibility to compactify the left and right components on different tori, T^{d_L} and T^{d_R} , and consider as the corresponding moduli the manifold

$$O(d_L, d_R; \mathbf{Z}) \backslash O(d_L, d_R) / O(d_L) \times O(d_R). \quad (3.117)$$

Before trying to find out the consistency of this picture, let us try to get a simple interpretation of moduli (3.117). The dimension of this moduli is $d_L \times d_R$, and we can separate it into $d_L \times d_L + d_L \times (d_R - d_L)$. Let us interpret the first part, $d_L \times d_L$, as

the standard moduli for compactifications on a torus T^{d_L} ; then, the second piece can be interpreted as the moduli of Wilson lines for a gauge group

$$U(1)^{d_R-d_L}. \quad (3.118)$$

With this simple interpretation, we already notice the interplay in heterosis when working with a gauge group that can be potentially non abelian, the gauge group (3.118), and differentiating left and right parts. When we were working with type II string theory, and considered toroidal compactifications, we were also adding, to the moduli space, the contribution of the Wilson lines for the RR gauge field, A_μ (in case we are in type IIA). However, in the case of type IIA on T^4 , taking into account the Wilson lines did not introduce any heterosis asymmetry in the moduli of the kind (3.117). However, T^4 is not the only Ricci flat four dimensional manifold; we can also consider $K3$ surfaces. It looks like if T^4 , $K3$, and its orbifold surface in between, $T^4/\mathbf{Z}_2$, saturate all compactification manifolds that can be thought in four dimensions. In the case of $K3$, the moduli of type IIA string (see equation (3.111)) really looks like the heterotic moduli, of the kind (3.117), we are looking for. Moreover, in this case, and based on the knowledge of the lattice of the second cohomology group of $K3$ (see equation (3.44)),

$$E_8 \perp E_8 \perp \mathcal{U} \perp \mathcal{U} \perp \mathcal{U}, \quad (3.119)$$

we can interpret the $16 = d_R - d_L$ units as corresponding precisely to Wilson lines of the $E_8 \times E_8$ gauge group appearing in (3.119). In other words, and following a very distant path from the historical one, what we are suggesting is interpreting moduli (3.111), of type IIA on $K3$, as some sort of heterosis, with $d_L = 4$ and $d_R = 20$. The magic of numbers is in fact playing in our team, as the numbers we get for d_L and d_R strongly suggest a left part, of critical dimension 10, and a right part, of precisely the critical dimension of the bosonic string, 26. This was, in fact, the original idea hidden under heterosis: working out a string theory looking, in its left components, as the standard superstring, and in its right components as the 26 dimensional bosonic string. However, we are still missing something in the “heterotic” interpretation of (3.111), which is the visualization, from $K3$ geometry, of the gauge group. In order to see this, some of the geometrical material introduced in subsection 3.1.5 will be needed; in terms of the concepts there introduced, we would claim that the (p_L, p_R) momentum is living in the lattice $\Gamma^{4,20}$. We can then think that p_L is in the space-like 4-plane where the holomorphic top form Ω , and the Kähler class J , are included. Recall that they define a space-like 3-plane. Now,

momentum vectors, orthogonal to this 4-plane, can be considered; they are of the type

$$(0, p_R). \quad (3.120)$$

Now, whenever $p_R^2 = -2$, this vector will define a rational curve inside $K3$, with vanishing volume (in fact, the volume is given by $p_R \cdot J = 0$). The points $p_R^2 = -2$ will be at the root lattice of $E_8 \times E_8$. Now, from the mass formulas (3.26) we easily observe that $p_R^2 = -2$ is the condition for massless vector particles. In fact, if we separate, in the spirit of heterosis, the p_R of a 26 dimensional bosonic string into $(p_R^{(16)}, p_R^{(10)})$, we get, from (3.26),

$$M^2 = 4(p_R^{(16)})^2 + 8(N - 1), \quad (3.121)$$

so that $M^2 = 0$, for $N = 0$, if $(p_R^{(16)})^2 = 2$. The sign difference appears here because (recall subsection 3.1.5) in the $K3$ construction used for the second cohomology lattice, the E_8 lattice was defined by minus the Cartan algebra of E_8 . Therefore, we observe that massless vector bosons in heterotic string are related to rational curves in $K3$ of vanishing volume, which allows to consider enhancement of symmetries when moving in moduli space [81, 89, 90]. Some of these rational curves can be blown up, which would be the geometrical analog of the Higgs mechanism, or either blown down, getting extra massless stuff. Moreover, for elliptically fibered $K3$ surfaces, the different Kodaira singularities reflect, in its Dynkin diagram, the kind of gauge symmetry to be found.

The above discussion summarizes what can be called the first quasi-theorem on string equivalence [78, 81],

Quasi-Theorem 1 Type IIA string on $K3$ is equivalent to $E_8 \times E_8$ heterotic string on T^4 .

Previous arguments were so general that we can probably obtain extra equivalences by direct inspection of the different $K3$ moduli spaces that have been discussed in subsection 3.1.5. In particular, let us consider the moduli space of complex structures for an elliptically fibered $K3$ surface, a fact represented, in terms of the Picard lattice, claiming that it is of $\Gamma^{1,1}$ type, generated by a section, and with the fiber satisfying relations (3.28). This moduli is

$$O(2, 18; \mathbf{Z}) \backslash O(2, 18) / O(2) \times O(18), \quad (3.122)$$

where we have used equation (3.62), and the fact that the transcendental lattice is of type $(2, 18)$. From the heterosis point of view, it would be reasonable to interpret (3.122) as heterotic $E_8 \times E_8$ string, compactified on a 2-torus, T^2 . In fact, we will have 4 real

moduli, corresponding to the Kähler class and complex structure of T^2 , and 16 extra complex moduli associated to the Wilson lines. However, now the type II interpretation of (3.122) is far from being clear, as (3.122) is just the part of the moduli space that is preserving the elliptic fibration. Now, in order to answer how (3.122) can be understood as a type II compactification a similar problem appears as we try to work out an heterotic interpretation of the type IIB moduli on $K3$, given in (3.116). A simple way to try to interpret (3.122), as some kind of type II compactification, is of course thinking of an elliptically fibered $K3$, where the volume of the fiber is fixed to be equal zero; generically,

$$J \cdot F = 0, \quad (3.123)$$

where F indicates the class of the fiber. Now, we can think that we are compactifying a type II string on the base space of the bundle. However, this does not lead to (3.122) for the type IIA case, as the RR fields are in H^1 and H^3 , which will vanish. But what about type IIB? In this case, we have the NS field ϕ , and the R field χ , and we should fix the moduli of possible configurations of these fields on the base space of the elliptic fibration. Here, type IIB S -duality, already implicit in moduli (3.116), can help enormously, mainly because we are dealing with an elliptically fibered $K3$ manifold [91, 92, 93]. To proceed, let us organize the fields ϕ and χ into the complex

$$\tau = \chi + ie^{-\phi}, \quad (3.124)$$

and identify this τ with the moduli of the elliptic fiber. Then, the 18 complex moduli dimension of (3.122) parametrizes the moduli of complex structures of the elliptic fibration, and therefore the moduli of τ field configurations on the base space (provided τ and $\frac{a\tau+b}{c\tau+d}$ are equivalent from the type IIB point of view). These moduli parametrize then the type IIB compactification on the base space B (it is $\mathbb{P}^1$; recall that in deriving (3.122) we have used a base space B such that $B \cdot B = -2$). There is still one moduli missing: the size of the base space B , that we can identify with the heterotic string coupling constant. Thus, we arrive to the following quasi-theorem,

Quasi-Theorem 2 Heterotic string on T^2 is equivalent to type IIB string theory on the base space of an elliptically fibered $K3$.

The previous discussion is known, in the physics literature, under the generic name of F-theory [94, 95, 96].

We have been considering, until now, type II strings on $K3$, and compared them to heterotic string on a torus. To find out what is the expected moduli for the heterotic string

on $K3$, we can use the following trick: if heterotic string on T^2 is type IIB on the base space of an elliptically fibered $K3$, by quasi-theorem 2 heterotic string on an elliptically fibered $K3$ should correspond to type IIB on the base space of an elliptically fibered Calabi-Yau manifold. More precisely, type IIB string should be compactified on the basis of an elliptic fibration, which is now four dimensional, and that can be represented as a fibration of a $\mathbb{P}^1$ space over another $\mathbb{P}^1$. This type of fibrations are known in the literature as Hirzebruch spaces, $\mathbf{F}_n$. Hirzebruch spaces can simply be determined through heterotic data, given by the $E_8 \times E_8$ bundle on the $K3$ manifold. The moduli of these bundles on $K3$ will put us in contact with yet another interesting topic: small instantons.

3.2.3 Heterotic Compactifications to Four Dimensions.

Before considering some definite examples, let us simply summarize the different supersymmetries we can get when compactifying to three dimensions, depending on the holonomy of the target manifold. In order to do that, we will need the results in subsection 3.2.1, on the maximum number of supersymmetries allowed for a given spacetime dimension.

Type of String	Target Manifold	Holonomy	Supersymmetry
II	$K3 \times T^2$	$SU(2)$	$N=4$
Heterotic	T^6	Trivial	$N=4$
II	Calabi-Yau	$SU(3)$	$N=2$
Heterotic	$K3 \times T^2$	$SU(2)$	$N=2$
II	$B_{SU(4)}$	$SU(4)$	$N=1$
Heterotic	Calabi-Yau	$SU(3)$	$N=1$

In the table above we have not differentiated between type IIA and type IIB¹⁶. The first two lines, corresponding to cases with $N=4$ and $N=2$ supersymmetry in four dimensional spacetime, will be the basic examples we will use to introduce the concept of dual pairs of string compactifications down to four dimensions.

Before entering a discussion on the ingredients of this table, we yet need to consider the holonomy of the moduli space. This holonomy will of course depend on the number of supersymmetries and the type (real, complex or quaternionic) of the representation. Hence, from subsection 3.2.1, we can complete the table below.

¹⁶This will be relevant when discussing the third line where, by $B_{SU(4)}$, we are thinking in the spirit of the discussion in the last part of previous section, where a Calabi-Yau fourfold of $SU(4)$ holonomy, elliptically fibered, and with a zero volume fiber, is used for compactification.

Spacetime Dimension	Supersymmetries	Type	Holonomy
$d = 6$	$N = 2$	$\mathbf{H}^4$	$Sp(1) \oplus Sp(1)$
$d = 4$	$N = 4$	$\mathbf{C}^2$	$U(4)$
$d = 4$	$N = 2$	$\mathbf{C}^2$	$U(2)$

Using this results, we can now decompose the tangent vectors to the moduli according to its transformation rules with respect to the holonomy group. Let us concentrate in the $d = 4$ case. For $U(4)$, we get

$$U(4) \simeq U(1) \oplus SO(6). \quad (3.125)$$

The matter multiplets will contain 6 (real) scalars each, i. e., the number of dimensions we compactify. Then, if we have m of these matter multiplets, the part of the moduli on which the $SO(6)$ part of the holonomy group is acting should be

$$O(6, m)/O(6) \times O(m). \quad (3.126)$$

The $U(1)$ part of (3.125) will act on the supergravity multiplet so we expect, just from holonomy arguments, a moduli of type

$$O(6, m)/O(6) \times O(m) \times Sl(2)/U(1). \quad (3.127)$$

Now, we need to compute m . For heterotic string, the answer is clear: $m = 22$, and the total dimension of (3.127) will be 134. Let us now consider the case of type IIA. From the table, we see that we should consider $K3 \times T^2$ as compactification manifold. Let us then first compute the dimension of the moduli space:

$$\begin{aligned}
\text{Moduli of metrics and B fields on } K3 &= 80 \\
\text{Moduli of metrics and B fields on } T^2 &= 4 \\
b_1(K3 \times T^2) &= 2 \\
b_3(K3 \times T^2) &= 44 \\
\text{Axion-Dilaton} &= 2 \\
\text{Duals in } \mathbf{R}^4 \text{ to 2-forms} &= 2
\end{aligned} \quad (3.128)$$

which sums up to 134. Notice that the 44 in $b_3(K3 \times T^2)$ is coming from the 3-cycles obtained from one S^1 of T^2 , and the 22 elements in $H^2(K3; \mathbf{Z})$. The 3-form of IIA can be compactified on the S^1 cycles of T^2 to give 2-forms in four dimensions. Now, the dual of a 2-form in $\mathbf{R}^4$ is scalar, so we get the last two extra moduli.

Now, we need to compare the two moduli spaces. If we expect S -duality in $N=4$ for the heterotic compactification, the moduli, once we have taken into account the $O(6, 22; \mathbf{Z})$ T -duality, will look like

$$O(6, 22; \mathbf{Z}) \backslash O(6, 22) / O(6) \times O(22) \quad Sl(2, \mathbf{Z}) \backslash Sl(2) / U(1). \quad (3.129)$$

Now, we have a piece in IIA looking naturally as the second term in (3.129), namely the moduli of the σ -model on T^2 , where $Sl(2, \mathbf{Z})$ will simply be part of the T -duality. Thus, it is natural to relate the moduli of IIA on the torus with the part of the moduli in (3.127) coming from the supergravity multiplet.

Let us now consider dual pairs in the second line of our table. There is a simple way to visualize under what general conditions on the Calabi-Yau manifold with $SU(3)$ holonomy such dual pairs can exist. In fact, imagine that $K3$ is elliptically fibered in $K3 \times T^2$; then, what we get is a fibration on $\mathbb{P}^1$ of the T^4 tori. Now, heterotic on T^4 is equivalent to type IIA on $K3$, so we expect that the Calabi-Yau manifold should be a $K3$ fibration on $\mathbb{P}^1$, and that duality works fiberwise. Therefore, from general arguments, we expect to get heterotic-type II dual pairs with $N = 2$ if we use Calabi-Yau manifolds which are $K3$ fibrations [97, 93]. In order to get a more precise picture, we need again to work out the holonomy, which is $U(2)$ in this case. In $N = 2$ we have two types of multiplets, vector and hypermultiplets. The vector multiplet contains two real scalars, and the hypermultiplet four real scalars. Then, we decompose $U(2)$ into $U(1) \oplus Sp(1)$, and the moduli into vector and hypermultiplet part.

Let us first consider type IIA string on the Calabi-Yau manifold. The moduli will contain $h^{1,1}$ deformations of B and J , $h^{2,1}$ complex deformations and b^3 RR deformations (b^1 does not contribute, as we are working with a Calabi-Yau manifold). The total number, in real dimension, is

$$2h^{1,1} + 4(h^{2,1} + 1), \quad (3.130)$$

where we have used that $b^3 = 2(h^{2,1} + 1)$, in real dimension. From (3.130) we conclude that we have $h^{1,1}$ vector multiplets, and $h^{2,1} + 1$ hypermultiplets. Notice that $4(h^{2,1} + 1)$ is counting the 2 coming from the dilaton and the axion so, for type II we have combined dilaton and axion into an hypermultiplet.

Now, let us consider heterotic string on $K3 \times T^2$. The moduli we must now consider, of $E_8 \times E_8$ bundles on $K3$, is much more elaborated than that of T^4 , or T^6 , that we have worked out. Part of the difficulty comes from anomaly conditions. However, we know, according to Mukai's theorem, that the moduli of holomorphic bundles on $K3$ is quaternionic, i. e., hyperkähler, and that the moduli of the σ -model on $K3$ is of dimension

80. We have yet the moduli on T^2 , that will be a manifold of $O(2, m)/O(2) \times O(m)$ type, and therefore a good candidate for representing the vector multiplet. Thus, we get

$$\begin{aligned} \text{Type IIA hypermultiplets} &\leftrightarrow K3 \text{ Heterotic,} \\ \text{Vector multiplets} &\leftrightarrow T^2. \end{aligned} \quad (3.131)$$

From our previous discussion we know that vector multiplets, in type IIA are related to $h^{1,1}$. Working fiberwise on a $K3$ fibered Calabi-Yau manifold we get, for $h^{1,1}$,

$$h^{1,1} = 1 + \rho, \quad (3.132)$$

with ρ the Picard number of the $K3$ manifold. Then, in order to get a dual pair in the sense of (3.131) we need m in the heterotic to satisfy

$$m = \rho. \quad (3.133)$$

In order to control the value of m , from the heterotic point of view, we need to watch out for possible Wilson lines that can be defined on T^2 after the gauge group has been fixed from the $K3$ piece. From (3.132) (and this was the logic for the identification (3.133)), the heterotic dilaton-axion is related to the 1 term contributing in (3.132), i. e., the 2-cycle defined by the base space of the $K3$ -fibration.

As can be observed from (3.133), if we do not freeze either the Kähler class or the complex structure of T^2 , the minimum value for ρ is 2. This is the contribution to the Picard lattice of a Dynkin diagram of type A_2 , i. e., $SU(3)$. A possible line of work opens here, in order to identify the moduli spaces of vector multiplets for type IIA theories with the quantum moduli, defined according to Seiberg and Witten, for gauge theories, with

$$\text{rank } G = \rho. \quad (3.134)$$

4 Chapter IV

4.1 M-Theory Compactifications.

Wittgenstein used to say that “meaning is use”. This is the kind of philosophical slogan able to make unhappy the platonic mathematician, but it is in essence the type of game we are going to play in order to begin the study of M-theory [98, 78, 80, 81, 86]. More precisely, we will start without saying what M-theory is from a microscopical point of view, giving instead a precise meaning to M-theory compactifications.

Recall that our first contact with the idea of M-theory was in connection with the interpretation of the moduli of type IIA string theory on T^4 . In that case the moduli, after including RR fields, was of the type

$$O(5, 5; \mathbf{Z}) \backslash O(5, 5) / O(5) \times O(5). \quad (4.1)$$

The M-theory interpretation of moduli (4.1) can be summarized according to the equivalence

$$\text{M-theory compactified on } T^5 \leftrightarrow \text{IIA on } T^4, \quad (4.2)$$

and therefore, more generically,

$$\text{M-theory compactified on } X \times S^1 \leftrightarrow \text{IIA on } X. \quad (4.3)$$

Let us now put rule (4.3) into work. In fact, one particular case of (4.3) will consist in considering M-theory on a manifold of type $B \times S^1 \times S^1$. Then, using T -duality, we can get

$$\begin{aligned} \text{M-theory compactified on } B \times S^1 \times S^1(R) &\leftrightarrow \\ \text{IIA on } B \times S^1(R) &\leftrightarrow \text{IIB on } B \times S^1\left(\frac{\alpha'}{R}\right). \end{aligned} \quad (4.4)$$

From (4.4) we see that in the $R \rightarrow \infty$ limit, we get type IIB string theory on B or, equivalently, M-theory on $B \times S^1$, since the second S^1 becomes uncompactified. This is in fact a very close example to the ones described in previous sections, under the generic name of F-theory compactifications. Namely, the $R \rightarrow \infty$ limit in (4.4) can be interpreted as defining a compactification of type IIB string theory on the base space B of an elliptic fibration $B \times S^1 \times S^1$, in the limit where the volume of the elliptic fiber becomes zero. Following that path, we get an interesting equivalence between M-theory on $B \times S^1 \times S^1$, as elliptic fibration, in the limit in which the volume of the elliptic fiber goes zero, and type IIB on B . This stands as a surprise, when compared to the result derived from the compactification rule (4.3). In fact, if B is, for instance, of dimension d , then we should expect that the compactification of an eleven dimensional theory, as M-theory, on $B \times S^1 \times S^1$, should lead to $11 - d - 2$ dimensions. However, type IIB, which is ten dimensional, would lead, when compactified on B , to a $10 - d$ dimensional theory, so that one dimension is missing. Getting rid off this contradiction requires knowledge of the microscopic nature of M-theory. The first thing to be required on M-theory is of course to have, as low energy limit, eleven dimensional supergravity. There is a connection between type IIA string theory and eleven dimensional supergravity, as the corresponding Kaluza-Klein dimensional reduction on an internal S^1 , which allows an identification of the string

theory spectrum with supergravity. In particular, the RR field in ten dimensions comes from the $g_{11,\mu}$ component of the metric, while the dilaton is obtained from $g_{11,11}$. The precise relation, in what is known as the string frame, is¹⁷

$$e^{-2\phi} = e^{-3\gamma}, \quad (4.5)$$

with ϕ the type IIA dilaton field. In terms of the radius R of the S^1 ,

$$R = e^{2\phi/3}. \quad (4.6)$$

Using now equation (3.21), we get a relation between the R of the internal manifold, S^1 , and the string coupling constant of type IIA string theory,

$$R = g^{2/3}. \quad (4.7)$$

From (4.7) it is obvious that, as $R \rightarrow \infty$, we properly enter the M-theory region when g is large, i. e., working in the strong coupling regime of string theory. Historically, this beautiful simple argument was put forward in 1995 by Witten [81]. It is astonishing that, with all the pieces around, nobody was able before to make at least the comment relating the R of eleven dimensional supergravity with the string coupling constant, and to derive from it such a striking conjecture as it is that strongly coupled IIA strings are described by eleven dimensional supergravity. In fact, there are good reasons for such a mental obstacle in the whole community: first of all, nobody did worry about type IIA dynamics, as it was a theory with only uninteresting pure abelian gauge physics. Secondly, the Kaluza-Klein modes coming from the compactification on S^1 , which have a mass of the order $\frac{1}{R}$, are charged with respect to the $U(1)$ gauge field defined by the $g_{11,\mu}$ piece of the metric. But this A_μ field in ten dimensional type IIA string is of RR type, so before the discovery of D-branes, there was no candidate in the string spectrum to be put in correspondence with these Kaluza-Klein modes, which can now be identified with D-0branes.

Witten's approach to M-theory can be the conceptual key to solve the problem concerning the missing dimension: in fact, something in the spectrum is becoming massless as the volume of the elliptic fiber, in the case of $B \times S^1 \times S^1$, is sent to zero. Moreover, the object becoming massless can be, as suggested by Sethi and Susskind, interpreted as a Kaluza-Klein mode of an opening dimension as the volume of the elliptic fiber goes zero. To understand the nature of this object we should look more carefully at M-theory. This theory is expected to contain a fundamental two dimensional membrane; if this membrane

¹⁷We have identified $g_{11,11} = e^{2\gamma}$.

wraps the 2-torus $S^1 \times S^1$, its mass becomes zero as the volume of the fiber goes zero. Then, all what is left is to relate the area with the standard Kaluza-Klein formula for compactifications on S^1 , which leads to

$$\frac{1}{R} \sim L_1 L_2 \quad (4.8)$$

solving our problem on the adequate interpretation of (4.4).

Let us now concentrate on a concrete example of (4.4): we will choose $X = B \times S^1 \times S_R^1$ as representing a Calabi-Yau fourfold of $SU(4)$ holonomy. After compactification, $SU(4)$ holonomy implies a three dimensional theory with $N = 2$ supersymmetry should be expected. Moreover, sending $R \rightarrow \infty$ leads to a four dimensional $N = 1$ theory. In order to work out the spectrum of the three dimensional theory, standard Kaluza-Klein techniques can be used. Compactification on the 2-cycles of $H^2(X; \mathbf{Z})$ of the 3-form $C_{\mu\nu\rho}$ of eleven dimensional supergravity leads to a vector in three dimensions. Moreover, the Kähler class can also be used to generate real scalars, from each 2-cycle. Thus, let us assume $r = \dim H^2(X; \mathbf{Z})$; then, the previous procedure produces r real scalars and r vector fields. In order to define r $N = 2$ vector multiplets in three dimensions, with these vector fields, another set of r scalars is yet needed, in order to build the complex fields. These extra r scalars can, as usual, be identified with the duals, in three dimensions, of the 1-form vector fields: the three dimensional dual photon.

Our next task will be reproducing, using M-theory, the well known instanton effects in $N = 2$ supersymmetric gauge theories in three dimensions.

4.2 M-Theory Instantons.

In order to define instantons in three dimensions, we will use 5-branes wrapped on the 6-cycles of a Calabi-Yau fourfold X [99]. The reason for using 6-cycles is understood as follows: the gauge bosons in three dimensions are obtained from the integration of the 3-form $C_{\mu\nu\rho}$ over 2-cycles. Thus, in order to define the dual photon, we should consider the dual, in the Calabi-Yau fourfold X , of 2-cycles, which are 6-cycles. However, not any 6-cycle can be interpreted as an instanton with topological charge equal one, and therefore no 6-cycle will contribute to the three dimensional superpotential.

If we interpret a 5-brane wrapped on a 6-cycle D of X as an instanton, we can expect a superpotential of the type

$$W = e^{-(V_D + i\phi_D)}, \quad (4.9)$$

with V_D the volume of D measured in units of the 5-brane tension, and ϕ_D the dual photon field, associated with the cycle D . In order to get, associated to D , a superpotential like (4.9), we need

- i) To define a $U(1)$ transformation with respect to which three dimensional fermions are charged.
- ii) To associate with the 6-cycle D a violation of $U(1)$ charge, in the adequate amount.
- iii) To prove that this $U(1)$ symmetry is not anomalous.
- iv) To interpret ϕ_D as the corresponding Goldstone boson.

Following these steps, we will extend to M-theory the instanton dynamics of three dimensional $N = 2$ gauge theories described in chapter I. We will start defining the $U(1)$ transformation. Let D be 6-cycle in the Calabi-Yau fourfold X , and let us denote by N the normal bundle of D in X . Since X is a Calabi-Yau manifold, its canonical bundle is trivial, and therefore we get

$$K_D \simeq N, \quad (4.10)$$

with K_D the canonical bundle of D . Locally, we can interpret X as the total space of the normal bundle. Denoting by z the coordinate in the normal direction, the $U(1)$ transformation can be defined as

$$z \rightarrow e^{i\theta} z. \quad (4.11)$$

The $U(1)$ transformation defined by (4.11) is very likely not anomalous, since it is part of the diffeomorphisms of the eleven dimensional theory; thus, it is a good candidate for the $U(1)$ symmetry we are looking for. Next, we need to get the $U(1)$ charge of the three dimensional fermions. However, before doing so, we will review some well known facts concerning fermions and Dirac operators on Kähler manifolds.

We will consider a Kähler manifold of complex dimension N . In holomorphic coordinates,

$$g_{ab} = g_{\bar{a}\bar{b}} = 0. \quad (4.12)$$

In these coordinates, the algebra of Dirac matrices becomes

$$\begin{aligned} \{\gamma^a, \gamma^b\} &= \{\gamma^{\bar{a}}, \gamma^{\bar{b}}\} = 0, \\ \{\gamma^a, \gamma^{\bar{b}}\} &= 2g^{a\bar{b}}. \end{aligned} \quad (4.13)$$

The $SO(2N)$ spinorial representations of (4.13) can be obtained in the standard Fock approach: a vacuum state is defined by condition

$$\gamma^a |\Omega\rangle = 0, \quad (4.14)$$

and n -particle states are defined by

$$\gamma^{\bar{a}} \gamma^{\bar{b}} \dots \gamma^{\bar{n}} |\Omega\rangle. \quad (4.15)$$

A spinor field $\psi(z, \bar{z})$ on the Kähler manifold takes values on the spinor bundle defined by this Fock representation:

$$\psi(z, \bar{z}) = \phi(z, \bar{z}) |\Omega\rangle + \phi_{\bar{a}}(z, \bar{z}) \gamma^{\bar{a}} |\Omega\rangle + \phi_{\bar{a}\bar{b}}(z, \bar{z}) \gamma^{\bar{a}} \gamma^{\bar{b}} |\Omega\rangle + \dots \quad (4.16)$$

The spaces $\Omega^{0,q}$ of $(0, q)$ -forms, generated by the Dirac operator, define the Dolbeault cohomology of the Kähler manifold. Using this notation, the two different chirality spinor bundles are

$$\begin{aligned} S^+ &= (K^{1/2} \otimes \Omega^{0,0}) \oplus (K^{1/2} \otimes \Omega^{0,2}) \oplus (K^{1/2} \otimes \Omega^{0,4}) \oplus \dots, \\ S^- &= (K^{1/2} \otimes \Omega^{0,1}) \oplus (K^{1/2} \otimes \Omega^{0,3}) \oplus (K^{1/2} \otimes \Omega^{0,5}) \oplus \dots, \end{aligned} \quad (4.17)$$

and the change of chirality (the index for the Dirac operator on the manifold X) will be given by the arithmetic genus,

$$\chi = \sum_{n=0}^N (-1)^n h_n, \quad (4.18)$$

where $h_n = \dim \Omega^{0,n}$.

The previous comments can be readily applied to the case of a six dimensional divisor D in a Calabi-Yau fourfold X . Now, we should take into account the normal bundle N , to D , in X . Using the fact that X is Calabi-Yau, i. e., with trivial canonical bundle, we conclude that N is isomorphic to K_D , the canonical bundle on D . The spinor bundle on N will be defined by

$$K_D^{1/2} \oplus K_D^{-1/2}. \quad (4.19)$$

In fact, in this case the complex dimension of N is one, and the vacuum and filled states have, respectively, $U(1)$ charges $\frac{1}{2}$ and $-\frac{1}{2}$. On the other hand, the spinor bundle on D will be defined by (4.17), with $K = K_D$. Thus, spinors on D are, up to the $SO(3)$ spacetime part, taking values in the positive and negative chirality bundles

$$\begin{aligned} \hat{S}^+ &= (K^{1/2} \oplus K^{-1/2}) \otimes [(K^{1/2} \otimes \Omega^{0,0}) \oplus (K^{1/2} \otimes \Omega^{0,2})], \\ \hat{S}^- &= (K^{1/2} \oplus K^{-1/2}) \otimes [(K^{1/2} \otimes \Omega^{0,1}) \oplus (K^{1/2} \otimes \Omega^{0,3})]. \end{aligned} \quad (4.20)$$

Now, we are interested in a change of $U(1)$ charge, with the $U(1)$ charge defined by the $\frac{1}{2}$ and $-\frac{1}{2}$ charges of the spinor bundle (4.19) on N .

For spinors of a given chirality, the change of $U(1)$ charge is given by

$$\dim (K \otimes \Omega^{0,0}) + \dim (K \otimes \Omega^{0,2}) - \dim (\Omega^{0,0}) - \dim (\Omega^{0,2}). \quad (4.21)$$

Using now Serre's duality,

$$\dim (K \otimes \Omega^{0,3-n}) = \dim (\Omega^{0,n}), \quad (4.22)$$

we get that the number of holomorphic $(0, k)$ -forms is equal to the number of holomorphic sections in $K \otimes \Omega^{0,3-k}$, and therefore the number of fermionic zero modes with $U(1)$ charge equal $\frac{1}{2}$ is given by $h_3 + h_1$, and the number of fermionic zero modes with $-\frac{1}{2}$ $U(1)$ charge, is given by $h_0 + h_2$ (here we have used the Dirac operator $\bar{\partial} + \bar{\partial}^*$, with $\bar{\partial}^*$ the adjoint of $\bar{\partial}$). Thus, the index for the twisted spin bundle (4.20) is given by the holomorphic Euler characteristic,

$$\chi(D) = h_0 - h_1 + h_2 - h_3. \quad (4.23)$$

Now, each of these fermionic zero modes is doubled once we tensor with spinors in $\mathbf{R}^3$. In summary, for each 6-cycle D we get an effective vertex with a net change of $U(1)$ charge equal to $\chi(D)$.

Therefore, in order to get the three dimensional in a three dimensional $N = 2$ theory, we need to look for 6-cycles D , with $\chi(D) = 1$, as the net change of $U(1)$ charge in that case is one, provided, as we did in (4.19), we normalize the $U(1)$ charge of the fermions to be $\frac{1}{2}$. More precisely, the number of fermionic zero modes for a three dimensional instanton, defined by a 6-cycle D , is $2\chi(D)$.

4.3 D-Brane Configurations in Flat Space.

We will consider a D-brane of dimension p , in flat ten dimensional Minkowski space, and with a flat $p + 1$ dimensional worldvolume. The quantization of the open superstring ending on the D-brane defines a low energy field theory, which is ten dimensional $N = 1$ supersymmetric Yang-Mills, with $U(1)$ gauge group. The dimensional reduction of this theory to $p + 1$ dimensions describes the massless excitations propagating on the worldvolume of the p dimensional D-brane. We will use as worldvolume coordinates $x^0, x^1, \dots, x^p$. The worldvolume lagrangian will contain a $U(1)$ massless gauge field $A_i(x_s)$, with $i, s = 0, \dots, p$, and a set of scalar fields $\phi_j(x_s)$, $j = p + 1, \dots, 9$, transforming in the adjoint representation. We can geometrically interpret the set of fields $\phi_j(x_s)$

as representing the “location” of the flat D-brane in transverse space. The simplest generalization of the previous picture corresponds to configurations of $k > 1$ parallel D- p branes. In this case we have, in addition to the massless excitations, a set of k massive excitations corresponding to open strings ending on different D-branes.

The field theory interpretation of this configuration of D-branes would be that of a gauge theory with $U(k)$ gauge group, spontaneously broken to $U(1)^k$, with the strings stretching between different D-branes representing charged massive vector bosons. To get such an interpretation, we can start with $N = 1$ $U(k)$ supersymmetric Yang-Mills in ten dimensions, and perform again dimensional reduction down to $p + 1$ dimensions. In this case, we will get a set of scalar fields, $X^j(x_s)$, with $j = p + 1, \dots, 9$, which are now $k \times k$ matrices, transforming in the adjoint of $U(k)$. Moreover, the kinetic term in ten dimensions produces a potential of the form

$$V = \frac{g^2}{2} \sum_{i,j=p+1}^9 \text{tr}[X^i, X^j]^2. \quad (4.24)$$

As we have already observed in many examples before, this potential possesses flat directions, corresponding to classical vacuum states. These flat directions are defined by diagonal X^i matrices,

$$X^i = \begin{pmatrix} \lambda_1^i & & \\ & \ddots & \\ & & \lambda_k^i \end{pmatrix}. \quad (4.25)$$

On each of these vacua, the $U(k)$ gauge symmetry is spontaneously broken to $U(1)^k$; thus, we can use these vacuum configurations to describe sets of k parallel p dimensional D-branes. In fact, as we observe for the simpler case of one D-brane, the set of scalars appearing by dimensional reduction has the geometrical interpretation of the position of the D-brane. In the case (4.25), we can in fact consider λ_l^i as defining the i^{th} -coordinate of the l^{th} -brane. This is consistent with the idea of interpreting the strings stretching between different D-branes as massive vector bosons. In fact, the mass of this string states would be

$$M \sim g|\lambda_l - \lambda_m|, \quad (4.26)$$

for a (l, m) string. This is, in fact, the Higgs mass of the corresponding massive charged boson. In summary, merging the previous comments into a lemma: the classical moduli space of the worldvolume lagrangian of a D-brane coincides with its transversal space. It is important realizing that only the minima of the potential (4.24), i. e., the moduli space of the worldvolume lagrangian, is the one possessing this simple geometrical interpretation.

In particular, the dimensional reduction, down to $p + 1$ dimensions, of $N = 1$ $U(k)$ gauge theory, describes a set of k parallel branes, but its full fledged dynamics is described by the complete matrix X^i , with non vanishing off diagonal terms. A nice way to think about the meaning of (4.25) is again in terms of ‘t Hooft’s abelian projection. In fact, we can think of (4.25) as a unitary gauge fixing, where we now allow λ_l^i to depend on the worldvolume coordinates. The case of flat parallel D-branes corresponds to a Higgs phase, with λ_l^i constant functions on the worldvolume. Moreover, we can even consider the existence of singularities, which will be points where two eigenvalues coincide,

$$\lambda_l^i = \lambda_{l+1}^i, \quad \forall i. \quad (4.27)$$

It is quite obvious realizing that (4.27) imposes three constraints, so we expect, for p -dimensional D-branes, that on a $p - 3$ -dimensional region of the worldvolume, two consecutive D-branes can overlap. The $p - 3$ region in the $p + 1$ dimensional worldvolume of the D-brane will represent, from the point of view of $p + 1$ dynamics, a monopole, in the very same sense as is the case in ‘t Hooft’s abelian projection.

Next we will consider some brane configurations for type IIA and type IIB string theory (some of the widely increasing references are those from [100] to [117]). In order to define these configurations we will first work out the allowed vertices for intersecting branes. Let us start with a vertex of type $(p, 1^F)$, corresponding to a Dirichlet p -brane, and a fundamental string ending on the D-brane worldvolume. In type IIA p should be even, and odd for type IIB. In fact, the RR fields for type IIA and type IIB string theory are

$$\begin{aligned} \text{IIA} &\rightarrow A_\mu A_{\mu\nu\rho}, \\ \text{IIB} &\rightarrow \chi B_{\mu\nu} A_{\mu\nu\rho\sigma}. \end{aligned} \quad (4.28)$$

The corresponding strength tensors are, respectively, two and four-forms for type IIA, and one, three and five-forms for type IIB. Thus, the sources are D-branes of dimensions zero and two, for type IIA string theory, and one and three for type IIB. In addition, we have the (Hodge) magnetic duals, which are six and four D-branes for type IIA string theory, and five and three D-branes for type IIB (notice that the threebrane in type IIB is self dual). Besides, for the χ field in type IIB, the source is a -1 extended object, and its dual is a D-7brane.

Let us then start with a vertex of type $(p, 1^F)$ in type IIB, i. e., with p odd. We can use the $Sl(2, \mathbf{Z})$ duality symmetry of type IIB strings to transform this vertex into a

$(p, 1)$ vertex, between a D- p brane and a D-1brane, or D-string. By performing j T-duality transformations on the spacetime directions orthogonal to the worldvolume of the D-brane and the D-string, we pass from $(p, 1^F)$ to a vertex $(p + j, 1^F + j)$ of two D-branes, sharing j common worldvolume coordinates. If j is even, we end up with a vertex in type IIB, and if j is odd with a vertex in type IIA. Namely, through a T-duality transformation we pass from type IIB string theory to type IIA. As an example, we will consider the vertex $(3, 1^F)$ in type IIB string theory. After a S-duality transformation in the $Sl(2, \mathbf{Z})$ duality group of type IIB strings, and two T-duality transformations, we get the vertex $(5, 3)$ for branes. As we are in type IIB, we can perform a duality transformation on it to generate the vertex $(5^{NS}, 3)$, between the solitonic Neveu-Schwarz fivebrane and a D-3brane.

Let us now consider some brane configurations build up using the vertices $(5, 3)$ and $(5^{NS}, 3)$ in type IIB theory [100]. In particular, we will consider solitonic fivebranes, with worldvolume coordinates x^0, x^1, x^2, x^3, x^4 and x^5 , located at some definite values of x^6, x^7, x^8 and x^9 . It is convenient to organize the coordinates of the fivebrane as $(x^6, \vec{\omega})$ where $\vec{\omega} = (x^7, x^8, x^9)$. By construction of the vertex, the D-3brane will share two worldvolume coordinates, in addition to time, with the fivebrane. Thus, we can consider D-3branes with worldvolume coordinates x^0, x^1, x^2 and x^6 . If we put a D-3brane in between two solitonic fivebranes, at x_2^6 and x_1^6 positions in the x^6 coordinate, then the worldvolume of the D-3brane will be finite in the x^6 direction (see Figure 1).




Figure 1: Solitonic fivebranes with n Dirichlet threebranes stretching along them.

Therefore, the macroscopic physics, i. e., for scales larger than $|x_2^6 - x_1^6|$, can be effectively described by a $2 + 1$ dimensional theory. In order to unravel what kind of $2 + 1$ dimensional theory, we are obtaining through this brane configuration, we must

first work out the type of constraint imposed by the fivebrane boundary conditions. In fact, the worldvolume low energy lagrangian for a D-3brane is a $U(1)$ gauge theory. Once we put the D-3brane in between two solitonic fivebranes we impose Neumann boundary conditions, in the x^6 direction, for the fields living on the D-3brane worldvolume. This means in particular that for scalar fields we impose

$$\partial_6 \phi = 0 \quad (4.29)$$

and, for gauge fields,

$$F_{\mu 6} = 0, \quad \mu = 0, 1, 2. \quad (4.30)$$

Thus, the three dimensional $U(1)$ gauge field, A_μ , with $\mu = 0, 1, 2$, is unconstrained which already means that we can interpret the effective three dimensional theory as a $U(1)$ gauge theory for one D-3brane, and therefore as a $U(n)$ gauge theory for n D-3branes. Next, we need to discover the amount of supersymmetry left unbroken by the brane configuration. If we consider Dirichlet threebranes, with worldvolume coordinates x^0, x^1, x^2 and x^6 , then we are forcing the solitonic fivebranes to be at positions $(x^6, \vec{\omega}_1)$ and $(x^6, \vec{\omega}_2)$, with $\vec{\omega}_1 = \vec{\omega}_2$. In this particular case, the allowed motion for the D-3brane is reduced to the space $\mathbf{R}^3$, with coordinates x^3, x^4 and x^5 . These are the coordinates on the fivebrane worldvolume where the D-3brane ends. Thus, we have defined on the D-3brane three scalar fields. By condition (4.29), the values of these scalar fields can be constrained to be constant on the x^6 direction. What this in practice means is that the two ends of the of the D-3brane have the same x^3, x^4 and x^5 coordinates. Now, if we combine these three scalar fields with the $U(1)$ gauge field A_μ , we get an $N = 4$ vector multiplet in three dimensions. Therefore, we can conclude that our effective three dimensional theory for n parallel D-3branes suspended between two solitonic fivebranes (Figure 1) is a gauge theory with $U(n)$ gauge group, and $N = 4$ supersymmetry. Denoting by $\vec{v}$ the vector (x^3, x^4, x^5) , the Coulomb branch of this theory is parametrized by the v_i positions of the n D-3branes (with i labelling each brane). In addition, we have, as discussed in chapter II, the dual photons for each $U(1)$ factor. In this way, we get the hyperkähler structure of the Coulomb branch of the moduli. Hence, a direct way to get supersymmetry preserved by the brane configuration is as follows. The supersymmetry charges are defined as

$$\epsilon_L Q_L + \epsilon_R Q_R, \quad (4.31)$$

where Q_L and Q_R are the supercharges generated by the left and right-moving worldsheet degrees of freedom, and ϵ_L and ϵ_R are ten dimensional spinors. Each solitonic p brane,

with worldvolume extending along $x^0, x^1, \dots, x^p$, imposes the conditions

$$\epsilon_L = \Gamma_0 \dots \Gamma_p \epsilon_L, \quad \epsilon_R = -\Gamma_0 \dots \Gamma_p \epsilon_R, \quad (4.32)$$

in terms of the ten dimensional Dirac gamma matrices, Γ_i ; on the other hand, the D- p branes, with worldvolumes extending along $x^0, x^1, \dots, x^p$, imply the constraint

$$\epsilon_L = \Gamma_0 \Gamma_1 \dots \Gamma_p \epsilon_R. \quad (4.33)$$

Thus, we see that NS solitonic fivebrane, with worldvolume located at x^0, x^1, x^2, x^3, x^4 and x^5 , and equal values of $\vec{\omega}$, and Dirichlet threebranes with worldvolume along x^0, x^1, x^2 and x^6 , preserve eight supersymmetries on the D-3brane worldvolume or, equivalently, $N = 4$ supersymmetry on the effective three dimensional theory.

The brane array just described allows a simple computation of the gauge coupling constant of the effective three dimensional theory: by standard Kaluza-Klein reduction on the finite x^6 direction, after integrating over the (compactified) x^6 direction to reduce the lagrangian to an effective three dimensional lagrangian, the gauge coupling constant is given by

$$\frac{1}{g_3^2} = \frac{|x_6^2 - x_6^1|}{g_4^2}, \quad (4.34)$$

in terms of the four dimensional gauge coupling constant. Naturally, (4.34) is a classical expression that is not taking into account the effect on the fivebrane position at x^6 of the D-3brane ending on its worldvolume. In fact, we can consider the dependence of x^6 on the coordinate $\vec{v}$, normal to the position of the D-3brane. The dynamics of the fivebranes should then be recovered when the Nambu-Goto action of the solitonic fivebrane is minimized. Far from the influence of the points where the fivebranes are located (at large values of x^3, x^4 and x^5), the equation of motion is simply three dimensional Laplace's equation,

$$\nabla^2 x^6(x^3, x^4, x^5) = 0, \quad (4.35)$$

with solution

$$x^6(r) = \frac{k}{r} + \alpha, \quad (4.36)$$

where k and α are constants depending on the threebrane tensions, and r is the spherical radius at the point (x^3, x^4, x^5) . From (4.36), it is clear that there is a well defined limit as $r \rightarrow \infty$; hence, the difference $x_2^6 - x_1^6$ is a well defined constant, $\alpha_2 - \alpha_1$, in the $r \rightarrow \infty$ limit.

Part of the beauty of brane technology is that it allows to obtain very strong results by simply performing geometrical brane manipulations. We will now present one example, concerning our previous model. If we consider the brane configuration from the point of view of the fivebrane, the n suspended threebranes will look like n magnetic monopoles. This is really suggesting since, as described in chapter II, we know that the Coulomb branch moduli space of $N = 4$ supersymmetric $SU(n)$ gauge theories is isomorphic to the moduli space of BPS monopole configurations, with magnetic charge equal n . This analogy can be put more precisely: the vertex $(5^{NS}, 3)$ can, as described above, be transformed into a $(3, 1)$ vertex. In this case, from the point of view of the threebrane, we have a four dimensional gauge theory with $SU(2)$ gauge group broken down to $U(1)$, and n magnetic monopoles. Notice that by passing from the configuration build up using $(5^{NS}, 3)$ vertices, to that build up with the $(3, 1)$ vertex, the Coulomb moduli remains the same.

Next, we will work out the same configuration, but now with the vertex $(5, 3)$ made out of two Dirichlet branes. The main difference with the previous example comes from the boundary conditions (4.29) and (4.30), which should now be replaced by Dirichlet boundary conditions. We will choose as worldvolume coordinates for the D-5branes x^0, x^1, x^2, x^7, x^8 and x^9 , so that they will be located at some definite values of x^3, x^4, x^5 and x^6 . As before, let us denote this positions by $(\vec{m}, x^6)$, where now $\vec{m} = (x^3, x^4, x^5)$. An equivalent configuration to the one studied above will be now a set of two D-5branes, at some points of the x^6 coordinate, that we will again call x_1^6 and x_2^6 , subject to $\vec{m}_1 = \vec{m}_2$, with D-3branes stretching between them along the x^6 coordinate, with worldvolume extending again along the coordinates x^0, x^1, x^2 and x^6 (Figure 2). Our task now will

be the description of the effective three dimensional theory on these threebranes. The end points of the D-3branes on the fivebrane worldvolumes will now be parametrized by values of x^7, x^8 and x^9 . This means that we have three scalar fields in the effective three dimensional theory. The scalar fields corresponding to the coordinates x^3, x^4, x^5 and x^6 of the threebranes are frozen to the constant values where the fivebranes are located. Next, we should consider what happens to the $U(1)$ gauge field on the D-3brane worldvolume. Imposing Dirichlet boundary conditions for this field is equivalent to

$$F_{\mu\nu} = 0, \quad \mu, \nu = 0, 1, 2, \quad (4.37)$$

i. e., there is no electromagnetic tensor in the effective three dimensional field theory. Before going on, it would be convenient summarizing the rules we have used to impose the different boundary conditions. Consider a D- p brane, and let M be its worldvolume




Figure 2: Dirichlet threebranes extending between a pair of Dirichlet fivebranes (in dashed lines).

manifold, and $B = \partial M$ the boundary of M . Neumann and Dirichlet boundary conditions for the gauge field on the D - p brane worldvolume are defined respectively by

$$\begin{aligned} N &\longrightarrow F_{\mu\rho} = 0, \\ D &\longrightarrow F_{\mu\nu} = 0, \end{aligned} \tag{4.38}$$

where μ and ν are directions of tangency to B , and ρ are the normal coordinates to B . If B is part of the worldvolume of a solitonic brane, we will impose Neumann conditions, and if it is part of the worldvolume of a Dirichlet brane, we will impose Dirichlet conditions. Returning to (4.37), we see that on the three dimensional effective theory, the only non vanishing component of the four dimensional strength tensor is $F_{\mu 6} \equiv \partial_\mu b$. Therefore, all together we have four scalar fields in three dimensions or, equivalently, a multiplet with $N = 4$ supersymmetry. Thus, the theory defined by the n suspended D-3branes in between a pair of D-5branes, is a theory of n $N = 4$ massless hypermultiplets.

There exists a different way to interpret the theory, namely as a magnetic dual gauge theory. In fact, if we perform a duality transformation in the four dimensional $U(1)$ gauge theory, and use magnetic variables $*F$, instead of the electric field F , what we get in three dimensions, after imposing D-boundary conditions, is a dual photon, or a magnetic $U(1)$ gauge theory.

The configuration chosen for the worldvolume of the Dirichlet and solitonic fivebranes yet allows a different configuration with D-3branes suspended between a D-5brane and a NS-5brane. This is in fact consistent with the supersymmetry requirements (4.32) and

(4.33). Namely, for the Dirichlet fivebrane we have

$$\epsilon_L = \Gamma_0 \Gamma_1 \Gamma_2 \Gamma_7 \Gamma_8 \Gamma_9 \epsilon_R. \quad (4.39)$$

The solitonic fivebrane imposes

$$\epsilon_L = \Gamma_0 \dots \Gamma_5 \epsilon_L, \quad \epsilon_R = -\Gamma_0 \dots \Gamma_5 \epsilon_R, \quad (4.40)$$

while the suspended threebranes imply

$$\epsilon_L = \Gamma_0 \Gamma_1 \Gamma_2 \Gamma_6 \epsilon_R, \quad (4.41)$$

which are easily seen to be consistent. The problem now is that the suspended D-3brane is frozen. In fact, the position (x^3, x^4, x^5) of the end point of the NS-5brane is equal to the position $\vec{m}$ of the D-5brane, and the position (x^7, x^8, x^9) of the end point on the D-5brane is forced to be equal to the position $\vec{\omega}$ of the NS-5brane. The fact that the D-3brane is frozen means that the theory defined on it has no moduli, i. e., possesses a mass gap.

Using the vertices between branes we have described so far we can build quite complicated brane configurations. When Dirichlet threebranes are placed to the right and left of a fivebrane, open strings can connect the threebranes at different sides of the fivebrane. They will represent hypermultiplets transforming as $(k_1, \bar{k}_2)$, with k_1 and k_2 the number of threebranes to the left and right, respectively, of the fivebrane. In case the fivebrane is solitonic, the hypermultiplets are charged with respect to an electric group, while in case it is a D-5brane, they are magnetically charged. Another possibility is that with a pair of NS-5branes, with D-3branes extending between them, and also a D-5brane located between the two solitonic fivebranes. A massless hypermultiplet will now appear whenever the (x^3, x^4, x^5) position of the D-3brane coincides with the $\vec{m} = (x^3, x^4, x^5)$ position of the D-5brane.

So far we have used brane configurations for representing different gauge theories. In these brane configurations we have considered two different types of moduli. For the examples described above, these two types of moduli are as follows: the moduli of the effective three dimensional theory, corresponding to the different positions where the suspended D-3branes can be located, and the moduli corresponding to the different locations of the fivebranes, which are being used as boundaries. This second type of moduli specifies, from the point of view of the three dimensional theory, different coupling constants; hence, we can move the location of the fivebranes, and follow the changes taking place in the effective three dimensional theory. Let us then consider a case with

two solitonic branes, and a Dirichlet fivebrane placed between them. Let us now move the NS-5brane on the left of the D-5brane to the right. In doing so, there is a moment when both fivebranes meet, sharing a common value of x^6 . If the interpretation of the hypermultiplet we have presented above is correct, we must discover what happens to the hypermultiplet after this exchange of branes has been performed. In order to maintain the hypermultiplet, a new D-3brane should be created after the exchange, extending from the right solitonic fivebrane to the Dirichlet fivebrane. To prove this we will need D-brane dynamics at work. Let us start considering two interpenetrating closed loops, C and C' , and suppose electrically charged particles are moving in C , while magnetically charged particles move in C' . The linking number $L(C, C')$ can be defined using the standard Wilson and 't Hooft loops. Namely, we can measure the electric flux passing through C' or, equivalently, compute $B(C')$, or measure the magnetic flux passing through C , i. e., the Wilson line $A(C)$. In both cases, what we are doing is integrating over C' and C the dual to the field created by the particle moving in C and C' , respectively. Let us now extend this simple result to the case of fivebranes. A fivebrane is a source of 7-form tensor field, and its dual is therefore a 3-form. We will call this 3-form H_{NS} for NS-5branes, and H_D for D-5branes. Now, let us consider the worldvolume of the two fivebranes,

$$\begin{aligned}\mathbf{R}^3 &\times Y_{NS}, \\ \mathbf{R}^3 &\times Y_D.\end{aligned}\tag{4.42}$$

We can now define the linking number as we did before, in the simpler case of a particle:

$$L(Y_{NS}, Y_D) = - \int_{Y_{NS}} H^D = \int_{Y_D} H^{NS}.\tag{4.43}$$

The 3-form H^{NS} is locally dB_{NS} . Since we have no sources for H^{NS} , we can use $H^{NS} = dB_{NS}$ globally; however, this requires B to be globally defined, or gauge invariant. In type IIB string theory, B is not gauge invariant; however, on a D-brane we can define the combination $B_{NS} - F_D$, which is invariant, with F_D the two form for the $U(1)$ gauge field on the D-brane. Now, when the D-5brane and the NS-5brane do not intersect, the linking number is obviously zero. When they intersect, this linking number changes, which means that (4.43) should, in that case, be non vanishing. Writing

$$\int_{Y_D} H^{NS} = \int_{Y_D} dB_{NS} - dF_D,\tag{4.44}$$

we observe that the only way to get linking numbers would be adding sources for F_D . These sources for F_D are point like on Y_D , and are therefore the D-3branes with world-

volume $\mathbf{R}^3 \times C$, with C ending on Y_D , which is precisely the required appearance of extra D-3branes.

4.4 D-Brane Description of Seiberg-Witten Solution.

In the previous example we have considered type IIB string theory and three and fivebranes. Now, let us consider type IIA strings, where we have fourbranes that can be used to define, by analogy with the previous picture, $N = 2$ four dimensional gauge theories [103]. The idea will again be the use of solitonic fivebranes, with sets of fourbranes in between. The only difference now is that the fivebrane does not create a RR field in type IIA string theory and, therefore, the physics of the two parallel solitonic fivebranes does not have the interpretation of a gauge theory, as was the case for the type IIB configuration above described [103].

Let us consider configurations of infinite solitonic fivebranes, with worldvolume coordinates x^0, x^1, x^2, x^3, x^4 and x^5 , located at $x^7 = x^8 = x^9 = 0$ and at some fixed value of the x^6 coordinate. In addition, let us introduce finite Dirichlet fourbranes, with worldvolume coordinates x^0, x^1, x^2, x^3 and x^6 , which terminate on the solitonic fivebranes; thus, they are finite in the x^6 direction. On the fourbrane worldvolume, we can define a macroscopic four dimensional field theory, with $N = 2$ supersymmetry. This four dimensional theory will, as in the type IIB case considered in previous section, be defined by standard Kaluza-Klein dimensional reduction of the five dimensional theory defined on the D-4brane worldvolume. Then, the bare coupling constant of the four dimensional theory will be

$$\frac{1}{g_4^2} = \frac{|x_6^2 - x_6^1|}{g_5^2}, \quad (4.45)$$

in terms of the five dimensional coupling constant. Moreover, we can interpret as classical moduli parameters of the effective field theory on the dimensionally reduced worldvolume of the fourbrane the coordinates x^4 and x^5 , which locate the points on the fivebrane worldvolume where the D-4branes terminate.

In addition to the Dirichlet fourbranes and solitonic fivebranes, we can yet include Dirichlet sixbranes, without any further break of supersymmetry on the theory in the worldvolume of the fourbranes. To prove this, we notice that each NS-5brane imposes the projections

$$\epsilon_L = \Gamma_0 \dots \Gamma_5 \epsilon_L, \quad \epsilon_R = -\Gamma_0 \dots \Gamma_5 \epsilon_R, \quad (4.46)$$

while the D-4branes, with worldvolume localized at x^0, x^1, x^2, x^3 and x^6 , imply

$$\epsilon_L = \Gamma_0 \Gamma_1 \Gamma_2 \Gamma_3 \Gamma_6 \epsilon_R. \quad (4.47)$$

Conditions (4.46) and (4.47) can be recombined into

$$\epsilon_L = \Gamma_0 \Gamma_1 \Gamma_2 \Gamma_3 \Gamma_7 \Gamma_8 \Gamma_9 \epsilon_R, \quad (4.48)$$

which shows that certainly sixbranes can be added with no additional supersymmetry breaking.

The solitonic fivebranes break half of the supersymmetries, while the D-6brane breaks again half of the remaining symmetry, leaving eight real supercharges, which leads to four dimensional $N = 2$ supersymmetry.

As we will discuss later on, the sixbranes of type IIA string theory can be used to add hypermultiplets to the effective macroscopic four dimensional theory. In particular, the mass of these hypermultiplets will become zero whenever the D-4brane meets a D-6brane.

One of the main achievements of the brane representations of supersymmetric gauge theories is the ability to represent the different moduli spaces, namely the Coulomb and Higgs branches, in terms of the brane motions left free. For a configuration of k fourbranes connecting two solitonic fivebranes along the x^6 direction, as the one we have described above, the Coulomb branch of the moduli space of the four dimensional theory is parametrized by the different positions of the transversal fourbranes on the fivebranes. When N_f Dirichlet sixbranes are added to this configuration, what we are describing is the Coulomb branch of a four dimensional field theory with $SU(N_c)$ gauge group (in case N_c is the number of D-4branes we are considering), with N_f flavor hypermultiplets. In this brane representation, the Higgs branch of the theory is obtained when each fourbrane is broken into several pieces ending on different sixbranes: the locations of the D-4branes living between two D-6branes determine the Higgs branch. However, we will mostly concentrate on the study of the Coulomb branch for pure gauge theories.

As we know from the Seiberg-Witten solution of $N = 2$ supersymmetric gauge theories, the classical moduli of the theory is corrected by quantum effects. There are two types of effects that enter the game: a non vanishing beta function (determined at one loop) implies the existence, in the asymptotically free regime, of a singularity at the infinity point in moduli space, and strong coupling effects, which imply the existence of extra singularities, where some magnetically charged particles become massless. The problem we are facing now is how to derive such a complete characterization of the quantum moduli space of four dimensional $N = 2$ supersymmetric field theory directly from the dynamics governing

the brane configuration. The approach to be used is completely different from a brane construction in type IIA string theory to a type IIB brane configuration. In fact, in the type IIB case, employed in the description of the preceding section of three dimensional $N = 4$ supersymmetric field theories, we can pass from weak to strong coupling through the standard $Sl(2, \mathbf{Z})$ duality of type IIB strings; hence, the essential ingredient we need is to know how brane configurations transform under this duality symmetry. In the case of type IIA string theory, the situation is more complicated, as the theory is not $Sl(2, \mathbf{Z})$ self dual. However, we know that the strong coupling limit of type IIA dynamics is described by the eleven dimensional M-theory; therefore, we should expect to recover the strong coupling dynamics of four dimensional $N = 2$ supersymmetric gauge theories using the M-theory description of strongly coupled type IIA strings.

Let us first start by considering weak coupling effects. The first thing to be noticed, concerning the above described configuration of N_c Dirichlet fourbranes extending along the x^6 direction between two solitonic fivebranes, where only a rigid motion of the transversal fourbranes is allowed, is that this simple image is missing the classical dynamics of the fivebranes. In fact, in this picture we are assuming that the x^6 coordinate on the fivebrane worldvolume is constant, which is in fact a very bad approximation. Of course, one physical requirement we should impose to a brane configuration, as we did in the case of the type IIB configurations of the previous section, is that of minimizing the total worldvolume action. More precisely, what we have interpreted as Coulomb or Higgs branches in term of free motions of some branes entering the configuration, should correspond to zero modes of the brane configuration, i. e., to changes in the configuration preserving the condition of minimum worldvolume action (in other words, changes in the brane configuration that do not constitute an energy expense). The coordinate x^6 can be assumed to only depend on the “normal” coordinates x^4 and x^5 , which can be combined into the complex coordinate

$$v \equiv x^4 + ix^5, \quad (4.49)$$

representing the normal to the position of the transversal fourbranes. Far away from the position of the fourbranes, the equation for x^6 reduces now to a two dimensional laplacian,

$$\nabla^2 x^6(v) = 0, \quad (4.50)$$

with solution

$$x^6(v) = k \ln |v| + \alpha, \quad (4.51)$$

for some constants k and α , that will depend on the solitonic and Dirichlet brane tensions. As we can see from (4.51), the value of x^6 will diverge at infinity. This constitutes, as a

difference with the type IIB case, a first problem for the interpretation of equation (4.45). In fact, in deriving (4.45) we have used a standard Kaluza-Klein argument, where the four dimensional coupling constant is defined by the volume of the internal space (in this occasion, the x^6 interval between the two solitonic fivebranes). Since the Dirichlet four branes will deform the solitonic fivebrane, the natural way to define the internal space would be as the interval defined by the values of the coordinate x^6 at v equal to infinity, which is the region where the disturbing effect of the four brane is very likely vanishing, as was the case in the definition of the effective three dimensional coupling in the type IIB case. However, equations (4.50) and (4.51) already indicate us that this can not be the right picture, since these values of the x^6 coordinate are divergent. Let us then consider a configuration with N_c transversal fourbranes. From equations (4.45) and (4.51), we get, for large v ,

$$\frac{1}{g_4^2} = -\frac{2kN_c \ln(v)}{g_5^2}, \quad (4.52)$$

where we have differentiated the direction in which the fourbranes pull the fivebrane. Equation (4.52) can have a very nice meaning if we interpret it as the one loop renormalization group equation for the effective coupling constant. In order to justify this interpretation, let us first analyze the physical meaning of the parameter k . From equation (4.51), we notice that if we move in v around a value where a fourbrane is located (that we are assuming is $v = 0$), we get the monodromy transformation

$$x^6 \rightarrow x^6 + 2\pi i k. \quad (4.53)$$

This equation can be easily understood in M-theory, where we add an extra eleventh dimension, x^{10} , that we use to define the complex coordinate

$$x^6 + ix^{10}. \quad (4.54)$$

Now, using the fact that the extra coordinate is compactified on a circle of radius R we can, from (4.53), identify k with R . From a field theory point of view, we have a similar interpretation of the monodromy of (4.52), but now in terms of a change in the theta parameter. Let us then consider the one loop renormalization group equation for $SU(N_c)$ $N = 2$ supersymmetric gauge theories without hypermultiplets,

$$\frac{4\pi}{g_4^2(u)} = \frac{4\pi}{g_0^2} - \frac{2N_c}{4\pi} \ln\left(\frac{u}{\Lambda}\right), \quad (4.55)$$

with Λ the dynamically generated scale, and g_0 the bare coupling constant. The bare coupling constant can be absorbed through a change in Λ ; in fact, when going from Λ to

a new scale Λ' , we get

$$\frac{4\pi}{g_4^2(u)} = \frac{4\pi}{g_0^2} - \frac{2N_c}{4\pi} \ln\left(\frac{u}{\Lambda'}\right) - \frac{2N_c}{4\pi} \ln\left(\frac{\Lambda'}{\Lambda}\right). \quad (4.56)$$

Thus, once we fix a reference scale Λ_0 , the dependence on the scale Λ of the bare coupling constant is given by

$$- \frac{2N_c}{4\pi} \ln\left(\frac{\Lambda}{\Lambda_0}\right). \quad (4.57)$$

It is important to distinguish the dependence on Λ of the bare coupling constant, and the dependence on u of the effective coupling. In the brane configuration approach, the coupling constant defined by (4.52) is the bare coupling constant of the theory, as determined by the definite brane configuration. Hence, it is (4.57) that we should compare with (4.52); naturally, some care is needed concerning units and scales. Once we interpret k as the radius of the internal S^1 of M-theory we can, in order to make contact with (4.57), identify g_5^2 with the radius of S^1 , which in M-theory units is given by

$$R = gl_s, \quad (4.58)$$

with g the string coupling constant, and l_s the string length, $\frac{1}{\alpha'}$. Therefore, (4.45) should be modified to

$$\frac{1}{g_4^2} = \frac{x_2^6 - x_1^6}{gl_s} = -2N_c \ln(v), \quad (4.59)$$

which should be dimensionless. Then, we should interpret v in (4.59) as a dimensionless variable or, equivalently, as $\frac{v}{R}$, with R playing the role of natural unit of the theory. Then, comparing (4.57) and (4.59), $\frac{v}{R}$ becomes the scale $\frac{\Lambda}{\Lambda_0}$ in the formula for the bare coupling constant. In summary, v fixes the scale of the theory. From the previous discussion, an equivalent interpretation follows, where R fixes Λ_0 , and therefore changes in the scale are equivalent to changes in the radius of the internal S^1 .

Defining now an adimensional complex variable,

$$s \equiv (x^6 + ix^{10})/R, \quad (4.60)$$

and a complexified coupling constant,

$$\tau = \frac{4\pi i}{g^2} + \frac{\theta}{2\pi}, \quad (4.61)$$

we can generalize (4.59) to

$$-i\tau_\alpha(v) = s_2(v) - s_1(v), \quad (4.62)$$

for the simple configuration of branes defining a pure gauge theory. Now, we can clearly notice how the monodromy, as we move around $v = 0$, means a change $\theta \rightarrow +2\pi N_c$.

Let us now come back, for a moment, to the bad behaviour of $x^6(v)$ at large values of v . A possible way to solve this problem is modifying the configuration of a single pair of fivebranes, with N_c fourbranes extending between them, to consider a larger set of solitonic fivebranes. Labelling this fivebranes by α , with $\alpha = 0, \dots, n$, the corresponding x^6_α coordinate will depend on v as follows:

$$x^6(v)_\alpha = R \sum_{i=1}^{q_L} \ln |v - a_i| - R \sum_{j=1}^{q_R} \ln |v - b_j|, \quad (4.63)$$

where q_L and q_R represent, respectively, the number of D-4branes to the left and right of the α^{th} fivebrane. As is clear from (4.63), a good behaviour at large v will only be possible if the numbers of fourbranes to the right and left of a fivebrane are equal, $q_L = q_R$, which somehow amounts to compensating the perturbation created by the fourbranes at the sides of a fivebrane. The four dimensional field theory represented now by this brane array will have a gauge group $\prod_\alpha U(k_\alpha)$, where k_α is the number of transversal fourbranes between the $\alpha - 1$ and α^{th} solitonic fivebranes. Now, minimization of the worldvolume action will require not only taking into account the dependence of x^6 on v , but also the fourbrane positions on the NS-5brane, represented by a_i and b_j in (4.63), on the four dimensional worldvolume coordinates x^0, x^1, x^2 and x^3 . Using (4.63), and the Nambu-Goto action for the solitonic fivebrane, we get, for the kinetic energy,

$$\int d^4x d^2v \sum_{\mu=0}^3 \partial_\mu x^6(v, a_i(x^\mu), b_j(x^\mu)) \partial^\mu x^6(v, a_i(x^\mu), b_j(x^\mu)). \quad (4.64)$$

Convergence of the v integration implies

$$\partial_\mu \left(\sum_i a_i - \sum_j b_j \right) = 0 \quad (4.65)$$

or, equivalently,

$$\sum_i a_i - \sum_j b_j = \text{constant}. \quad (4.66)$$

This “constant of motion” is showing how the average of the relative position between left and right fourbranes must be hold constant. Since the Coulomb branch of the $\prod_\alpha U(k_\alpha)$ gauge theory will be associated with different configurations of the transversal fourbranes, constraint (4.66) will reduce the dimension of this space. As we know from

our general discussion on D-branes, the $U(1)$ part of the $U(k_\alpha)$ gauge group can be associated to the motion of the center of mass. Constraint (4.66) implies that the center of mass is frozen in each sector. With no semi-infinite fourbranes to the right, we have that $\sum_i a_i = 0$; now, this constraint will force the center of mass of all sectors to vanish, which means that the field theory we are describing is $\prod_\alpha SU(k_\alpha)$, instead of $\prod_\alpha U(k_\alpha)$. The same result can be derived if we include semi-infinite fourbranes to the left and right of the first and last solitonic fivebranes: as they are infinitely massive, we can assume that they do not move in the x^4 and x^5 directions. An important difference will appear if we consider periodic configurations of fivebranes, upon compactification of the x^6 direction to a circle: in this case, constraint (4.66) is now only able to reduce the group to $\prod_\alpha SU(k_\alpha) \times U(1)$, leaving alive a $U(1)$ factor.

Hypermultiplets in this gauge theory are understood as strings connecting fourbranes on different sides of a fivebrane; therefore, whenever the positions of the fourbranes to the left and right of a solitonic brane become coincident, a massless hypermultiplet arises. As the hypermultiplets are charged under the gauge groups at both sides of a certain $\alpha + 1$ fivebrane, they will transform as $(k_\alpha, \bar{k}_{\alpha+1})$.

However, as the position of the fourbranes on both sides of a fivebrane varies as a function of x^0, x^1, x^2 and x^3 , the existence of a well defined hypermultiplet can only be accomplished thanks to the fact that its variation rates on both sides are the same, as follows again from (4.65): $\partial_\mu(\sum_i a_{i,\alpha}) = \partial_\mu(\sum_j a_{j,\alpha+1})$. The definition of the bare masses comes then naturally from constraint (4.66):

$$m_\alpha = \frac{1}{k_\alpha} \sum_i a_{i,\alpha} - \frac{1}{k_{\alpha+1}} \sum_j a_{j,\alpha+1}. \quad (4.67)$$

With this interpretation, the constraint (4.66) becomes very natural from a physical point of view: it states that the masses of the hypermultiplets do not depend on the spacetime position.

The consistency of the previous definition of hypermultiplets can be checked using the previous construction of the one-loop beta function. In fact, from equation (4.62), we get, for large values of v ,

$$-i\tau_\alpha(v) = (2k_\alpha - k_{\alpha-1} - k_{\alpha+1}) \ln v. \quad (4.68)$$

The number k_α of branes in the α^{th} is, as we know, the number of colours, N_c . Comparing with the beta function for $N = 2$ supersymmetric $SU(N_c)$ gauge theory with N_f flavors, we conclude that

$$N_f = k_{\alpha-1} + k_{\alpha+1}, \quad (4.69)$$

so that the number of fourbranes (hypemultiplets) at both sides of a certain pair of fivebranes, $k_{\alpha+1} + k_{\alpha-1} \equiv N_f$, becomes the number of flavors.

Notice, from (4.67), that the mass of all the hypermultiplets associated with fourbranes at both sides of a solitonic fivebrane are the same. This implies a global flavor symmetry. This global flavor symmetry is the gauge symmetry of the adjacent sector. This explains the physical meaning of (4.67).

Let us now come back to equation (4.59). What we need in order to unravel the strong coupling dynamics of our effective four dimensional gauge theory is the u dependence of the effective coupling constant, dependence that will contain non perturbative effects due to instantons. It is from this dependence that we read the Seiberg-Witten geometry of the quantum moduli space. Strong coupling effects correspond to u in the infrared region, i. e., small u or, equivalently, large Λ . From our previous discussion of (4.59), we conclude that the weak coupling regime corresponds to the type IIA string limit, $R \rightarrow 0$, and the strong coupling regime to the M-theory regime, at large values of R (recall that changes of scale in the four dimensional theory correspond to changes of the radius of the internal S^1). This explains our hopes that M-theory could describe the strong coupling regime of the four dimensional theory). We will then see now how M-theory is effectively working.

4.4.1 M-Theory and Strong Coupling.

From the M-theory point of view, the brane configuration we are considering can be interpreted in a different way. In particular, the D-4branes we are using to define the four dimensional macroscopic gauge theory can be considered as fivebranes wrapping the eleven dimensional S^1 . Moreover, the trick we have used to make finite these fourbranes in the x^6 direction can be directly obtained if we consider fivebranes with worldvolume $\mathbf{R}^4 \times \Sigma$, where $\mathbf{R}^4$ is parametrized by the coordinates x^0, x^1, x^2 and x^4 , and Σ is two dimensional, and embedded in the four dimensional space of coordinates x^4, x^5, x^6 and x^{10} . If we think in purely classical terms, the natural guess for Σ would be a cylinder with the topology $S^1 \times [x_2^6, x_1^6]$, for a configuration of k D-4branes extending along the x^6 direction between two solitonic fivebranes. This is however a very naive compactification, because there is no reason to believe that a fivebrane wrapped around this surface will produce, on the four dimensional worldvolume $\mathbf{R}^4$, any form of non abelian gauge group. In fact, any gauge field on $\mathbf{R}^4$ should come from integrating the chiral antisymmetric tensor field β of the M-theory fivebrane worldvolume, on some one-cycle of Σ . If we want to reproduce, in four dimensions, some kind of $U(k)$ or $SU(k)$ gauge theory, we should better consider a surface Σ with a richer first homology group. However, we can try to do something

better when including the explicit dependence of the x^6 coordinate on v . In this case, we will get a picture that is closer to the right answer, but still far away from the true solution. Including the v dependence of the x^6 coordinate leads to a family of surfaces, parametrized by v , Σ_v , defined by $S^1 \times [x_2^6, x_1^6](v)$. The nice feature about this picture is that v , which is the transverse coordinate of Σ in the space $\mathcal{Q}$, defined by the coordinates x^4, x^5, x^6 and x^{10} , becomes now similar to the moduli of Σ_v ; however, we have yet the problem of the of the genus or, in more general terms, the first homology group of Σ . The reason for following the previous line of thought, is that we are trying to keep alive the interpretation of the v coordinate as moduli, or coordinate of the Coulomb branch. This is, in fact, the reason giving rise to the difficulties with the genus, as we are using just one complex coordinate, independently of the rank of the gauge group, something we are forced to do because of the divergences in equation (4.51).

The right M-theory approach is quite different. In fact, we must try to get Σ directly from the particular brane configuration we are working with, and define the Coulomb branch of the theory by the moduli space of brane configurations. Let us then define the single valued coordinate t ,

$$t \equiv \exp -s, \quad (4.70)$$

and define the surface Σ we are looking for through

$$F(t, v) = 0. \quad (4.71)$$

From the classical equations of motion of the fivebrane we know the asymptotic behaviour for very large t ,

$$t \sim v^k, \quad (4.72)$$

and for very small t ,

$$t \sim v^{-k}. \quad (4.73)$$

Conditions (4.72) and (4.73) imply that $F(t, v)$ will have, for fixed values of t , k roots, while two different roots for fixed v . It must be stressed that the asymptotic behaviour (4.72) and (4.73) corresponds to the one loop beta function for a field theory with gauge group $SU(k)$, and without hypermultiplets. A function satisfying the previous conditions will be of the generic type

$$F(t, v) = A(v)t^2 + B(v)t + C(v), \quad (4.74)$$

with A , B and C polynomials in v of degree k . From (4.72) and (4.73), the function (4.74) becomes

$$F(t, v) = t^2 + B(v)t + \text{constant}, \quad (4.75)$$

with one undetermined constant. In order to kill this constant, we can rescale t to $t/\text{constant}$. The meaning of this rescaling can be easily understood in terms of the one loop beta function, written as (4.72) and (4.73). In fact, these equations can be read as

$$s = -k \ln \left(\frac{v}{R} \right), \quad (4.76)$$

and therefore the rescaling of R goes like

$$s \rightarrow -k \ln \left(\frac{v}{R'} \frac{R'}{R} \right) \quad (4.77)$$

or, equivalently,

$$t \rightarrow t \left(\frac{R'}{R} \right)^k. \quad (4.78)$$

Thus, and based on the above discussion on the definition of the scale, we observe that the constant in (4.75) defines the scale of the theory. With this interpretation of the constant, we can get the Seiberg-Witten solution for $N = 2$ pure gauge theories, with gauge group $SU(k)$. If $B(v)$ is chosen to be

$$B(v) = v^k + u_2 v^{k-2} + u_3 v^{k-3} + \cdots + u_k, \quad (4.79)$$

we finally get the Riemann surface

$$t^2 + B(v)t + 1 = 0, \quad (4.80)$$

a Riemann surface of genus $k - 1$, which is in fact the rank of the gauge group. Moreover, we can now try to visualize this Riemann surface as the worldvolume of the fivebrane describing our original brane configuration: each v -plane can be compactified to $\mathbf{P}^1$, and the transversal fourbranes can be interpreted as gluing tubes, which clearly represents a surface with $k - 1$ handles. This image corresponds to gluing two copies of $\mathbf{P}^1$, with k disjoint cuts on each copy or, equivalently, $2k$ branch points. Thus, as can be observed from (4.80), to each transversal D-4brane there correspond two branch points and one cut on $\mathbf{P}^1$.

If we are interested in $SU(k)$ gauge theories with hypermultiplets, then we should first replace (4.72) and (4.73) by the corresponding relations,

$$t \sim v^{k-k_{\alpha}-1}, \quad (4.81)$$

and

$$t \sim v^{-k-k_{\alpha}+1}, \quad (4.82)$$

for t large and small, respectively. These are, in fact, the relations we get from the beta functions for these theories. If we take $k_{\alpha_1} = 0$, and $N_f = k_{\alpha+1}$, the curve becomes

$$t^2 + B(v)t + C(v) = 0, \quad (4.83)$$

with $C(v)$ a polynomial in v , of degree N_f , parametrized by the masses of the hypermultiplets,

$$C(v) = f \prod_{j=1}^{N_f} (v - m_j), \quad (4.84)$$

with f a complex constant.

Summarizing, we have been able to find a moduli of brane configurations reproducing four dimensional $N = 2$ supersymmetric $SU(k)$ gauge theories. The exact Seiberg-Witten solution is obtained by reduction of the worldvolume fivebrane dynamics on the surface $\Sigma_{\vec{u}}$ defined at (4.80) and (4.82). Obviously, reducing the fivebrane dynamics to $\mathbf{R}^4$ on $\Sigma_{\vec{u}}$ leads to an effective coupling constant in $\mathbf{R}^4$, the $k - 1 \times k - 1$ Riemann matrix $\tau(\vec{u})$ of $\Sigma_{\vec{u}}$.

Before finishing this section, it is important to stress some peculiarities of the brane construction. First of all, it should be noticed that the definition of the curve Σ , in terms of the brane configuration, requires working with uncompactified x^4 and x^5 directions. This is part of the brane philosophy, where we must start with a particular configuration in flat spacetime. A different approach will consist in directly working with a spacetime $\mathcal{Q} \times \mathbf{R}^7$, with $\mathcal{Q}$ some Calabi-Yau manifold, and consider a fivebrane worldvolume $\Sigma \times \mathbf{R}^4$, with $\mathbf{R}^4 \subset \mathbf{R}^7$, and Σ a lagrangian submanifold of $\mathcal{Q}$. Again, by Mc Lean's theorem, the $N = 2$ theory defined on $\mathbf{R}^4$ will have a Coulomb branch with dimension equal to the first Betti number of Σ , and these deformations of Σ in $\mathcal{Q}$ will represent scalar fields in the four dimensional theory. Moreover, the holomorphic top form Ω of $\mathcal{Q}$ will define the meromorphic λ of the Seiberg-Witten solution. If we start with some Calabi-Yau manifold $\mathcal{Q}$, we should provide some data to determine Σ (this is what we did in the brane case, with $\mathcal{Q}$ non compact and flat. If, on the contrary, we want to select Σ directly from $\mathcal{Q}$, we can only do it in some definite cases, which are those related to the *geometric mirror construction* [118, 119]. Let us then recall some facts about the geometric mirror. The data are

- The Calabi-Yau manifold $\mathcal{Q}$.
- A lagrangian submanifold $\Sigma \rightarrow \mathcal{Q}$.

- A $U(1)$ flat bundle on Σ .

The third requirement is equivalent to interpreting Σ as a D-brane in $\mathcal{Q}$. This is a crucial data, in order to get from the above points the structure of abelian manifold of the Seiberg-Witten solution. Namely, we first use Mc Lean's theorem to get the moduli of deformations of $\Sigma \rightarrow \mathcal{Q}$, preserving the condition of lagrangian submanifold. This space is of dimension $b_1(\Sigma)$. Secondly, on each of these points we fiber the jacobian of Σ , which is of dimension g . This family of abelian varieties defines the quantum moduli of a gauge theory, with $N = 2$ supersymmetry, with a gauge group of rank equal $b_1(\Sigma)$. Moreover, this family of abelian varieties is the moduli of the set of data of the second and third points above, i. e., the moduli of Σ as a D-2brane. In some particular cases, this moduli is $\mathcal{Q}$ itself or, more properly, the geometric mirror of $\mathcal{Q}$. This will be the case for Σ of genus equal one, i. e., for the simple $SU(2)$ case. In this cases, the characterization of Σ in $\mathcal{Q}$ is equivalent to describing $\mathcal{Q}$ as an elliptic fibration. The relation between geometric mirror and T-duality produces a completely different physical picture. In fact, we can, when Σ is a torus, consider in type IIB a threebrane with classical moduli given by $\mathcal{Q}$. After T-duality or mirror, we get the type IIA description in terms of a fivebrane. In summary, it is an important problem to understand the relation of quantum mirror between type IIA and type IIB string theory, and the M-theory strong coupling description of type IIA strings.

4.5 Brane Description of $N = 1$ Four Dimensional Field Theories.

In order to consider field theories with $N = 1$ supersymmetry, the first thing we will study will be R -symmetry. Let us then recall the way R -symmetries were defined in the case of four dimensional $N = 2$ supersymmetry, and three dimensional $N = 4$ supersymmetry, through compactification of six dimensional $N = 1$ supersymmetric gauge field theories. The $U(1)_R$ in four dimensions, or $SO(3)_R$ in three dimensions, are simply the euclidean group of rotations in two and three dimensions, respectively. Now, we have a four dimensional space $\mathcal{Q}$, parametrized by coordinates t and v , and a Riemann surface Σ , embedded in $\mathcal{Q}$ by equations of the type (4.74). To characterize R -symmetries, we can consider transformations on $\mathcal{Q}$ which transform non trivially its holomorphic top form Ω . The unbroken R -symmetries will then be rotations in $\mathcal{Q}$ preserving the Riemann surface defined by the brane configuration. If we consider only the asymptotic behaviour of type

(4.72), or (4.81), we get $U(1)_R$ symmetries of type

$$\begin{aligned} t &\rightarrow \lambda^k t, \\ v &\rightarrow \lambda v. \end{aligned} \tag{4.85}$$

This $U(1)$ symmetry is clearly broken by the curve (4.80). This spontaneous breakdown of the $U(1)_R$ symmetry is well understood in field theory as an instanton induced effect. If instead of considering $\mathcal{Q}$, we take the larger space $\hat{\mathcal{Q}}$, containing the x^7, x^8 and x^9 coordinates, we see that the $N = 2$ curve is invariant under rotations in the (x^7, x^8, x^9) space.

Let us now consider a brane configuration which reproduces $N = 1$ four dimensional theories [117]. We will again start in type IIA string theory, and locate a solitonic fivebrane at $x^6 = x^7 = x^8 = x^9 = 0$ with, as usual, worldvolume coordinates x^0, x^1, x^2, x^3, x^4 and x^5 . At some definite value of x^6 , say x_0^6 , we locate another solitonic fivebrane, but this time with worldvolume coordinates x^0, x^1, x^2, x^3, x^7 and x^8 , and $x^4 = x^5 = x^9 = 0$. As before, we now suspend a set of k D-4branes in between. They will be parametrized by the positions $v = x^4 + ix^5$, and $w = x^7 + ix^8$, on the two solitonic fivebranes. The worldvolume coordinates on this D-4branes are, as in previous cases, x^0, x^1, x^2 and x^3 . The effective field theory defined by the set of fourbranes is macroscopically a four dimensional gauge theory, with coupling constant

$$\frac{1}{g^2} = \frac{x_0^6}{gl_s}. \tag{4.86}$$

Moreover, now we have only $N = 1$ supersymmetry, as no massless bosons can be defined on the four dimensional worldvolume (x^0, x^1, x^2, x^3) . In fact, at the line $x^6 = 0$ the only possible massless scalar would be v , since $w = 0$ and $x^9 = 0$, so that we project out x^9 and w . On the other hand, at x_0^6 we have $v = 0$ and $x^9 = 0$ and, therefore, we have projected out all massless scalars. Notice that by the same argument, in the case of two solitonic fivebranes located at different values of x^6 but at $x^7 = x^8 = x^9 = 0$, we have one complex massless scalar that is not projected out, which leads to $N = 2$ supersymmetry in four dimensions. The previous discussion means that v , w and x^9 are projected out as four dimensional scalar fields; however, w and v are still classical moduli parameters of the brane configuration.

Now, we return to a comment already done in previous section: each of the fourbranes we are suspending in between the solitonic fivebranes can be interpreted as a fivebrane wrapped around a surface defined by the eleven dimensional S^1 of M-theory, multiplied by the segment $[0, x_0^6]$. Classically, the four dimensional theory can be defined through

dimensional reduction of the fivebrane worldvolume on the surface Σ . The coupling constant will be given by the moduli τ of this surface,

$$\frac{1}{g^2} = \frac{2\pi R}{S}, \quad (4.87)$$

with S the length of the interval $[0, x_0^6]$, in M-theory units. In $N = 1$ supersymmetric field theories, on the contrary of what takes place in the $N = 2$ case, we have not a classical moduli and, therefore, we can not define a wilsonian coupling constant depending on some mass scale fixed by a vacuum expectation value. This fact can produce some problems, once we take into account the classical dependence of x^6 on v and w . In principle, this dependence should be the same as that in the case studied in previous section,

$$\begin{aligned} x^6 &\sim \tilde{k} \ln v, \\ x^6 &\sim \tilde{k} \ln w. \end{aligned} \quad (4.88)$$

Using the t coordinate defined in (4.70), equations (4.88) become

$$\begin{aligned} t &\sim v^k, \\ t &\sim w^k, \end{aligned} \quad (4.89)$$

for large and small t , respectively, or, equivalently, $t \sim v^k$, $t^{-1} \sim w^k$. Now, we can use these relations in (4.86). Taking into account the units, we can write

$$\frac{1}{g^2} \sim N_c \left[\ln \frac{v}{R} + \ln \frac{w}{R} \right], \quad (4.90)$$

with $k \equiv N_c$. As we did in the $N = 2$ case, we can try to compare (4.90) with the one loop beta function for $N = 1$ supersymmetric $SU(N_c)$ pure Yang-Mills theory,

$$\Lambda = \mu \exp - \frac{8\pi^2}{3N_c g(\mu)^2}. \quad (4.91)$$

In order to get the scale from (4.90) we impose

$$v = \zeta w^{-1}, \quad (4.92)$$

with ζ some constant with units of $(\text{length})^2$. Using (4.92) and (4.90) we get

$$\frac{1}{g^2} \sim N_c \ln \left(\frac{\zeta}{R^2} \right). \quad (4.93)$$

In order to make contact with (4.91) we must impose

$$\frac{\zeta}{R^2} = (\Lambda R)^3, \quad (4.94)$$

where we have used $\frac{1}{R}$ in order to measure Λ . Using (4.92), we get the curve associated to four dimensional $N = 1$ field theory,

$$\begin{aligned} t &= v^k, \\ \zeta^k t^{-1} &= w^k \\ v &= \zeta w^{-1}. \end{aligned} \quad (4.95)$$

The curve defined by (4.95) will only depend on ζ^k . The different set of brane configurations compatible with (4.95) are given by values of ζ , with fixed ζ^k . These N_c roots parametrize the N_c different vacua predicted by $\text{tr}(-1)^F$ arguments. It is important to observe that the coupling constant $\frac{1}{g^2}$ we are defining is the so called wilsonian coupling. We can interpret it as a complex number with $\text{Im } \frac{1}{g^2} \equiv \frac{\theta}{8\pi^2}$. Hence, the value of $\text{Im } \zeta^k$ fixes the θ parameter of the four dimensional theory.

For a given value of ζ , (4.95) defines a Riemann surface of genus zero, i. e., a rational curve. This curve is now embedded in the space of (t, v, w) coordinates. We will next observe that these curves, (4.95), are the result of “rotating” [106] the rational curves in the Seiberg-Witten solution, corresponding to the singular points. However, before doing that let us comment on $U(1)_R$ symmetries. As mentioned above, in order to define an R -symmetry we need a transformation on variables (t, v, w) not preserving the holomorphic top form,

$$\Omega = dv \wedge dw \wedge \frac{dt}{t} R. \quad (4.96)$$

A rotation in the w -plane, compatible with the asymptotic conditions (4.89), and defining an R -symmetry, is

$$\begin{aligned} v &\rightarrow v, \\ t &\rightarrow t, \\ w &\rightarrow e^{2\pi i/k} w. \end{aligned} \quad (4.97)$$

Now, it is clear that this symmetry is broken spontaneously by the curve (4.95). More interesting is an exact $U(1)$ symmetry, that can be defined for the curve (4.95):

$$\begin{aligned} v &\rightarrow e^{i\delta} v, \\ t &\rightarrow e^{i\delta k} t, \\ w &\rightarrow e^{-i\delta} w. \end{aligned} \quad (4.98)$$

As can be seen from (4.96), this is not an R -symmetry, since Ω is invariant. Fields charged with respect to this $U(1)$ symmetry should carry angular momentum in the v or w plane, or linear momentum in the eleventh dimension interval (i. e., zero branes). The fields of $N = 1$ SQCD do not carry any of these charges, so all fields with $U(1)$ charge should be decoupled from the $N = 1$ SQCD degrees of freedom. This is equivalent to the way we have projected out fields in the previous discussion on the definition of the effective $N = 1$ four dimensional field theory.

4.5.1 Rotation of Branes.

A different way to present the above construction is by performing a rotation of branes. We will now concentrate on this procedure. The classical configuration of NS-5branes with worldvolumes extending along x^0, x^1, x^2, x^3, x^4 and x^5 , can be modified to a configuration where one of the solitonic fivebranes has been rotated, from the $v = x^4 + ix^5$ direction, to be also contained in the (x^7, x^8) -plane, so that, by moving it a finite angle μ , it is localized in the (x^4, x^5, x^7, x^8) space. Using the same notation as in previous section, the brane configuration, where a fivebrane has been moved to give rise to an angle μ in the (v, w) -plane, the rotation is equivalent to imposing

$$w = \mu v. \tag{4.99}$$

In the brane configuration we obtain, points on the rotated fivebrane are parametrized by the (v, w) coordinates in the (x^4, x^5, x^7, x^8) space. We can therefore impose the following asymptotic conditions [116]:

$$\begin{aligned} t &= v^k, & w &= \mu v, \\ t &= v^{-k}, & w &= 0, \end{aligned} \tag{4.100}$$

respectively for large and small t . Let us now assume that this brane configuration describes a Riemann surface, $\hat{\Sigma}$, embedded in the space $(x^6, x^{10}, x^4, x^5, x^7, x^8)$, and let us denote by Σ the surface in the $N = 2$ case, i. e., for $\mu = 0$. In these conditions, $\hat{\Sigma}$ is simply the graph of the function w on Σ . We can interpret (4.99) as telling us that w on Σ possesses a simple pole at infinity, extending holomorphically over the rest of the Riemann surface. If we impose this condition, we get that the projected surface Σ , i. e., the one describing the $N = 2$ theory, is of genus zero. In fact, it is a well known result in the theory of Riemann surfaces that the order of the pole at infinity depends on the genus of the surface in such a way that for genus larger than zero, we will be forced to

replace (4.99) by $w = \mu v^a$ for some power a depending on the genus. A priori, there is no problem in trying to rotate using, instead of $w = \mu v$, some higher pole modification of the type $w = \mu v^a$, for $a > 1$. This would provide Σ surfaces with genus different from zero; however, we would immediately find problems with equation (4.90), and we will be unable to kill all dependence of the coupling constant on v and w . Therefore, we conclude that the only curves that can be rotated to produce a four dimensional $N = 1$ theory are those with zero genus. This is in perfect agreement with the physical picture we get from the Seiberg-Witten solution. Namely, once we add a soft breaking term of the type $\mu \text{tr} \phi^2$, the only points remaining in the moduli space as real vacua of the theory are the singular points, where the Seiberg-Witten curve degenerates.

4.5.2 QCD Strings and Scales.

In all our previous discussion we have not been careful enough in separating arguments related to complex or *holomorphic structure*, and those related to *Kähler structure*. The M-theory description contains however relevant information on both aspects. For instance, in our previous derivation of curves, we were mostly interested in reproducing the complex structure of the Seiberg-Witten solution, as is, for instance, the moduli dependence on vacuum expectation values, i. e., the effective wilsonian coupling constant. However, we can also ask ourselves on BPS masses and, in that case, we will need the definite embedding of Σ in the ambient space $\mathcal{Q}$, and the holomorphic top form defined on $\mathcal{Q}$. As is clear from the fact that we are working in M-theory, the holomorphic top form on $\mathcal{Q}$ will depend explicitly on R , i. e., on the string coupling constant, and we will therefore find BPS mass formulas that will depend explicitly on R . We will discuss this type of dependence on R first in the case of $N = 1$ supersymmetry. The $N = 1$ four dimensional field theory we have described contains, in principle, two parameters. One is the constant ζ introduced in equation (4.92) which, as we have already mentioned, is, because of (4.90), intimately connected with Λ , and the radius R of the eleven dimensional S^1 . Our first task would be to see what kind of four dimensional dynamics is dependent on the particular value of R , and in what way. The best example we can of course use is the computation of gaugino-gaugino condensates. In order to do that, we should try to minimize a four dimensional superpotential for the $N = 1$ theory. Following Witten, we will define this superpotential W as an holomorphic function of Σ , and with critical points precisely when the surface Σ is a holomorphic curve in $\mathcal{Q}$. The space $\mathcal{Q}$ now is the one with coordinates x^4, x^5, x^6, x^7, x^8 and x^{10} (notice that this second condition was the one used to prove that rotated curves are necessarily of genus equal zero). Moreover, we

need to work with a holomorphic curve because of $N = 1$ supersymmetry. A priori, there are two different ways we can think about this superpotential: maybe the simplest one, from a physical point of view, is as a functional defined on the volume of Σ , where this volume is given by

$$\text{Vol}(\Sigma) = J.\Sigma, \quad (4.101)$$

with J the Kähler class of $\mathcal{Q}$. The other possibility is defining

$$W(\Sigma) = \int_B \Omega, \quad (4.102)$$

with B a 3-surface such that $\Sigma = \partial B$, and Ω the holomorphic top form in $\mathcal{Q}$. Definition (4.102) automatically satisfies the condition of being stationary, when Σ is a holomorphic curve in $\mathcal{Q}$. Notice that the holomorphy condition on Σ means, in mathematical terms, that Σ is an element of the Picard lattice of $\mathcal{Q}$, i. e., an element in $H_{1,1}(\mathcal{Q}) \cap H_2(\mathcal{Q})$. This is what allows us to use (4.101), however, and this is the reason for temporarily abandoning the approach based on (4.101). What we require to W is being stationary for holomorphic curves, but it should, in principle, be defined for arbitrary surfaces Σ , even those which are not part of the Picard group. Equation (4.102) is only well defined if Σ is contractible, i. e., if the homology class of Σ in $H_2(\mathcal{Q}; \mathbf{Z})$ is trivial. If that is not the case, a reference surface Σ_0 needs to be defined, and (4.102) is modified to

$$W(\Sigma) - W(\Sigma_0) = \int_B \Omega, \quad (4.103)$$

where now $\partial B = \Sigma \cup \Sigma_0$. For simplicity, we will assume $H_3(\mathcal{Q}; \mathbf{Z}) = 0$. From physical arguments we know that the set of zeroes of the superpotential should be related by $\mathbf{Z}_k$ symmetry, with k the number of transversal fourbranes. Therefore, if we choose Σ_0 to be $\mathbf{Z}_k$ invariant, we can write $W(\Sigma_0) = 0$, and $W(\Sigma) = \int_B \Omega$. Let us then take B as the complex plane multiplied by an interval $I = [0, 1]$, and let us first map the complex plane into Σ . Denoting r the coordinate on this complex plane, Σ , as given by (4.95), is defined by

$$\begin{aligned} t &= r^k, \\ v &= r, \\ w &= \zeta r^{-1}. \end{aligned} \quad (4.104)$$

Writing $r = e^\rho e^{i\theta}$, we can define Σ_0 as

$$\begin{aligned} t &= r^k, \\ v &= f(\rho)r, \\ w &= \zeta f(-\rho)r^{-1}, \end{aligned} \quad (4.105)$$

with $f(\rho) = 1$ for $\rho > 2$, and $f(\rho) = 0$ for $\rho < 1$. The $\mathbf{Z}_k$ transformation $t \rightarrow t$, $w \rightarrow e^{2\pi i/k}w$ and $v \rightarrow v$, is a symmetry of (4.105) if, at the same time, we perform the reparametrization of the r -plane

$$\begin{aligned}\rho &\rightarrow \rho, \\ \theta &\rightarrow \theta + b(\rho),\end{aligned}\tag{4.106}$$

with $b(\rho) = 0$ for $\rho \geq 1$, and $b(\rho) = -\frac{2\pi}{k}$ for $\rho \leq -1$. Thus, the 3-manifold entering the definition of B , is given by

$$\begin{aligned}t &= r^k, \\ v &= g(\rho, \sigma)r, \\ w &= \zeta g(-\rho, \sigma)r^{-1},\end{aligned}\tag{4.107}$$

such that for $\sigma = 0$ we have $g = 1$, and for $\sigma = 1$, we get $g(\rho) = f(\rho)$. Now, with

$$\Omega = R dv \wedge dw \wedge \frac{dt}{t},\tag{4.108}$$

we get

$$W(\Sigma) = kR \int_B dv \wedge dw \wedge \frac{dr}{r}.\tag{4.109}$$

The dependence on R is already clear from (4.109). In order to get the dependence on ζ we need to use (4.107),

$$W(\Sigma) = kR\zeta \int d\sigma d\theta d\rho \left(\frac{\partial g_+}{\partial \sigma} \frac{\partial g_-}{\partial \rho} - \frac{\partial g_+}{\partial \rho} \frac{\partial g_-}{\partial \sigma} \right),\tag{4.110}$$

for $g_{\pm} = g(\pm\rho, \sigma)$. Thus we get

$$W(\Sigma) \sim kR\zeta,\tag{4.111}$$

Notice that the superpotential (4.111) is given in units $(\text{length})^3$, as corresponds to the volume of a 3-manifold. In order to make contact with the gaugino-gaugino condensate, we need to obtain $(\text{length})^{-3}$ units. We can do this multiplying by $\frac{1}{R^6}$; thus, we get

$$< \lambda \lambda > \sim kR\zeta \frac{1}{R^6} \sim \Lambda^3,\tag{4.112}$$

where we have used equation (4.108). A different way to connect ζ with Λ is defining, in the M-theory context, the QCD string and computing its tension. Following Witten, we will then try an interpretation of ζ independent of (4.90), by computing in terms of ζ the

tension of the QCD string. We will then, to define the tension, consider the QCD string as a membrane, product of a string in $\mathbf{R}^4$, and a string living in $\mathcal{Q}$. Let us then denote by C a curve in $\mathcal{Q}$, and assume that C ends on Σ in such a way that a membrane wrapped on C defines a string in $\mathbf{R}^4$ ¹⁸. Moreover, we can simply think of C as a closed curve in $\mathcal{Q}$, going around the eleven dimensional S^1 ,

$$\begin{aligned} t &= t_0 \exp(-2\pi i \sigma), \\ v &= t_0^{1/k}, \\ w &= \zeta v^{-1}. \end{aligned} \tag{4.113}$$

This curve is a non trivial element in $H_1(\mathcal{Q}; \mathbf{Z})$, and a membrane wrapped on it will produce an ordinary type IIA string; however, we can not think that the QCD string is a type IIA string. If $\mathcal{Q} = \mathbf{R}^3 \times S^1$, then $H_1(\mathcal{Q}; \mathbf{Z}) = \mathbf{Z}$, and curves of type (4.113) will be the only candidates for non trivial 1-cycles in $\mathcal{Q}$. However, we can define QCD strings using cycles in the relative homology, $H_1(\mathcal{Q}/\Sigma; \mathbf{Z})$, i. e., considering non trivial cycles ending on the surface Σ . To compute $H_1(\mathcal{Q}/\Sigma; \mathbf{Z})$, we can use the exact sequence

$$H_1(\Sigma; \mathbf{Z}) \rightarrow H_1(\mathcal{Q}; \mathbf{Z}) \xrightarrow{\iota} H_1(\mathcal{Q}/\Sigma; \mathbf{Z}), \tag{4.114}$$

which implies

$$H_1(\mathcal{Q}/\Sigma; \mathbf{Z}) = H_1(\mathcal{Q}; \mathbf{Z}) / \iota H_1(\Sigma; \mathbf{Z}). \tag{4.115}$$

The map ι is determined by the map defining Σ ($t = v^k$), and thus we can conclude that, very likely,

$$H_1(\mathcal{Q}/\Sigma; \mathbf{Z}) = \mathbf{Z}_k. \tag{4.116}$$

A curve in $H_1(\mathcal{Q}/\Sigma; \mathbf{Z})$ can be defined as follows:

$$\begin{aligned} t &= t_0, \\ v &= t_0^{1/k} e^{2\pi i \sigma/k}, \\ w &= \zeta v^{-1}, \end{aligned} \tag{4.117}$$

with $t_0^{1/k}$ one of the k roots. The tension of (4.117), by construction, is independent of R , because t is fixed. Using the metric on $\mathcal{Q}$, the length of (4.117) is given by

$$\left(\frac{\zeta^2 t^{-2/n}}{n^2} + \frac{t^{2/n}}{n^2} \right)^{1/2}, \tag{4.118}$$

¹⁸Notice that if we were working in type IIB string theory, we would have the option to wrap a threebrane around Σ , in order to define a string on $\mathbf{R}^4$.

and its minimum is obtained when $t^{2/n} = \zeta$. Thus, the length of the QCD string should be

$$\frac{|\zeta|^{1/2}}{n}, \quad (4.119)$$

which has the right length units, as ζ behaves as $(\text{length})^2$. In order to define the tension we need to go to $(\text{length})^{-1}$ units, again using $\frac{1}{R^2}$. Then, if we identify this tension with Λ , we get

$$\Lambda \sim \frac{|\zeta|^{1/2}}{n} \frac{1}{R^2} \quad (4.120)$$

or, equivalently,

$$\Lambda^2 \sim \Lambda^3 R. \quad (4.121)$$

Thus, consistency with QCD results requires $\Lambda \sim \frac{1}{R}$. These are not good news, as they imply that the theory we are working with, in order to match QCD, possesses 0-brane modes, with masses of the order of Λ , and therefore we have not decoupled the M-theory modes.

Next, we would like to compare the superpotential described above with the ones obtained using standard instanton techniques in M-theory. However, before doing that we will conclude this brief review on brane configurations with the description of models with $N = 4$ supersymmetry.

4.5.3 $N = 2$ Models with Vanishing Beta Function.

Let us come back to brane configurations with $n+1$ solitonic fivebranes, with k_α Dirichlet fourbranes extending between the α^{th} pair of NS-5branes. The beta function, derived in (4.68), is

$$-2k_\alpha + k_{\alpha+1} + k_{\alpha-1}, \quad (4.122)$$

for each $SU(k_\alpha)$ factor in the gauge group. In this section, we will compactify the x^6 direction to a circle of radius L . Imposing the beta function to vanish in all sectors immediately implies that all k_α are the same. Now, the compactification of the x^6 direction does not allow to eliminate all $U(1)$ factors in the gauge group: one of them can not be removed, so that the gauge group is reduced from $\prod_{\alpha=1}^n U(k_\alpha)$ to $U(1) \times SU(k)^n$. Moreover, using the definition (4.67) of the mass of the hypermultiplets we get, for periodic configurations,

$$\sum_{\alpha} m_{\alpha} = 0. \quad (4.123)$$

The hypermultiplets are now in representations of type $k \otimes \bar{k}$, and therefore consists of a copy of the adjoint representation, and a neutral singlet.

Let us consider the simplest case, of $N = 2$ $SU(2) \times U(1)$ four dimensional theory, with one hypermultiplet in the adjoint representation [103]. The corresponding brane configuration contains a single solitonic fivebrane, and two Dirichlet fourbranes. The mass of the hypermultiplet is clearly zero, and the corresponding four dimensional theory has vanishing beta function. A geometric procedure to define masses for the hypermultiplets is a fibering of the v -plane on the x^6 S^1 direction, in a non trivial way, so that the fourbrane positions are identified modulo a shift in v ,

$$\begin{aligned} x^6 &\rightarrow x^6 + 2\pi L, \\ v &\rightarrow v + m, \end{aligned} \tag{4.124}$$

so that now, the mass of the hypermultiplet, is the constant m appearing in (4.124), as $\sum_{\alpha} m_{\alpha} = m$.

From the point of view of M-theory, the x^{10} coordinate has also been compactified on a circle, now of radius R . The (x^6, x^{10}) space has the topology of $S^1 \times S^1$. This space can be made non trivial if, when going around x^6 , the value of x^{10} is changed as follows:

$$\begin{aligned} x^6 &\rightarrow x^6 + 2\pi L, \\ x^{10} &\rightarrow x^{10} + \theta R, \end{aligned} \tag{4.125}$$

and, in addition, $x^{10} \rightarrow x^{10} + 2\pi R$. Relations (4.125) define a Riemann surface of genus one, and moduli depending on L and θ for fixed values of R . θ in (4.125) can be understood as the θ -angle of the four dimensional field theory: the θ -angle can be defined as

$$\frac{x_1^{10} - x_2^{10}}{R}, \tag{4.126}$$

with $x_2^{10} = x^{10}(2\pi L)$, and $x_1^{10} = x^{10}(0)$. Using (4.125), we get θ as the value of (4.126). This is the bare θ -angle of the four dimensional theory.

A question immediately appears concerning the value of the bare coupling constant: the right answer should be

$$\frac{1}{g^2} = \frac{2\pi L}{R}. \tag{4.127}$$

It is therefore clear that we can move the bare coupling constant of the theory keeping fixed the value of R , and changing L and θ . Let us now try to solve this model for the massless case. The solution will be given by a Riemann surface Σ , living in the space

$E \times C$, where E is the Riemann surface defined by (4.125), and C is the v -plane. Thus, all what we need is defining Σ through an equation of the type

$$F(x, y, z) = 0, \quad (4.128)$$

with x and y restricted by the equation of E ,

$$y^2 = (x - e_1(\tau))(x - e_2(\tau))(x - e_3(\tau)), \quad (4.129)$$

with τ the bare coupling constant defined by (4.126) and (4.127) [120]. In case we have a collection of k fourbranes, we will require F to be a polynomial of degree k in v ,

$$F(x, y, z) = v^k - f_1(x, y)v^{k-1} + \dots \quad (4.130)$$

The moduli parameters of Σ are, at this point, hidden in the functions $f_i(x, y)$ in (4.130). Let us denote $v_i(x, y)$ the roots of (4.130) at the point (x, y) in E . Notice that (4.130) is a spectral curve defining a branched covering of E , i. e., (4.130) can be interpreted as a spectral curve in the sense of Hitchin's integrable system [121]. If f_i has a pole at some point (x, y) , then the same root $v_i(x, y)$ should go to infinity. These poles have the interpretation of locating the position of the solitonic fivebranes. In the simple case we are considering, with a single fivebrane, the Coulomb branch of the theory will be parametrized by meromorphic functions on E with a simple pole at one point, which is the position of the fivebrane. As we have k functions entering (4.130), the dimension of the Coulomb branch will be k , which is the right one for a theory with $U(1) \times SU(k)$ gauge group.

Now, after this discussion of the model with massless hypermultiplets, we will introduce the mass. The space where now we need to define Σ is not $E \times C$, but the non trivial fibration defined through

$$\begin{aligned} x^6 &\rightarrow x^6 + 2\pi L, \\ x^{10} &\rightarrow x^{10} + \theta R, \\ v &\rightarrow v + m \end{aligned} \quad (4.131)$$

or, equivalently, the space obtained by fibering C non trivially on E . We can flat this bundle over all E , with the exception of one point p_0 . Away from this point, the solution is given by (4.130). If we write (4.130) in a factorized form,

$$F(x, y, z) = \prod_{i=1}^k (v - v_i(x, y)), \quad (4.132)$$

we can write f_1 in (4.130) as the sum

$$f_1 = \sum_{i=1}^k v_i(x, y); \quad (4.133)$$

therefore, f_1 will have poles at the positions of the fivebrane. The mass of the hypermultiplet will be identified with the residue of the differential $f_1\omega$, with ω the abelian differential, $\omega = \frac{dx}{y}$. As the sum of the residues is zero, this means that at the point at infinity, that we identify with p_0 , we have a pole with residue m .

4.6 M-Theory and String Theory.

In this section we will compare the M-theory description of $N = 2$ and $N = 1$ four dimensional gauge theories, with that obtained in string theory upon performing the point particle limit [122, 123, 124, 125]. Let us then return for a moment to the brane representation of $N = 2$ four dimensional gauge theories. In the M-theory approach, we will consider M-theory on flat spacetime, $\mathbf{R}^7 \times \mathcal{Q}$, with $\mathcal{Q} = \mathbf{R}^5 \times S^1$. The S^1 stands for the (compactified) eleventh dimension, with the radius R proportional to the string coupling constant. The brane configuration in $\mathbf{R}^7 \times \mathcal{Q}$ turns out to be equivalent to a solitonic fivebrane, with worldvolume $\Sigma \times \mathbf{R}^4$, where Σ is a complex curve in $\mathcal{Q}$, defined by

$$F(t, v) = 0. \quad (4.134)$$

This is equivalent to defining an embedding

$$\Phi : \Sigma \hookrightarrow \mathcal{Q}. \quad (4.135)$$

If Σ is a lagrangian manifold of $\mathcal{Q}$, then we can interpret the moduli space of the effective four dimensional $N = 2$ theory as the space of deformations of Φ in (4.135) preserving the condition of lagrangian submanifold¹⁹. By Mc Lean's theorem, we know that the dimension of this space of deformations is $b_1(\Sigma)$, in agreement with the existing relation between the genus of Σ and the rank of the gauge group in the effective four dimensional theory. It is important keeping in mind that in the M-theory approach two ingredients

¹⁹Recall that a lagrangian manifold is defined by the condition that

$$\int_{\Sigma} \Phi^*(\Omega) = \text{Vol}(\Sigma),$$

with Φ^* such that $\Phi^*(\omega) = 0$ (where ω is the Kähler class of $\mathcal{Q}$), and Ω the holomorphic top form of $\mathcal{Q}$.

are being used: the curve defined by (4.134), and the holomorphic top form Ω of $\mathcal{Q}$, which explicitly depends on the radius R of the eleventh dimension. This will be very important, as already noticed in the discussion of the $N = 1$ superpotentials, because an explicit dependence on the string coupling constant will be induced in the BPS mass formulas.

A different approach to (4.134) and (4.135) is that based on geometric engineering [127]. In this case, the procedure is based on the following set of steps:

1. String theory is compactified on a Calabi-Yau threefold X , with the appropriate number of vector multiplets in four dimensions.
2. A point corresponding to classical enhancement of gauge symmetry in the moduli space of the Calabi-Yau threefold must be localized.
3. A rigid Calabi-Yau threefold is defined by performing a point particle limit.
4. The rigid Calabi-Yau manifold is used to define the Seiberg-Witten surface Σ .
5. Going from type IIB to type IIA string theory represents a brane configuration corresponding to an ALE space with singularity of some Dynkin type into a set of fivebranes that can be interpreted as a fivebrane with worldvolume $\Sigma \times \mathbf{R}^4$.
6. The BPS states are defined through the meromorphic one-form λ , derived from the the Calabi-Yau holomorphic top form, in the rigid point particle limit.

As we can see from the previous set of steps, that we will explicitly show at work in one definite example, the main difference between both approaches is at the level of the meromorphic form in Seiberg-Witten theory. There is also an important difference in the underlying philosophy, related to the implicit use in the string approach, described in the above steps, of the heterotic-type II dual pairs, driving us to the choice of a particular Calabi-Yau manifold. The most elaborated geometric engineering approach uses, instead of a certain heterotic-type II dual pair, a set of local geometrical data, determined by the type of gauge symmetry we are interested on, and generalizes mirror maps to this set of local data. In all these cases, the four dimensional field theory we are going to obtain will not depend on extra parameters, as the string coupling constant. On the other hand, the M-theory approach, where field theories are obtained depending explicitly on the string coupling constant, might be dynamically rich enough as to provide a direct explanation of

phenomena that can not be easily understood in the more restricted context of the point particle limit of string theory.

Next, we will follow steps 1 to 6 through an explicit example [124]. In order to obtain a field theory with gauge group $SU(n)$ we should start with a Calabi-Yau manifold with $h_{2,1} = n$, and admitting the structure of a $K3$ -fibered threefold (see chapter II for definitions, and additional details). We will consider the $SU(3)$ case, corresponding to a Calabi-Yau manifold whose mirror is the weighted projective space $\mathbb{P}_{1,1,2,8,12}^{24}$,

$$\frac{1}{24}(x_1^{24} + x_2^{24}) + \frac{1}{12}x_3^{12} + \frac{1}{2}x_5^2 - \psi_0 x_1 x_2 x_3 x_4 x_5 - \frac{1}{6}(x_1 x_2 x_3)^6 - \frac{1}{12}(x_1 x_2)^{12} = 0. \quad (4.136)$$

In order to clearly visualize (4.136) as a $K3$ -fibration we will perform the change of variables

$$x_1/x_2 \equiv \hat{z}^{1/12} b^{-1/24}, \quad x_1^2 \equiv x_0 \hat{z}^{1/12}, \quad (4.137)$$

so that (4.136) can be rewritten in the form

$$\frac{1}{24}(\hat{z} + \frac{b}{\hat{z}} + 2)x_0^{12} + \frac{1}{12}x_3^{12} + \frac{1}{3}x_4^3 + \frac{1}{2}x_5^2 + \frac{1}{6\sqrt{c}}(x_0 x_3)^6 + \left(\frac{a}{\sqrt{c}}\right)^{1/6} x_0 x_3 x_4 x_5 = 0, \quad (4.138)$$

which represents a $K3$ surface, fibered over a $\mathbb{P}^1$ space parametrized by the coordinate z . Parameters in (4.138) are related to those in (4.136) through

$$a = -\psi_0^6/\psi_1, \quad b = \psi_2^{-2}, \quad c = \psi_2/\psi_1^2. \quad (4.139)$$

The parameter b can be interpreted as the volume of $\mathbb{P}^1$:

$$-\log b = \text{Vol}(\mathbb{P}^1). \quad (4.140)$$

Next, we should look for the points $\hat{z}$ in $\mathbb{P}^1$ over which the $K3$ surface is singular. The discriminant can be written as

$$\Delta_{K3} = \prod_{i=0}^2 (\hat{z} - e_i^+(a, b, c))(\hat{z} - e_i^-(a, b, c)), \quad (4.141)$$

where

$$\begin{aligned} e_0^\pm &= -1 \pm \sqrt{1-b}, \\ e_1^\pm &= \frac{1-c \pm \sqrt{(1-c)^2 - bc^2}}{c}, \\ e_2^\pm &= \frac{(1-a)^2 - c \pm \sqrt{((1-a)^2 - c)^2 - bc^2}}{c}. \end{aligned} \quad (4.142)$$

The Calabi-Yau manifold will be singular whenever two roots e_i coalesce, as

$$\Delta_{\text{Calabi-Yau}} = \prod_{i < j} (e_i - e_j)^2. \quad (4.143)$$

We will consider the singular point in the moduli space corresponding to $SU(3)$ symmetry. Around this point we will introduce new coordinates, through

$$\begin{aligned} a &= -2(\alpha' u)^{3/2}, \\ b &= \alpha' \Lambda^6, \\ c &= 1 - \alpha'^{3/2}(-2u^{3/2} + 3\sqrt{3}v). \end{aligned} \quad (4.144)$$

Going now to the $\alpha' \rightarrow 0$ limit in (4.143), we get a set of roots $e_i(u, v; \Lambda^6)$ on a z -plane, with z defined in $\alpha'^{3/2}z \equiv \hat{z}$:

$$\begin{aligned} e_0 &= 0, \quad e_\infty = \infty, \\ e_1^\pm &= 2u^{3/2} + 3\sqrt{3}v \pm \sqrt{(2u^{3/2} + 3\sqrt{3}v)^2 - \Lambda^6}, \\ e_2^\pm &= -2u^{3/2} + 3\sqrt{3}v \pm \sqrt{(2u^{3/2} - 3\sqrt{3}v)^2 - \Lambda^6}. \end{aligned} \quad (4.145)$$

Now, we can use (4.145) as the definition of a Riemann surface Σ , defined by the Calabi-Yau data at the singular $SU(3)$ point, and in the point particle limit. There exists a natural geometrical picture for understanding the parameters u and v in (4.144), which is the definition of the blow up, in the moduli space of complex structures of $\mathbb{P}_{1,1,2,8,12}^{24}$, of the $SU(3)$ singular point. From this point of view, the parameters u and v in (4.144) will be related to the volume of the set of vanishing two-cycles associated with a *rational* singularity, i. e., an orbifold singularity of type A_{n-1} (in the case we are considering, $n = 3$). These vanishing cycles, as is the case with rational singularities, are associated with Dynkin diagrams of non affine type. The branch points (4.145) on the z -plane define the curve

$$y^2 = \prod_i (x - e_i(u, v; \Lambda^6)), \quad (4.146)$$

which can also be represented as the vanishing locus of a polynomial $F(x, z) = 0$, with F given by [129, 130]

$$F(x, z) = z + \frac{\Lambda^6}{z} + B(x), \quad (4.147)$$

where $B(x)$ is a polynomial in x of degree three; in the general case of $SU(n)$ theories, the polynomial will be of degree n .

This has exactly the same look as what we have obtained using brane configurations, with the space $\mathcal{Q}$ replaced by the (x, z) space. The difference is that now we are not considering the (x, z) space as a part of spacetime, and Σ as embedded in it, but we use Σ as defined in (4.147) to define a Calabi-Yau space in a rigid limit by the equation

$$F(x, z) + y^2 + w^2 = 0, \quad (4.148)$$

which defines a threefold in the (x, y, z, w) space. And, in addition, we think of (4.148) as a Calabi-Yau representation of the point particle limit. In order to get the meromorphic one-form λ , and the BPS states, we need to define a map from the third homology group, $H_3(CY)$, of the Calabi-Yau manifold, into $H_1(\Sigma)$. This can be done as follows. The three-cycles in $H_3(CY)$ of the general type $S^2 \times S^1$, with S^2 a vanishing cycle of $K3$, correspond to S^1 circles in the z -plane. The three-cycles with the topology of S^3 can be interpreted as a path from the north to the south pole of S^3 , starting with a vanishing two-cycle, and ending at another vanishing two-cycle of $K3$. This corresponds, in the z -plane, to paths going from e_i^+ to e_i^- . Once we have defined this map,

$$f : H_3(CY) \longrightarrow H_1(\Sigma), \quad (4.149)$$

we define

$$\lambda(f(C)) = \Omega(C), \quad (4.150)$$

with Ω the holomorphic top form.

A similar analysis can be done for computing the mass of BPS states, and the meromorphic one-form λ in the brane framework. In fact, we can consider a two-cycle C in $\mathcal{Q}$ such that

$$\partial C \subset \Sigma, \quad (4.151)$$

or, in other words, $C \in H_2(\mathcal{Q}/\Sigma; \mathbf{Z})$. The holomorphic top form on $\mathcal{Q}$ is given by

$$\Omega = R \frac{dt}{t} \wedge dv, \quad (4.152)$$

and thus the BPS mass will be given by

$$M \sim R \int_C \frac{dt}{t} \wedge dv = R \int_{\partial C} \frac{dt}{t} v(t), \quad (4.153)$$

with $v(t)$ given by

$$F(t, v) = 0, \quad (4.154)$$

for the corresponding Seiberg-Witten curve, Σ . Notice that the same analysis, using (4.150) and the holomorphic top form for (4.148) will give, by contrast to the brane case, a BPS mass formula independent of R .

Next, we will compare the brane construction and geometric engineering in the more complicated case of $N = 1$ [126, 128].

4.7 Local Models for Elliptic Fibrations.

Let V be an elliptic fibration,

$$\Phi : V \rightarrow \Delta, \quad (4.155)$$

with Δ an algebraic curve, and $\Phi^{-1}(a)$, with a any point in Δ , an elliptic curve. Let us denote $\{a_\rho\}$ the finite set of points in Δ such that $\Phi^{-1}(a_\rho) = \mathcal{C}_\rho$ is a singular fiber. Each singular fiber $\mathcal{C}_\rho$ can be written as

$$\mathcal{C}_\rho = \sum_i n_{i\rho} \Theta_{i\rho}, \quad (4.156)$$

where $\Theta_{i\rho}$ are non singular rational curves, with $\Theta_{i\rho}^2 = -2$, and $n_{i\rho}$ are integer numbers. Different types of singularities are characterized by (4.156) and the intersection matrix $(\Theta_{i\rho} \cdot \Theta_{j\rho})$. All different types of Kodaira singularities satisfy the relation

$$\mathcal{C}_\rho^2 = 0. \quad (4.157)$$

Let $\tau(u)$ be the elliptic modulus of the elliptic fiber at the point $u \in \Delta$. For each path α in $\Pi_1(\Delta')$, with $\Delta' = \Delta - \{a_\rho\}$, we can define a monodromy transformation S_α , in $Sl(2, \mathbf{Z})$, acting on $\tau(u)$ as follows:

$$S_\alpha \tau(u) = \frac{a_\alpha \tau(u) + b_\alpha}{c_\alpha \tau(u) + d_\alpha}. \quad (4.158)$$

Each type of Kodaira singularity is characterized by a particular monodromy matrix.

In order to define an elliptic fibration [45], the starting point will be an algebraic curve Δ , that we will take, for simplicity, to be of genus zero, and a meromorphic function $\mathcal{J}(u)$ on Δ . Let us assume $\mathcal{J}(u) \neq 0, 1, \infty$ on $\Delta' = \Delta - \{a_\rho\}$. Then, there exists multivalued holomorphic function $\tau(u)$, with $\text{Im } \tau(u) > 0$, satisfying $\mathcal{J}(u) = j(\tau(u))$, with j the elliptic modular j -function on the upper half plane. As above, for each $\alpha \in \Pi_1(\Delta')$ we define a monodromy matrix S_α , acting on $\tau(u)$ in the form defined by (4.158). Associated to these data we will define an elliptic fibration, (4.155). In order to do that, let us first define the universal covering $\tilde{\Delta}'$, of Δ' , and let us identify the covering transformations of

$\tilde{\Delta}'$ over Δ' , with the elements in $\Pi_1(\Delta')$. Denoting by $\tilde{u}$ a point in $\tilde{\Delta}'$, we define, for each $\alpha \in \Pi_1(\Delta')$, the covering transformation $\tilde{u} \rightarrow \alpha\tilde{u}$, by

$$\tau(\alpha\tilde{u}) = S_\alpha\tau(\tilde{u}); \quad (4.159)$$

in other words, we consider τ as a single valued holomorphic function on $\tilde{\Delta}'$. Using (4.158), we define

$$f_\alpha(\tilde{u}) = (c_\alpha\tau(\tilde{u}) + d_\alpha)^{-1}. \quad (4.160)$$

Next, we define the product $\tilde{\Delta}' \times \mathbf{C}$ and, for each (α, n_1, n_2) , with $\alpha \in \Pi_1(\Delta')$, and n_1, n_2 integers, the automorphism

$$g(\alpha, n_1, n_2) : (\tilde{u}, \lambda) \rightarrow (\alpha\tilde{u}, f_\alpha(\tilde{u})(\lambda + n_1\tau(\tilde{u}) + n_2)). \quad (4.161)$$

Denoting by $\mathcal{G}$ the group of automorphisms (4.161), we define the quotient space

$$B' \equiv (\tilde{\Delta}' \times \mathbf{C})/\mathcal{G}. \quad (4.162)$$

This is a non singular surface, since g , as defined by (4.161), has no fixed points in $\tilde{\Delta}'$. From (4.161) and (4.162), it is clear that B' is an elliptic fibration on Δ' , with fiber elliptic curves of elliptic modulus $\tau(u)$. Thus, by the previous construction, we have defined the elliptic fibration away from the singular points a_ρ .

Let us denote E_ρ a local neighbourhood of the point a_ρ , with local coordinate t , and such that $t(a_\rho) = 0$. Let S_ρ be the monodromy associated with a small circle around a_ρ . By $\mathcal{U}_\rho$ we will denote the universal covering of $E'_\rho = E_\rho - a_\rho$, with coordinate ρ defined by

$$\rho = \frac{1}{2\pi i} \log t. \quad (4.163)$$

The analog of (4.159) will be

$$\tau(\rho + 1) = S_\rho\tau(\rho). \quad (4.164)$$

If we go around the points a_ρ , k times, we should act with S_ρ^k ; hence, we parametrize each path by the winding number k . The group of automorphisms (4.161), reduced to small closed paths around a_ρ , becomes

$$g(k, n_1, n_2)(\rho, \lambda) = (\rho + k, f_k(\rho)[\lambda + n_1\tau(\rho) + n_2]). \quad (4.165)$$

Denoting by $\mathcal{G}_\rho$ the group (4.165), we define the elliptic fibration around a_ρ as

$$(\mathcal{U}_\rho \times \mathbf{C})/\mathcal{G}_\rho. \quad (4.166)$$

Next, we will extend the elliptic fibration to the singular point a_ρ . We can consider two different cases, depending on the finite or infinite order of S_ρ .

4.8 Singularities of Type $\hat{D}_4$: $\mathbf{Z}_2$ Orbifolds.

Let us assume S_ρ is of finite order,

$$(S_\rho)^m = \mathbf{1}_d. \quad (4.167)$$

In this case, we can extend (4.166) to the singular points, simply defining a new variable σ as

$$\sigma^m = t. \quad (4.168)$$

Let us denote D a local neighbourhood in the σ -plane of the point $\sigma = 0$, and define the group G_D of automorphisms

$$g(n_1, n_2) : (\sigma, \lambda) = (\sigma, \lambda + n_1\tau(\sigma) + n_2), \quad (4.169)$$

and the space

$$F = (D \times \mathbf{C})/G_D. \quad (4.170)$$

Obviously, F defines an elliptic fibration over D , with fiber F_σ at each point $\sigma \in D$, an elliptic curve of modulus $\tau(\sigma)$. From (4.167) and (4.160), it follows that

$$f_k(\sigma) = 1, \quad (4.171)$$

with $k = O(m)$. Thus, we can define a normal subgroup $\mathcal{N}$ of $\mathcal{G}_\rho$ as the set of transformations (4.165):

$$g(k, n_1, n_2) : (\rho, \lambda) \rightarrow (\rho + k, \lambda + n_1\tau(\rho) + n_2). \quad (4.172)$$

Comparing now (4.169) and (4.172), we get

$$(\mathcal{U}_\rho \times \mathbf{C})/\mathcal{N} = (D' \times \mathbf{C})/G_D \equiv F - F_0. \quad (4.173)$$

Using (4.172) and (4.165) we get

$$\mathcal{C} = \mathcal{G}/\mathcal{N}, \quad (4.174)$$

with $\mathcal{C}$ the cyclic group of order m , defined by

$$g_k : (\sigma, \lambda) \rightarrow (e^{2\pi i k/m} \sigma, f_k(\sigma) \lambda). \quad (4.175)$$

From (4.174) and (4.173), we get the desired extension to a_ρ , namely

$$F/\mathcal{C} = (\mathcal{U}_\rho \times \mathbf{C})/\mathcal{G}_\rho \cup F_0/\mathcal{C}. \quad (4.176)$$

Thus, the elliptic fibration extended to a_ρ , in case S_ρ is of finite order, is defined by $F/\mathcal{C}$. Now, $F/\mathcal{C}$ can have singular points that we can regularize. The simplest example corresponds to

$$S_\rho = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (4.177)$$

i. e., a parity transformation. In this case, the order is $m = 2$, and we define σ by $\sigma^2 \equiv t$. The cyclic group (4.175) in this case simply becomes

$$(\sigma, \lambda) \rightarrow (-\sigma, -\lambda), \quad (4.178)$$

since from (4.177) and (4.160) we get $f_1 = -1$. At the point $\sigma = 0$ we have four fixed points, the standard $\mathbf{Z}_2$ orbifold points,

$$(0, \frac{a}{2}\tau(0) + \frac{b}{2}), \quad (4.179)$$

with $a, b = 0, 1$. The resolution of these four singular points will produce four irreducible components, $\Theta^1, \dots, \Theta^4$. In addition, we have the irreducible component Θ_0 , defined by the curve itself at $\sigma = 0$. Using the relation $\sigma^2 = t$, we get the $\hat{D}_4$ cycle,

$$\mathcal{C} = 2\Theta_0 + \Theta^1 + \Theta^2 + \Theta^3 + \Theta^4, \quad (4.180)$$

with $(\Theta_0, \Theta^1) = (\Theta_0, \Theta^2) = (\Theta_0, \Theta^3) = (\Theta_0, \Theta^4) = 1$. In general, the four external points of D -diagrams can be associated with the four $\mathbf{Z}_2$ orbifold points of the torus.

4.9 Singularities of Type $\hat{A}_{n-1}$.

We will now consider the case

$$S_\rho = \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}, \quad (4.181)$$

which is of infinite order. A local model for this monodromy can be defined by

$$\tau(t) = \frac{1}{2\pi i} n \log t. \quad (4.182)$$

Using the variable ρ defined in (4.163), we get, for the group $\mathcal{G}_\rho$ of automorphisms,

$$g(k, n_1, n_2) : (\rho, \lambda) \rightarrow (\rho + k, \lambda + n_1 n \rho + n_2), \quad (4.183)$$

and the local model for the elliptic fibration, out of the singular point,

$$(\mathcal{U}_\rho \times \mathbf{C})/\mathcal{G}_\rho, \quad (4.184)$$

i. e., fibers of the type of elliptic curves, with elliptic modulus $n\rho$. A simple way to think about these elliptic curves is in terms of cyclic unramified coverings [63]. Let us recall that a cyclic unramified covering, $\Pi : \hat{C} \rightarrow C$, of order n , of a curve C of genus g , is a curve $\hat{C}$ of genus

$$\hat{g} = ng + 1 - n. \quad (4.185)$$

Thus, for $g = 1$, we get $\hat{g} = 1$, for arbitrary n . Denoting by τ the elliptic modulus of C , in case $g = 1$, the elliptic modulus of $\hat{C}$ is given by

$$\hat{\tau} = n\tau. \quad (4.186)$$

Moreover, the generators $\hat{\alpha}$ and $\hat{\beta}$ of $H_1(\hat{C}; \mathbf{Z})$ are given in terms of the homology basis α, β of C as

$$\begin{aligned} \Pi\hat{\alpha} &= \alpha, \\ \Pi\hat{\beta} &= n\beta, \end{aligned} \quad (4.187)$$

with Π the projection $\Pi : \hat{C} \rightarrow C$. From (4.186) and (4.183), we can interpret the elliptic fibration (4.184) as one with elliptic fibers given by n -cyclic unramified coverings of a curve C with elliptic modulus ρ or, equivalently, $\frac{1}{2\pi i} \log t$. There exists a simple way to define a family of elliptic curves, with elliptic modulus given by $\frac{1}{2\pi i} \log t$, which is the plumbing fixture construction. Let D_0 be the unit disc around $t = 0$, and let C_0 be the Riemann sphere. Define two local coordinates, $z_a : \mathcal{U}_a \rightarrow D_0$, $z_b : \mathcal{U}_b \rightarrow D_0$, in disjoint neighbourhoods $\mathcal{U}_a, \mathcal{U}_b$, of two points P_a and P_b of C_0 . Let us then define

$$\begin{aligned} W = \{ (p, t) | t \in D_0, p \in C_0 - \mathcal{U}_a - \mathcal{U}_b, \text{ or } p \in \mathcal{U}_a, \text{ with } |z_a(p)| > |t|, \text{ or} \\ p \in \mathcal{U}_b, \text{ with } |z_b(p)| > |t| \}, \end{aligned} \quad (4.188)$$

and let S be the surface

$$S = \{ xy = t; (x, y, t) \in D_0 \times O_0 \times D_0 \}. \quad (4.189)$$

We define the family of curves through the following identifications

$$\begin{aligned} (p_a, t) \in W \cap \mathcal{U}_a \times D_0 &\simeq (z_a(p_a), \frac{t}{z_a(p_a)}, t) \in S, \\ (p_b, t) \in W \cap \mathcal{U}_b \times D_0 &\simeq (\frac{t}{z_b(p_b)}, z_b(p_b), t) \in S. \end{aligned} \quad (4.190)$$

For each t we get a genus one curve, and at $t = 0$ we get a nodal curve by pinching the non zero homology cycles. The pinching region is characterized by

$$xy = t, \quad (4.191)$$

which defines a singularity of type A_0 . The elliptic modulus of the curves is given by

$$\tau(t) = \frac{1}{2\pi i} \log t + C_1 t + C_2, \quad (4.192)$$

for some constants C_1 and C_2 . We can use an appropriate choice of coordinate t , such that $C_1 = C_2 = 0$. The singularity at $t = 0$ is a singularity of type $\hat{A}_0$, in Kodaira's classification, corresponding to

$$S_\rho = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}. \quad (4.193)$$

Using now (4.186) and (4.192) we get, for the cyclic covering of order n , the result (4.182), and the group (4.183). The pinching region of the cyclic unramified covering is given by

$$xy = t^n, \quad (4.194)$$

instead of (4.191), i. e., for the surface defining the A_{n-1} singularity, $\mathbf{C}^2/\mathbf{Z}_n$. Now, we can proceed to the resolution of the singularity at $t = 0$. The resolution of the singularity (4.194) requires $n-1$ exceptional divisors, $\Theta_1, \dots, \Theta_{n-1}$. In addition, we have the rational curve Θ_0 , defined by the complement of the node. Thus, we get, at $t = 0$,

$$\mathcal{C} = \Theta_0 + \dots + \Theta_{n-1}, \quad (4.195)$$

with $(\Theta_0, \Theta_1) = (\Theta_0, \Theta_{n-1}) = 1$, and $(\Theta_i, \Theta_{i+1}) = 1$, which is the $\hat{A}_{n-1}$ Dynkin diagram. The group of covering transformations of the n^{th} order cyclic unramified covering is $\mathbf{Z}_n$, and the action over the components (4.195) is given by

$$\begin{aligned} \Theta_i &\rightarrow \Theta_{i+1}, \\ \Theta_{n-1} &\rightarrow \Theta_0. \end{aligned} \quad (4.196)$$

4.10 Singularities of Type $\hat{D}_{n+4}$.

This case is a combination of the two previous examples. Through the same reasoning as above, the group $\mathcal{G}_\rho$ is given, for

$$S_\rho = \begin{pmatrix} -1 & -n \\ 0 & -1 \end{pmatrix}. \quad (4.197)$$

by

$$g(k, n_1, n_2) : (\rho, \lambda) \rightarrow (k + \rho, (-1)^k(\lambda + n_1 n \rho + n_2)). \quad (4.198)$$

Using a new variable $\sigma^2 = t$, what we get is a set of irreducible components $\Theta_0, \dots, \Theta_{2n}$, with the identifications $\Theta_i \rightarrow \Theta_{2n-i}$. In addition, we get the four fixed $\mathbf{Z}_2$ orbifold points described above. The singular fiber is then given by

$$\mathcal{C} = 2\Theta_0 + \dots + 2\Theta_n + \Theta^1 + \Theta^2 + \Theta^3 + \Theta^4, \quad (4.199)$$

with the intersections of the $\hat{D}_{n+4}$ affine diagram. It is easy to see that in this case we also get

$$(\mathcal{C})^2 = 0. \quad (4.200)$$

Defining the genus of the singular fiber by $\mathcal{C}^2 = 2g - 2$, we conclude that $g = 1$, for all singularities of Kodaira type. Notice that for rational singularities, characterized by non affine Dynkin diagrams of ADE type [44], we get self intersection $\mathcal{C}^2 = -2$, which corresponds to genus equal zero.

4.11 Decompactification and Affinization.

The general framework in which we are working in order to get four dimensional $N = 1$ gauge theories is that of M-theory compactifications on elliptically fibered Calabi-Yau fourfolds, in the limit $\text{Vol}(E) = 0$, with E the elliptic fiber. As described above, we can interpolate between $N = 2$ supersymmetry in three dimensions, and $N = 1$ in four dimensions, by changing the radius R through

$$\text{Vol}(E) = \frac{1}{R}. \quad (4.201)$$

The three dimensional limit then corresponds to $\text{Vol}(E) \rightarrow \infty$, and the four dimensional to $\text{Vol}(E) \rightarrow 0$. Now, we will work locally around a singular fiber of Kodaira $\hat{A}\hat{D}\hat{E}$ type. As we know, for the Calabi-Yau fourfold X ,

$$E \rightarrow X \xrightarrow{\Pi} B, \quad (4.202)$$

the locus C in B , where the fiber is singular, is of codimension one in B , i. e., of real dimension four. Let us now see what happens to the singular fiber in the three dimensional limit. In this case, we have $\text{Vol}(E) = \infty$. A possible way to represent this phenomenon is by simply extracting the point at infinity. In the case of $\hat{A}_{n-1}$ singularities, as described in previous subsection, taking out the point at infinity corresponds to decompactifying the

irreducible component Θ_0 , that was associated with the curve itself. As was clear in this case, we then pass from the affine diagram, $\hat{A}_{n-1}$, to the non affine, A_{n-1} . More generally, as the elliptic fibration we are considering possesses a global section, we can select the irreducible component we are going to decompactify as the one intersecting with the basis of the elliptic fibration. When we decompactify, in the $\text{Vol}(E) = 0$ limit, what we are doing, at the level of the fiber, is precisely compactifying the extra irreducible component, which leads to the affine Dynkin diagram.

4.12 M-Theory Instantons and Holomorphic Euler Characteristic.

Using the results of reference [99] a vertical instanton in a Calabi-Yau fourfold, of the type (4.202), will be defined by a divisor D of X , such that $\Pi(D)$ is of codimension one in B , and with holomorphic Euler characteristic

$$\chi(D, \mathcal{O}_D) = 1. \quad (4.203)$$

It is in case (4.203) that we have two fermionic zero modes [99], and we can define a superpotential contribution associated to D . For N , the normal bundle to D in X , which is locally a complex line bundle on D , we define the $U(1)$ transformation

$$t \rightarrow e^{i\alpha} t, \quad (4.204)$$

with t a coordinate of the fiber of N . The two fermionic zero modes have $U(1)$ charge equal one half. Associated to the divisor D , we can define a scalar field ϕ_D that, together with $\text{Vol}(D)$ defines the imaginary and real parts of a chiral superfield. Under $U(1)$ rotations (4.204), ϕ_D transforms as

$$\phi_D \rightarrow \phi_D + \chi(D)\alpha. \quad (4.205)$$

In three dimensions, this is precisely the transformation of the dual photon field as Goldstone boson [24]. However, transformation (4.205) has perfect sense, for vertical instantons, in the four dimensional decompactification limit.

Let us now consider an elliptically fibered Calabi-Yau fourfold, with singular fiber of $\hat{A}_{n-1}$ type, over a locus C of codimension one in B . We will assume that the singular fiber is constant over C . Moreover, in the geometrical engineering spirit, we will impose

$$h_{1,0}(C) = h_{2,0}(C) = 0 \quad (4.206)$$

and, thus, $\Pi_1(C) = 0$. This prevents us from having non trivial transformations on the fiber by going, on C , around closed loops, since all closed loops are contractible. In addition, we will assume, based on (4.206), that C is an Enriques surface. After imposing these assumptions, we will consider divisors D_i , with $i = 0, \dots, n-1$, defined by the fibering over C , in a trivial way, of the irreducible components Θ_i of the $\hat{A}_{n-1}$ singularity [126]. Ussing the Todd representation of the holomorphic Euler class [137],

$$\chi(D) = \frac{1}{24} \int_{D_i} c_1(\Theta_i) c_2(C) \quad (4.207)$$

we get, for C an Enriques surface,

$$\chi(D_i) = 1. \quad (4.208)$$

Interpreting now the t variable (4.204) on the fiber of the normal bundle N of D in X as the t variable used in our previous description of Kodaira singularities of type $\hat{A}_{n-1}$, we can derive the transformation law, under the $\mathbf{Z}_n$ subgroup of $U(1)$, of the scalar fields ϕ_{D_i} associated to these divisors. Namely, from (4.196) we get

$$\mathbf{Z}_n : \phi_{D_i} \rightarrow \phi_{D_{i+1}}, \quad (4.209)$$

with the $\mathbf{Z}_n$ transformation being defined by

$$t \rightarrow e^{2\pi i/n} t. \quad (4.210)$$

Using now (4.205), we get

$$\mathbf{Z}_n : \phi_{D_i} \rightarrow \phi_{D_i} + \frac{2\pi}{n}. \quad (4.211)$$

Combining (4.209) and (4.211) we get, modulo 2π ,

$$\phi_{D_j} = \frac{2\pi j}{n} + c, \quad (4.212)$$

with $j = 0, \dots, n-1$, and c a constant independent of j .

Let us now consider the divisor $\mathcal{D}$ obtained by fibering over C the singular fiber $\mathcal{C} = \sum_{j=0}^{n-1} \Theta_j$, defined in (4.195). In this case we need to be careful in order to compute (4.207). If we naively consider the topological sum of components Θ_j in (4.207), we will get the wrong result $\chi(\mathcal{D}) = n$. This result would be correct topologically, but not for the holomorphic Euler characteristic we are interested in. In fact, what we should write in (4.207) for $\int c_1(\sum_{j=0}^{n-1} \Theta_j)$ is $2(1 - g(\sum_{j=0}^{n-1} \Theta_j))$, with g the genus of the cycle (4.195),

as defined by $\mathcal{C}^2 = 2g - 2$, with $\mathcal{C}^2$ the self intersection of the cycle (4.195) which, as for any other Kodaira singularity, is zero. Thus, we get $g = 1$, and [131]

$$\chi(\mathcal{D}) = 0. \quad (4.213)$$

We can try to interpret the result (4.213) in terms of the fermionic zero modes of each component Θ_i , and the topology of the cycle. In fact, associated to each divisor D_i we have, as a consequence of (4.208), two fermionic zero modes. In the case of the $\hat{A}_{n-1}$ singularity, we can soak up all zero modes inside the graph, as shown in Figure 1,




Figure 3: Soaking up of zero modes for $\hat{A}_{n-1}$.

where from each node, representing one Θ_i , we have two fermionic zero mode lines. The soaking up of fermionic zero modes represented in the figure is an heuristic interpretation of the result (4.213).

4.13 θ -Parameter and Gaugino Condensates.

We will, in this section, only consider singularities of $\hat{A}_{n-1}$ and $\hat{D}_{n+4}$ type. In both cases, and for each irreducible component Θ_i , we get a divisor D_i , with $\chi(D_i) = 1$. Associated to this divisor, we can get a superpotential term of the order [99]

$$\int d^2\theta e^{-(V(D_i)) + i\phi_{D_i}}, \quad (4.214)$$

where $V(D_i)$ means the volume of the divisor D_i . As explained above, we are using vertical instanton divisors D_i , defined by a trivial fibering of Θ_i over the singular locus $C \subset B$, satisfying conditions (4.206). In order to get the four dimensional $N = 1$ limit, we will take the limit $\text{Vol}(E) = \frac{1}{R} \rightarrow 0$. Since the singular fibers are, topologically, the union of irreducible components (see (4.195) and (4.199)), we can write

$$\text{Vol}(\Theta_i) = \frac{1}{R\text{Cox}}, \quad (4.215)$$

with Cox the Coxeter number of the corresponding singularity, which equals the total number of irreducible components. Therefore, we will define $\text{Vol} (D_i)$ as

$$\text{Vol} (D_i) = \lim_{R \rightarrow \infty} \text{Vol} (C) \frac{1}{R \text{Cox}}. \quad (4.216)$$

If we first consider the $N = 2$ supersymmetric three dimensional theory obtained by compactifying M-theory on the Calabi-Yau fourfold X , i. e., in the limit $R \rightarrow 0$, we know that only the divisor Θ_0 , for the $\hat{A}_{n-1}$ case, is decompactified, passing from the affine diagram describing an elliptic singularity to the non affine diagram describing a rational, Artin like, singularity [44]. In that case, the volumes of the Θ_i components, for $i \neq 0$, are free parameters, corresponding to the Coulomb branch of the $N = 2$ three dimensional theory. In the three dimensional theory, the factor $\text{Vol} (C)$ corresponds to the bare coupling constant in three dimensions,

$$\text{Vol} (C) = \frac{1}{g_3^2}, \quad (4.217)$$

and $\text{Vol} (D_i) = \frac{1}{g_3^2} \chi_i$, for $i \neq 0$, with χ_i the three dimensional Coulomb branch coordinates. In the four dimensional case, we must use (4.216), that becomes

$$\text{Vol} (D_i) = \lim_{R \rightarrow \infty} \frac{1}{g_3^2} \frac{1}{R \text{Cox}} = \frac{1}{g_4^2 \text{Cox}}. \quad (4.218)$$

Let us now concentrate on the $\hat{A}_{n-1}$ case, where $\text{Cox} = n$. Using (4.214) we get the following superpotential for each divisor D_j ,

$$\exp - \left(\frac{1}{g_4^2 n} + i \left(\frac{2\pi j}{n} + c \right) \right). \quad (4.219)$$

Let us now fix the constant c in (4.219). In order to do that, we will use the transformation rules (4.205). From the four dimensional point of view, these are the transformation rules with respect to the $U(1)_R$ symmetry. From (4.205) we get, that under $t \rightarrow e^{i\alpha} t$,

$$\sum_{i=0}^{n-1} \phi_{D_i} \rightarrow \sum_{i=0}^{n-1} \phi_{D_i} + n\alpha. \quad (4.220)$$

This is precisely the transformation rule under $U(1)_R$ of the $N = 1$ θ -parameter,

$$\theta \rightarrow \theta + n\alpha. \quad (4.221)$$

In fact, (4.220) is a direct consequence of the $U(1)$ axial anomaly equation: if we define θ as

$$\frac{\theta}{32\pi^2} F\tilde{F}, \quad (4.222)$$

the anomaly for $SU(n)$ is given by

$$\partial_\mu j_5^\mu = \frac{n}{16\pi^2} F\tilde{F}. \quad (4.223)$$

The factor 2, differing (4.222) from (4.223), reflects the fact that we are assigning $U(1)_R$ charge $\frac{1}{2}$ to the fermionic zero modes. Identifying the θ -parameter with the topological sum $\sum_{i=0}^{n-1} \phi_{D_i}$ we get that the constant c in (4.219) is simply

$$c = \frac{\theta}{n}, \quad (4.224)$$

so that we then finally obtain the superpotential

$$\exp - \left(\frac{1}{g_4^2 n} + i \left(\frac{2\pi j}{n} + \frac{\theta}{n} \right) \right) \simeq \Lambda^3 e^{2\pi i j/n} e^{i\theta/n}, \quad (4.225)$$

with $j = 0, \dots, n-1$, which is the correct value for the gaugino condensate.

Let us now try to extend the previous argument to the $\hat{D}_{n+4}$ type of singularities. Defining again the four dimensional θ -parameter as the topological sum of ϕ_{D_i} for the whole set of irreducible components we get, for the cycle (4.199), the transformation rule

$$\theta \rightarrow \theta + \text{Cox} \cdot \alpha \quad (4.226)$$

where now the Coxeter for $\hat{D}_{n+4}$ is $2n+6$. Interpreting $\hat{D}_{n+4}$ as $O(N)$ gauge groups, with $N = 2n+8$, we get $\text{Cox}(\hat{D}_{n+4}) = N-2$. Since θ is defined modulo 2π we get that for $\hat{D}_{n+4}$ singularities the value of ϕ_{D_i} , for any irreducible component, is

$$\frac{2\pi k}{N-2} + \frac{\theta}{N-2}, \quad (4.227)$$

with $k = 1, \dots, N-2$. However, now we do not know how to associate a value of k to each irreducible component Θ_i of the $\hat{D}_{n+4}$ diagram. Using (4.227), we get a set of $N-2$ different values for the gaugino condensate for $O(N)$ groups:

$$\exp \left(-\frac{1}{g_4^2(N-2)} + i \left(\frac{2\pi k}{N-2} + \frac{\theta}{N-2} \right) \right), \quad (4.228)$$

with $k = 1, \dots, N-2$. However, we still do not know how to associate to each Θ_i a particular value of k . A possibility will be associating consecutive values of k to components with non vanishing intersection; however, the topology of diagrams of type D

prevents us from doing that globally. Notice that the problem we have is the same sort of puzzle we find for $O(N)$ gauge groups, concerning the number of values for $\langle \lambda \lambda \rangle$, and the value of the Witten index, which in diagrammatic terms is simply the number of nodes of the diagram. In order to unravel this puzzle, let us consider more closely the way fermionic zero modes are soaked up on a $\hat{D}_{n+4}$ diagram. We will use the cycle (4.199); for the components Θ^1 to Θ^4 , associated to the $\mathbf{Z}_2$ orbifold points, we get divisors with $\chi = 1$. Now, for the components $2\Theta_0, \dots, 2\Theta_n$ we get, from the Todd representation of the holomorphic Euler characteristic,

$$\chi = 4. \quad (4.229)$$

The reason for this is that the cycle 2Θ , with $\Theta^2 = -2$, has self intersection -8 . Of course, (4.229) refers to the holomorphic Euler characteristic of the divisor obtained when fibering over C any of the cycles $2\Theta_i$, with $i = 0, \dots, n$. Equation (4.229) implies 8 fermionic zero modes, with the topology of the soaking up of zero modes of the $\hat{D}_{n+4}$ diagram, as represented in Figure 2.




Figure 4: Soaking up of zero modes for $\hat{D}_{n+4}$.

Notice that the contribution to χ of 2Θ is different from that of $(\Theta_1 + \Theta_2)$, with $(\Theta_1 \cdot \Theta_2) = 0$; namely, for the first case $\chi = 4$, and $\chi = 2$ for the second. For the $\hat{D}_{n+4}$ diagram, we can define: *i*) The Witten index $\text{tr} (-1)^F$, as the number of nodes, i. e., $5 + n$; *ii*) The Coxeter number, which is the number of irreducible components, i. e., $2n + 6$ and *iii*) The number of intersections as represented by the dashed lines in Figure 4, i. e., $8 + 4n$. From the point of view of the Cartan algebra, used to define the vacuum configurations in [22], we can only feel the number of nodes. The θ -parameter is able to feel the Coxeter number; however, we now find a new structure related to the intersections

of the graph. In the Witten index case, the nodes corresponding to cycles $2\Theta_i$, with $i = 0, \dots, n$ contribute with one, in the number of $\langle \lambda\lambda \rangle$ values with two, and in the number of intersections with four. This value four calls for an orientifold interpretation of these nodes. The topological definition of the θ -parameter implicitly implies the split of this orientifold into two cycles, a phenomena recalling the F-theory description [138] of the Seiberg-Witten splitting [35]. Assuming this splitting of the orientifold, the only possible topology for the soaking up of zero modes is the one represented in Figure 3, where the “splitted orientifold” inside the box is associated to four zero modes, corresponding to $\chi = 2$ for a cycle $\Theta_1 + \Theta_2$, with $\Theta_1.\Theta_2 = 0$.




Figure 5: Orientifold splitting.

On the other hand, each node surrounded by a circle in Figure 5 represents itself the disconnected sum of two non singular rational curves; thus, we represent each “orientifold” mode by four rational curves, with the intersections depicted inside the box of Figure 5. When we forget about internal lines in Figure 5, we get the cyclic $\mathbf{Z}_{2n+6} \equiv \mathbf{Z}_{N-2}$ structure of equation (4.228). It is clear that much more is necessary in order to reach a complete description of the $O(N)$ vacuum structure.

4.14 Domain Walls and Intersections.

The discussion in the previous section already raises the problem known as θ -puzzle. In fact, and discussing again only the $SU(n)$ case, the transformation law (4.221) together with the very definition for the θ -angle as the topological sum $\sum_{i=0}^{n-1} \phi_{D_i}$ would imply that θ is the scalar field $\phi_{\mathcal{D}}$ of the 6-cycle associated to the $\hat{A}_{n-1}$ cycle, $\mathcal{C} = \sum_{i=0}^{n-1} \Theta_i$. On the basis of (4.205), this will be equivalent to saying that $\chi(\mathcal{D}) = n$, instead of zero. This is, in mathematical terms, the θ -puzzle. The mathematical solution comes from the fact

that $\chi(\mathcal{D}) = 0$. In this section we will relate this result, on the value of the holomorphic Euler chareacteristic, to the appearance of domain walls [132, 133, 134]. To start with, let us consider a cycle $\mathcal{C} = \Theta_1 + \Theta_2$, with $(\Theta_1.\Theta_2) = 1$. The self intersection can be expressed as

$$(\mathcal{C}.\mathcal{C}) = -2 - 2 + 2, \quad (4.230)$$

where the -2 contributions come from Θ_1^2 and Θ_2^2 , and the $+2$ comes from the intersection between Θ_1 and Θ_2 . As usual, we can consider $\mathcal{C}$ trivially fibered on an Enriques surface. The holomorphic Euler chracteristic of the corresponding six cycle can be written as

$$\chi = \frac{1}{2}(-\mathcal{C}^2). \quad (4.231)$$

Using now the decomposition (4.230) we get two contributions of one, coming from the components Θ_1 and Θ_2 , considered independently, and a contribution of -1 from the intersection term $+2$ in (4.230). In this sense, the intersection term can be associated to two fermionic zero modes, and net change of chiral charge oposite to that of the Θ_i components. When we do this for the cycle $\mathcal{C}$ of $\hat{A}_{n-1}$ singularities, we get that each intersection is soaking up two zero modes, leading to the result that $\chi(\mathcal{C}) = 0$. A graphical way to represent equation (4.230) is presented in Figure 4.




Figure 6: Intersection term.

Now, we will wonder about the physical interpretation of the intersection terms leading to $\chi(\mathcal{C}) = 0$ for all Kodaira singularities. The simplest, and most natural answer, is certainly domain walls extending between different vacua, or values of $\langle \lambda \lambda \rangle$.

From the point of view of zero mode counting, the “intersection term” behaves effectively as an anti-instanton with two fermionic zero modes. One of these fermionic zero modes, let us say $\psi_{j,j+1}$, is associated to the intersection of Θ_j with Θ_{j+1} and the other $\psi_{j+1,j}$ with the intersection of Θ_j and Θ_{j+1} . Thus, extending naively the computation done for irreducible components, the contribution of the black box in Figure 4 should be of the order

$$\Lambda^3 e^{2\pi i j/n} (1 - e^{2\pi i/n}). \quad (4.232)$$

In result (4.232), interpreted as the contribution of the intersection term, the most surprising fact is the appearance of Λ^3 , since now we are geometrically considering simply a point; the factor Λ^3 in the computation of the gaugino condensate comes from the volume of the divisor. In the same way as we interpret M-theory instantons as fivebranes wrapped on the six-cycles used to define the instanton, we can think of the intersection terms as fivebranes wrapped on the cycle $C \times \{(\Theta_i, \Theta_{i+1})\}$, i. e., the product of the singular locus C and the intersection point. The fivebrane wrapped on this cycle defines, in four dimensions, a domain wall, let us say intertwining between the vacua i , at $x_3 = +\infty$, and the vacua $i + 1$, at $x_3 = -\infty$, where the coordinate x_3 is identified with the unwrapped direction. It is in this sense that we should use (4.232) to define the energy density, or tension, of the domain wall. In the four dimensional limit, $\text{Vol}(E)$ goes zero as $\frac{1}{R}$; moreover, the local engineering approach works in the limit where the volume of the singular locus C is very large, so that we can very likely assume that intersection terms behave like (4.232), with Λ^3 , but only in the four dimensional limit. Cyclicity of the $\hat{A}_{n-1}$ diagram allows us to pass from the j to the $j + 1$ vacua in two different ways: $n - 1$ steps, or a single one. The sum of both contributions should define the physical domain wall; thus the energy density will behave as

$$n\Lambda^3|e^{2\pi ij/n}(1 - e^{2\pi i/n})|. \quad (4.233)$$

The extension of the previous argument to the case of $O(N)$ groups is certainly more involved, due to the topology of the $\hat{D}$ diagram, and the presence of orientifolds. It would certainly be interesting studying the interplay between orientifolds and domain walls in this case.

Finally, we will say some words on the QCD string. In reference [117], the geometry of QCD strings is intimately related to the topological fact that

$$H_1(Y/Z; \mathbf{Z}) = \mathbf{Z}_n, \quad (4.234)$$

where Σ is a rational curve associated to the configuration of fourbranes, and $Y = S^1 \times \mathbf{R}^5$ is the ambient space where Σ is embedded (see [117] for details). The QCD string is then associated to a partially wrapped membrane on a non trivial element of $H_1(Y/\Sigma; \mathbf{Z})$. Recall that $H_1(Y/\Sigma; \mathbf{Z})$ is defined by one-cycles in Y , with boundary on Σ . The previous discussion was done for $SU(N)$ gauge groups. Using our model of $\hat{A}_{n-1}$ singularities, described in section 2, the analog in our framework of (4.234) is equation (4.187). Then we can, in the same spirit as in reference [117], associate the QCD string to paths going from p_k to p_{k+1} , where p_k are the intersection points,

$$\Theta_k \cdot \Theta_{k-1} = p_k. \quad (4.235)$$

Geometrically, it is clear that the tension of this QCD string is the square root of the domain wall tension. By construction, the QCD string we are suggesting here ends on domain walls, i. e., on intersection points.

To end up, let us include some comments on the existence of extra vacua, as suggested in [135]. It is known that the strong coupling computation of $\langle \lambda\lambda \rangle$ does not coincide with the weak coupling computation; more precisely[136],

$$\langle \lambda\lambda \rangle_{\text{sc}} < \langle \lambda\lambda \rangle_{\text{wc}} . \quad (4.236)$$

In the framework of M-theory instanton computations, the numerical factors will depend in particular on the moduli of complex structures of the Calabi-Yau fourfold. In the strong coupling regime we must consider structures preserving the elliptic fibration structure and the Picard lattice. In the weak coupling regime, where the computation is performed in the Higgs phase, the amount of allowed complex structures contributing to the value of $\langle \lambda\lambda \rangle$ is presumably larger. Obviously, the previous argument is only suggesting a possible way out of the puzzle (4.236).

Equally, at a very speculative level, the extra vacua, with no chiral symmetry breaking, could be associated to the cycle $\mathcal{D}$ defining the singular fiber, a cycle that we know leads to $\chi = 0$, and therefore does not produce any gaugino condensate. Notice that any other cycle with $\chi \neq 0$ will lead, if clustering is used, to some non vanishing gaugino condensates, so that $\mathcal{D}$ with $\chi = 0$ looks like a possible candidate to the extra vacua suggested in [135]. If this argument is correct this extra vacua will appear for any Kodaira singularity i. e. in any ADE $N = 1$ four dimensional gauge theory.

It is important to stress that the θ -puzzle is not exclusive of $N = 1$ gluodynamics. In the $N = 0$ case the Witten-Veneziano formula [139, 140] for the η' mass also indicates a dependence of the vacuum energy on θ in terms of $\frac{\theta}{N}$, which means a set of entangled "vacuum" states. In our approach to $N = 1$ the origin of this entanglement is due to the fact that $\chi = 0$ for the singular cycle. In fact, $\chi(\mathcal{D}) = 0$ means that the set of divisors D_i , plus the intersections, i. e., the domain walls, are invariant under $U(1)$, as implied by equation (4.205). If we naively think of something similar in $N = 0$ and we look for the origin of vacuum entanglement in intersections we maybe should think in translating the topology of intersections into topological properties of abelian projection gauges [58].

A M(atrix) Theory.

A.1 The Holographic Principle.

The holomorphic principle was originally suggested by 't Hooft in [141]. Let us work first in four dimensional spacetime. Let $S^{(2)}$ be a surface with the topology of two sphere in $\mathbf{R}^3$, and let us wonder about how many orthogonal quantum states can $S^{(2)}$ contain: we will find an upper bound for this number of states. In order to do this, we will use the Bekenstein-Hawking relation between the entropy and the horizon area of the black hole. Let us then call $\mathcal{N}$ the number of states inside $S^{(2)}$; the entropy can be defined as

$$\exp S = \mathcal{N}. \quad (\text{A.1})$$

If S^2 is the horizon of a black hole, we have [142]

$$S \sim \frac{1}{4} \frac{A}{l_p^2}, \quad (\text{A.2})$$

with A the area of the horizon, in Planck length units. Now, let us translate all the physical information contained inside the $S^{(2)}$ surface, in terms of states of q -bits, defined as quantum systems of two states. The number n of q -bits we need is given by

$$N = 2^n. \quad (\text{A.3})$$

Using (A.1) to (A.3) we then get

$$n = \frac{1}{4 \ln 2} \frac{A}{l_p^2}, \quad (\text{A.4})$$

which is essentially the number of cells, of area l_p^2 , covering the surface $S^{(2)}$. What we learn from this is that all three dimensional physics inside $S^{(2)}$ can be described using states of q -bits, living on the two dimensional surface $S^{(2)}$. We will call these q -bits the holographic degrees of freedom. What we need now is the two dimensional dynamics governing these two dimensional degrees of freedom, able to reproduce, in holographic projection, the three dimensional physics taking place $S^{(2)}$. We can even consider, instead of $S^{(2)}$, an hyperplane of dimension two, dividing space into two regions. The extension, to this extreme situation, of the holomorphic principle, will tell us that the $3 + 1$ dynamics can be described in terms of some $2 + 1$ dynamics for the holomorphic degrees of freedom living on the hypersurface.

This picture of the holographic principle allows to introduce M(atrix) theory [143] as the holographic projection of M-theory. In this case, we will pass from eleven dimensional

to ten dimensional physics. What we will need, in order to formulate M(atrix) theory, will be

- i) An explicit definition of the holographic projection.
- ii) Identifying the holomorphic degrees of freedom.
- iii) Providing a ten dimensional dynamics for these degrees of freedom.

The conjectured answer in [143] to *i*), *ii*) and *iii*) are

- i) The infinite momentum frame.
- ii) D-0branes as degrees of freedom.
- iii) The dynamics is implemented through the worldvolume of the lagrangian of D-0branes.

These set of conjectures define M(atrix) theory at present. The idea of the infinite momentum frame is boosting in the eleventh direction in eleven dimensional spacetime, in such a way that p_{11} , the eleventh component of the momentum, becomes larger than any scale in the problem. In this frame, we associate, with an eleven dimensional massless system of momentum $\vec{p} = (p_{11}, p_{\perp})$, a ten dimensional galilean system with mass p_{11} , and energy

$$E = \frac{p_{\perp}^2}{2p_{11}}. \quad (\text{A.5})$$

If we introduce an infrared cut off, by compactifying the eleventh dimension on a circle S^1 , of radius R , the p_{11} is measured in units of $\frac{1}{R}$. Then,

$$p_{11} = \frac{n}{R}. \quad (\text{A.6})$$

We will interpret n as the number of partons which are necessary to describe a system with value of p_{11} given by (A.6). These partons are the ten dimensional degrees of freedom we are going to consider as holographic variables.

Using (A.4), we can define the ten dimensional size of an eleven dimensional massless particle, in eleven dimensions, with some given p_{11} . In fact, $n = p_{11}R$ is the number of holographic degrees of freedom which, by (A.4), means that

$$\frac{r^9}{l_p^9} \sim p_{11}R, \quad (\text{A.7})$$

and the radius r , characterizing the size, will be $(p_{11}R)^{1/9}l_p$. Now, we can look for objects in ten dimensions with mass equal to the mass of a parton, i. e., $\frac{1}{R}$. Natural candidates are D-0branes. From this, it seems natural to conjecture that the worldvolume dynamics of D-0branes will be a good candidate for the holographic description of M-theory.

M(atrrix) theory would hence simply be defined as the worldvolume theory of D-0branes. As for any other type of D-branes, this worldvolume theory is defined as the dimensional reduction down to 0+1 dimensions of ten dimensional Yang-Mills with $N = 1$ supersymmetry. If we consider a set of N D-0branes, we have to introduce matrices X^i , with $i = 1, \dots, 9$. As usual, the diagonal part of this matrices can be interpreted in terms of the classical positions of the N D-0branes, and the off diagonal terms as representing the exchange of open strings. Thus, the worldvolume lagrangian we get for N D-0branes is $U(N)$ Yang-Mills quantum mechanics. Using units in which $\alpha' = 1$, the bosonic part of this lagrangian is simply

$$\mathcal{L} = \frac{1}{2g} [\text{tr} \dot{X}^i \dot{X}^i - \frac{1}{2} \text{tr} [X^i, X^j]^2], \quad (\text{A.8})$$

in units where $l_p = 1$, and with g the string coupling constant. In this definition we have simply consider D-0branes in a ten dimensional type IIA string theory. However, we know that D-0branes are in fact Kaluza-Klein modes of an eleven dimensional theory named M-theory. In the eleven dimensional spacetime, the D-0branes have a momentum p_{11} given by

$$p_{11} = \frac{1}{R}, \quad (\text{A.9})$$

with R the radius of the eleventh dimension. The way to relate (A.8) to the physics of the partons defined in the infinite momentum frame is observing that the kinetic term in (A.8) coincides with the equation (A.5) for p_{11} given in (A.9). In fact, using the relation

$$R = gl_s, \quad (\text{A.10})$$

and choosing units where $l_s = 1$, we notice that (A.8) is precisely the galilean lagrangian for particles of mass $\frac{1}{g}$. Thus, we will interpret the worldvolume dynamics of D-0branes, (A.8), as the infinite momentum frame of the M-theory D-0branes.

Our main task now will consist in deriving the brane spectrum directly from the M(atrrix) lagrangian (A.8), interpreting the different branes as collective excitations of D-0branes. In order to achive this, it will be necessary to work in the $N \rightarrow \infty$ of (A.8). Using the relation

$$l_s = g^{-1/3}l_p, \quad (\text{A.11})$$

between the string length, and the Plank scale, we can pass to Plank units by defining $Y = \frac{X}{g^{1/3}}$. In Y variables, and with $l_p = 1$, we get

$$\mathcal{L} = \text{tr} \left[\frac{1}{2R} D_t Y^i D_t Y^i - \frac{1}{4} R [Y^i, Y^j]^2 \right], \quad (\text{A.12})$$

where $D_t = \partial_t + iA$, with A equal the A_0 piece of the ten dimensional Yang-Mills theory. In order to get (A.8), going to the temporal gauge $A_0 = 0$ is all what is needed.

Now, some of the ingredients introduced in chapter I will be needed; namely, the matrices P and Q defined in (1.87). In terms of this basis of matrices, any matrix Z can be written as

$$Z = \sum_{n,m=1}^N z_{n,m} P^n Q^m. \quad (\text{A.13})$$

Taking into account that

$$PQ = QPe^{2\pi i/N}, \quad (\text{A.14})$$

we can define

$$P = e^{i\hat{p}}, \quad Q = e^{i\hat{q}}, \quad (\text{A.15})$$

with

$$[\hat{p}, \hat{q}] = \frac{2\pi i}{N}. \quad (\text{A.16})$$

Replacing (A.15) in (A.13) we get

$$Z = \sum_{n,m} z_{n,m} e^{in\hat{p}} e^{im\hat{q}}, \quad (\text{A.17})$$

which looks like the Fourier transform of a function $Z(p, q)$. The only difference is that this function is defined on a quantum phase space defined by $\hat{p}$ and $\hat{q}$ variables satisfying (A.16). In the $N \rightarrow \infty$ limit, we can interpret this quantum space as classical, since in this limit $[\hat{p}, \hat{q}] = 0$. Thus, in the $N \rightarrow \infty$ limit, the matrices can be replaced by functions $Z(p, q)$, as defined by (A.17). The matrix operations become, in this limit,

$$\begin{aligned} \text{tr} Z &\rightarrow N \int Z(p, q) dp dq, \\ [X, Y] &\rightarrow \frac{1}{N} [\partial_q X \partial_p Y - \partial_q Y \partial_p X], \end{aligned} \quad (\text{A.18})$$

i. e., the conmutator becomes the Poisson bracket. Now, we can use (A.18) in (A.12); what we then get is

$$\mathcal{L} = \frac{p_{11}}{2} \left[\int dp dq \dot{Y}^i(p, q) \dot{Y}^i(p, q) - \frac{1}{P_{11}} \int dp dq [\partial_q Y^i \partial_p Y^j - \partial_q Y^j \partial_p Y^i] \right], \quad (\text{A.19})$$

where $p_{11} = \frac{N}{R}$. The interest of (A.19) is that this result coincides with the eleven dimensional lagrangian for the eleven dimensional supermembrane in the light cone frame. Notice that i in (A.19) goes from 1 to 9, which can be interpreted as the transversal directions to the supermembrane worldvolume. The previous result is already a good indication of the consistency of M(atrix) theory as a microscopic description of M-theory. Next, we will try to define toroidal compactifications of (A.8).

A.2 Toroidal Compactifications.

The definition of toroidal compactifications [145] of M(atrix) theory is quite simple. We will consider the worldvolume of lagrangian of D-0branes, starting with ten dimensional $N = 1$ supersymmetric Yang-Mills in $\mathbf{R}^9 \times S^1$. In order to clear up the procedure, we will keep all indices for a while, so that we will write

$$X_{k,l}^i \tag{A.20}$$

for the matrix X^i . The indices k and l will hence label different D-0branes. Now, if we force D-0branes to live in $\mathbf{R}^9 \times S^1$, and interpret S^1 as

$$\mathbf{R}/\Gamma, \tag{A.21}$$

with Γ a one dimensional lattice defined by a vector $\vec{e} = 2\pi R$, we can think of copies of each D-0brane, parametrized by integers n , depending on the cell of $\mathbf{R}/\Gamma$ where they are. Then, (A.20) should be changed to

$$X_{k,m;l,m}^i. \tag{A.22}$$

We can now forget about the indices k and l , to write X_{nm}^i , where n and m are integers. The lagrangian (A.8) then becomes

$$\mathcal{L} = \frac{1}{2g} [\text{tr} \dot{X}_{mn}^i \dot{X}_{mn}^i + \frac{1}{2} \text{tr} (X_{mq}^i X_{qn}^j - X_{mq}^j X_{qn}^i) (X_{nr}^i X_{rm}^j - X_{nr}^j X_{rm}^i)]. \tag{A.23}$$

Now, we should impose symmetry with respect to the action of Γ , which implies

$$\begin{aligned} X_{mn}^i &= X_{m-1 \ n-1}^i \quad i > 1, \\ X_{mn}^1 &= X_{m-1 \ n-1}^1 \quad m \neq n, \\ X_{mn}^1 &= 2\pi R \mathbf{I} + X_{m-1 \ n-1}^1. \end{aligned} \tag{A.24}$$

The meaning of (A.24) is that the coordinate X^1 is periodic, so that the difference in X^1 for n D-0branes, and $n + 1$ D-0branes, is simply the length of the compactified direction.

Using (A.24), matrices can be simply labelled by one index, $X_{0,n}^i \equiv X_n^i$, and the lagrangian (A.23) becomes

$$\mathcal{L} = \frac{1}{2g} \left[\sum_{i=1}^9 \text{tr} \dot{X}_n^i \dot{X}_{-n}^i - \sum_{j=2}^9 \text{tr} S_n^j (S_n^j)^+ - \frac{1}{2} \sum_{j,k=2}^9 \text{tr} T_n^{jk} (T_n^{jk})^+ \right], \quad (\text{A.25})$$

where

$$\begin{aligned} S_n^j &= \sum_q ([X_q^1, X_{n-q}^j]) - 2\pi R n X_n^j, \\ T_n^{jk} &= \sum_q [X_q^j, X_{n-q}^k]. \end{aligned} \quad (\text{A.26})$$

Once we get lagrangian (A.21), we can compare it with the worldvolume lagrangian for D-1branes. In fact, for D-1branes the worldvolume lagrangian is 1 + 1 dimensional super Yang-Mills theory, with gauge fields A^1 and A^0 , and matter fields Y^j (with $j = 2, \dots, 9$) in the adjoint representation. We can then work in the temporal gauge, fixing $A^0 = 0$. On the other hand, performing T-duality on S^1 takes form D-0branes to D-1branes. Hence, on the dual S^1 the worldvolume lagrangian for D-1branes should coincide with that of D-0branes in $\mathbf{R}^9 \times S^1$, i. e, with lagrangian (A.25). The D-1brane worldvolume lagrangian in the dual S^1 , with radius $R' = \frac{1}{2\pi R}$,

$$\mathcal{L} = \int dx \frac{dt}{2\pi R'} \frac{1}{2g} \left[\text{tr} \dot{Y}^i \dot{Y}^i + \text{tr} \dot{A}^1 \dot{A}^1 + \frac{1}{2} \text{tr} [Y^i, Y^j]^2 - \text{tr} [\partial_1 Y^i - i[A^1, Y^i]]^2 \right], \quad (\text{A.27})$$

can be compared with (A.26) if we just interpret X_n^i as the Fourier modes of $Y^i(x)$, and X_n^1 as the Fourier modes of $A^1(x)$:

$$\begin{aligned} A^1(x) &= \sum_n e^{inx/R'} X_n^1, \\ Y^i(x) &= \sum_n e^{inx/R'} X_n^i. \end{aligned} \quad (\text{A.28})$$

Hence, we can readily induce the following result: M(atrix) theory compactified on T^d is equivalent to $d+1$ supersymmetric Yang-Mills on the dual $\hat{T}^d \times \mathbf{R}$, with $\mathbf{R}$ standing for the time direction, and the supersymmetric Yang-Mills theory defined through dimensional reduction from $N = 1$ ten dimensional Yang-Mills theory. This is a surprising result, connecting M(atrix) compactifications with Yang-Mills theories, a relation with far reaching consequences, some of which we will consider in what follows.

A.3 M(atrix) Theory and Quantum Directions.

We can then represent M(atrix) theory, compactified on T^d , as supersymmetric Yang-Mills theory defined on $\hat{T}^d \times \mathbf{R}$, i. e., as the worldvolume lagrangian of d D-branes wrapped on the dual torus, $\hat{T}^d$. Let us then work out some simple cases. We will first compactify M(atrix) on T^4 , which will be an interesting case concerning the U-duality symmetry.

Let L_i be the lengths of T^4 . The dual torus $\hat{T}$ will then be defined with sides of length

$$\Sigma_i = \frac{l_s^2}{L_i}, \quad (\text{A.29})$$

with l_s the string length. In terms of the eleven dimensional Planck scale, l_P , we have

$$\frac{l_P^3}{R} = l_s^2, \quad (\text{A.30})$$

and therefore

$$\Sigma_i = \frac{l_P^3}{L_i R}. \quad (\text{A.31})$$

Let us now consider the infinite momentum frame energy of a state with one unit of p_{11} , and one unit of momentum in some internal direction, L_i ,

$$E = \frac{p_\perp}{2P_{11}} = \frac{R}{2L_i^2}. \quad (\text{A.32})$$

This state corresponds, in supersymmetric Yang-Mills, to a gauge configuration with a non trivial Wilson line $A(C_i)$ (recall that in the toroidal compactification the compactified components X^i behave as Yang-Mills fields. This non trivial Wilson line means a flux through C_i . This energy is given by

$$\frac{g_{SYM}^2 \Sigma_i^2}{\Sigma_1 \Sigma_2 \Sigma_3 \Sigma_4}. \quad (\text{A.33})$$

Identifying (A.32) and (A.33) we get

$$g_{SYM}^2 = \frac{R}{2L_i^2} \frac{\Sigma_1 \Sigma_2 \Sigma_3 \Sigma_4}{\Sigma_i^2}. \quad (\text{A.34})$$

Using (A.31) we get

$$g_{SYM}^2 = \frac{R^3 \Sigma_1 \Sigma_2 \Sigma_3 \Sigma_4}{2l_P^6} = \frac{l_P^6}{2L_1 L_2 L_3 L_4 R}, \quad (\text{A.35})$$

which means that g^2 , as expected in $4 + 1$ dimensions, has units of length.

From the definition of M(atrix) compactifications, we expect that M(atrix) on T^4 will reproduce type IIA string theory on T^4 that, has been derived in chapter III, is invariant under the U-duality group, $Sl(5, \mathbf{Z})$. Thus, our task is to unravel this U-duality invariance, considering supersymmetric Yang-Mills on $\hat{T}^4 \times \mathbf{R}$. From (A.35), we observe a clear $Sl(4, \mathbf{Z})$ invariance of the gauge theory. These transformations exchange all Σ_i , leaving their product invariant. In order to extend this symmetry to $Sl(5, \mathbf{Z})$, an extra dimension Σ_5 needs to be defined. A way to do this is using as such direction the coupling constant itself in $4 + 1$ directions that, as can be clearly seen from (A.35), has dimensions of length. In this way, we can think that M(atrix) on T^4 is described by a $5 + 1$ dimensional theory, with space dimensions a torus T^5 , of dimensions Σ_i , with $i = 1, 2, 3, 4$, and $\Sigma_5 = \frac{l_p^6}{L_1 L_2 L_3 L_4 R}$. This is exactly the same picture we have in M-theory, understood as the strong coupled limit of type IIA string theory. There, we associated the RR D-0branes with Kaluza-Klein modes of the extra dimension. In the gauge theory context we should look for objects in $4 + 1$ dimensions, that can be interpreted as Kaluza-Klein modes of the extra dimension required by U -duality. As candidates to these states, we can use instantons. Instantons are associated with the Π_3 homotopy group of the gauge group so that, in $4 + 1$ dimensions, they look like particles. Moreover, their mass is given by $\frac{1}{g^2}$, with the gauge coupling constant (recall that $\frac{1}{g^2}$ is the action for the instanton in $3 + 1$ dimensions). Therefore, using (A.35), we get the desired result, namely that instantons are the Kaluza-Klein modes of the extra dimension.

We can, in fact, try to understand what kind of dynamics is playing the role here, using string theory language. The supersymmetric Yang-Mills theory on T^4 , with gauge group $U(N)$, can be interpreted as the worldvolume lagrangian for N fourbranes of type IIA, wrapped around T^4 . In M-theory, we can interpret this fourbranes as fivebranes partially wrapped in around the internal eleventh dimension. When we move to strong coupling, we open the extra direction, and we effectively get a $5 + 1$ dimensional gauge theory. If this is the correct picture, we can check it by comparison of the mass of the instanton and the expected mass of the wrapped around T^4 and the internal eleventh dimension. The energy of the fivebrane would then be

$$E = \frac{L_1 L_2 L_3 L_4 R}{l_p^6}, \quad (\text{A.36})$$

which is exactly the mass of the instanton,

$$\frac{1}{g^2} \equiv \frac{1}{\hat{g}} \frac{L_1 L_2 L_3 L_4 R}{l_p^6}. \quad (\text{A.37})$$

In order to understand the effect described above, it would be convenient to discuss briefly the scales entering the theory. Using relations (A.32) to (A.35), and $gl_s = R$ we get, for generic dimension d ,

$$g_{SYM}^2 = \frac{R^{3-d} l_s^{3d-6}}{L_i^d g^{d-3}}. \quad (\text{A.38})$$

It is clear from (A.38) that for $d \leq 3$ the limit of string coupling constant equal zero gives a weak coupled supersymmetric Yang-Mills theory. However, a barrier appears in $d = 4$. In fact, for $d \geq 4$ the limit $g \rightarrow 0$ leads to strong coupling in the field theory. One of these strong coupling effects is the generation of the quantum dimension needed for U -duality.

Acknowledgments

This work is partially supported by European Community grant ERBFMRXCT960012, and by grant AEN-97-1711.

References

- [1] P. A. M. Dirac, “Quantized Singularities in the Electromagnetic Field”, Proc. Roy. Soc. Lond. **A133** (1931), 60.
- [2] T. T. Wu and C. N. Yang, “Concept of Nonintegrable Phase Factors and Global Formulation of Gauge Fields”, Phys. Rev. **D12** (1975), 3845.
- [3] P. Goddard and D. Olive, Prog. Rep. Phys. **41** (1978), 1357.
- [4] P. Goddard, J. Nuyts and D. Olive, “Gauge Theories and Magnetic Charge”, Nucl. Phys. **B125** (1977), 1.
- [5] C. Montonen and D. Olive, “Magnetic Monopoles as Gauge Particles ?”, Phys. Lett. **B72** (1977), 177.
- [6] H. Georgi and S. Glashow, “Unified Weak and Electromagnetic Interactions Without Neutral Currents”, Phys. Rev. Lett. **28** (1972), 1494.
- [7] G. ‘t Hooft, “Magnetic Monopoles in Unified Gauge Theories”, Nucl. Phys. **B79** (1974), 276.
A. M. Polyakov, “Particle Spectrum in the Quantum Field Theory”, JETP Lett. **20** (1974), 194.
- [8] M. K. Prasad and C. M. Sommerfeld, “An Exact Classical Solution for the ‘t Hooft Monopole and the Julia-Zee Dyon”, Phys. Rev. Lett. **35** (1975), 760.
- [9] E. B. Bogomolny, “Stability of Classical Solutions”, Sov. J. Nucl. Phys. **24** (1976), 449.
- [10] A. A. Belavin, A. M. Polyakov, A. S. Swartz and Y. S. Tyupkin, “Pseudoparticle Solutions of the Yang-Mills Equations”, Phys. Lett. **B59** (1975), 85.
- [11] G. ‘t Hooft, “Computation of the Quantum Effects due to a Four Dimensional Pseudoparticle”, Phys. Rev. **D14** (1977), 3432.
- [12] R. Jackiw and C. Rebbi, “Degree of Freedom in Pseudoparticle Physics”, Phys. Lett. **B67** (1977), 189.
- [13] M. F. Atiyah N. Hitchin and I. M. Singer, “Self-Duality in Four Dimensional Riemannian Geometry”, Proc. Roy. Soc. Lond. **A362** (1978), 475.

- [14] M. F. Atiyah and I. M. Singer, “Index of Elliptic Operators I”, Ann. Math. **87** (1968), 485.
M. F. Atiyah and G. B. Segal, “Index of Elliptic Operators II”, Ann. Math. **87** (1968), 531.
M. F. Atiyah and I. M. Singer, “Index of Elliptic Operators III”, Ann. Math. **87** (1968), 546.
M. F. Atiyah and I. M. Singer, “Index of Elliptic Operators IV”, Ann. Math. **93** (1971), 119.
M. F. Atiyah and I. M. Singer, “Index of Elliptic Operators V”, Ann. Math. **87** (1971), 139.
- [15] G. ‘t Hooft, “Symmetry Breaking Through Bell-Jackiw Anomalies”, Phys. Rev. Lett. **37** (1976), 8.
- [16] R. Jackiw and C. Rebbi, “Vacuum Periodicity in a Yang-Mills Quantum Theory”, Phys. Rev. Lett. **37** (1976), 172.
- [17] C. G. Callan, R. Dashen and D. J. Gross, “Towards a Theory of the Strong Interactions”, Phys. Rev. **D17** (1978), 2717.
- [18] E. Witten, “Dyons of Charge $e\theta/2\pi$ ”, Phys. Lett. **B86** (1979), 283.
- [19] V. A. Rubakov, “Adler-Bell-Jackiw Anomaly and Fermionic-Number Breaking in the Presence of a Magnetic Monopole”, Nucl. Phys. **B203** (1982), 311.
- [20] G. ‘t Hooft, “A Property of Electric and Magnetic Charges in Non Abelian Gauge Theories”, Nucl. Phys. **B153** (1979), 141.
- [21] G. ‘t Hooft, “Some Twisted Self-Dual Solutions for the Yang-Mills Equations on a Hypertorus”, Commun. Math. Phys. **81** (1981), 267.
- [22] E. Witten, “Constraints on Supersymmetry Breaking”, Nucl. Phys. **B202** (1982), 253.
- [23] A. M. Polyakov, “Quark Confinement and Topology of Gauge Groups”, Nucl. Phys. **B120** (1977), 429.
- [24] I. Affleck, J. A. Harvey and E. Witten, “Instantons and Supersymmetry Breaking in $2 + 1$ Dimensions”, Nucl. Phys. **B206** (1982), 413.

- [25] C. Callias, “Axial Anomalies and Index Theorems on Open Spaces”, *Comm. Math. Phys.* **62** (1978), 213.
- [26] J. Wess and J. Bagger, “*Supersymmetry and Supergravity*”, Princeton University Press, Princeton, 1984.
- [27] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, “Exact Gell-Mann-Low Function of Supersymmetric Yang-Mills Theories From Instanton Calculus”, *Nucl. Phys.* **B229** (1983), 381.
- [28] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, “Instanton Effects in Supersymmetric Theories”, *Nucl. Phys.* **B229** (1983), 407.
- [29] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, “On Gluino Condensation in Supersymmetric Gauge Theories. $SU(N)$ and $O(N)$ Gauge Groups”, *Nucl. Phys.* **B296** (1988), 445.
- [30] D. Amati, K. Konishi, Y. Meurice, G. C. Rossi and G. Veneziano, “Non Perturbative Aspects in Supersymmetric Gauge Theories”, *Phys. Rep.* **162** (1988), 169.
- [31] I. Affleck, M. Dine and N. Seiberg, “Dynamical Supersymmetry Breaking in Supersymmetric QCD”, *Nucl. Phys.* **B241** (1984), 493.
- [32] I. Affleck, M. Dine and N. Seiberg, “Dynamical Supersymmetry Breaking in Four-Dimensions and its Phenomenological Implications”, *Nucl. Phys.* **B256** (1985), 557.
- [33] M. A. Shifman and A. I. Vainshtein, “On Gluino Condensation in Supersymmetric Gauge Theories, $SU(N)$ and $O(N)$ Gauge Groups”, *Nucl. Phys.* **B296** (1988), 445.
- [34] E. Cohen and C. Gómez, “Chiral Symmetry Breaking in Supersymmetric Yang-Mills”, *Phys. Rev. Lett.* **52** (1984), 237.
- [35] N. Seiberg and E. Witten, “Electric-Magnetic Duality, Monopole Condensation and Confinement in $N = 2$ Supersymmetric Yang-Mills Theory”, *Nucl. Phys.* **B426** (1994), 19.
- [36] N. Seiberg and E. Witten, “Monopoles, Duality and Chiral Symmetry Breaking in $N = 2$ Supersymmetric QCD”, *Nucl. Phys.* **B431** (1994), 484.
- [37] N. Seiberg and E. Witten, “Gauge Dynamics and Compactification to Three Dimensions”, **hep-th/9607163**.

- [38] M. Shifman and A. I. Vainshtein, “On Holomorphic Dependence and Infrared Effects in Supersymmetric Gauge Theories”, Nucl. Phys. **B359** (1991), 571.
- [39] N. Seiberg, “Supersymmetry and Nonperturbative Beta Functions”, Phys. Lett. **B206** (1988), 75.
- [40] K. G. Wilson and J. Kogut, “The Renormalization Group and the ϵ -Expansion”, Phys. Rep. **12** (1974), 75.
- [41] L. Adler and W. A. Bardeen, “Absence of Higher Order Corrections in the Anomalous Axial Vector divergence Equation”, Phys. Rev. **182** (1969), 1517.
- [42] M. Atiyah and N. Hitchin, “*The Geometry and Dynamics of Magnetic Monopoles*”, Princeton University Press, Princeton, 1988.
- [43] V. I. Arnold, “*Singularity Theory*”, London Math. Soc. Lecture Note Series **53**, Cambridge University Press, Cambridge, 1981.
- [44] M. Artin, “On Isolated Rational Singularities of Surfaces”, Amer. J. Math. **88** (1966), 129.
- [45] K. Kodaira, Ann. Math. **77**, 3 (1963), 563.
- [46] A. Klemm, W. Lerche, S. Theisen and S. Yankielowicz, “Simple Singularities and $N = 2$ Supersymmetric Yang-Mills”, Phys. Lett. **B344** (1995), 169.
- [47] P. Argyres and A. Faraggi, “The Vacuum Structure and Spectrum of $N = 2$ Supersymmetric $SU(N)$ Gauge Theory”, Phys. Rev. Lett. **73** (1995), 3931.
- [48] A. Hanany and Y. Oz, “On the Quantum Moduli Space of Vacua of $N = 2$ Supersymmetric $SU(N_c)$ Gauge Theories”, Nucl. Phys. **B452** (1995), 283.
- [49] P. Argyres, M. Plesser and A. Shapere, “The Coulomb Phase of $N = 2$ Supersymmetric QCD”, Phys. Rev. Lett. **75** (1995), 1699.
- [50] U. Danielsson and B. Sundborg, “The Moduli Space and Monodromies of $N = 2$ Supersymmetric $SO(2r + 1)$ Yang-Mills Theory”, Phys. Lett. **B358** (1995), 273.
- [51] A. Brandhuber and K. Landsteiner, “On the Monodromies of $N = 2$ Supersymmetric Yang-Mills Theory with Gauge Group $SO(2n)$ ”, Phys. Lett. **B358** (1995), 73.

- [52] K. Intriligator and N. Seiberg, “Lectures on Supersymmetric Gauge Theories and Electric-Magnetic Duality”, Nucl. Phys. Proc. Supl. **45** (1996).
- [53] C. Gómez and R. Hern’andez, “Electric-Magnetic Duality and Effective Field Theories”, **hep-th/9510023**; published in “**Advanced School on Effective Theories**”, F. Cornet and M. J. Herrero eds. World Scientific (1997).
A. Bilal, “Duality in $N = 2$ SUSY $SU(2)$ Yang-Mills Theory: A Pedagogical Introduction to the Work of Seiberg and Witten”, **9601007**.
W. Lerche, “Introduction to Seiberg-Witten Theory and its Stringy Origin”, Nucl. Phys. Proc. Supl. **B55** (1997), 83.
L. Álvarez-Gaumé and F. Zamora, “Duality in Quantum Field Theory (and String Theory)”, **hep-th/9709180**.
- [54] O. J. Ganor, D. R. Morrison and N. Seiberg, “Branes, Calabi-Yau Spaces, and Toroidal Compactification of the $N = 1$ Six-Dimensional E_8 Theory”, Nucl. Phys. **B487** (1997), 93.
- [55] Mandelstam, “Vortices and Quark Confinement in Non Abelian Gauge Theories”, Phys. Rep. **23** (1976), 245.
- [56] G. ‘t Hooft, “On the Phase Transition Towards Permanent Quark Confinement”, Nucl. Phys. **B138** (1978), 1.
- [57] H. B. Nielsen and P. Olesen, “Vortex Line Models for Dual Strings”, Nucl. Phys. **B61** (1973), 45.
- [58] G. ‘t Hooft, “Topology of the Gauge Condition and New Confinement Phases in Non Abelian Gauge Theories”, Nucl. Phys. **B190** (1981), 455.
- [59] M. B. Green, J. H. Schwarz and E. Witten, “*Superstring Theory*”, Cambridge University Press, Cambridge, 1987.
- [60] J. Polchinski, “What is String Theory?”, **hep-th/9411028**.
- [61] C. Vafa, “Lectures on Strings and Dualities”, **hep-th/9702201**.
- [62] E. Kiritsis, “Introduction to Superstring Theory”, **hep-th/9709062**.

- [63] J. D. Fay, “Theta Functions on Riemann Surfaces”, Lecture Notes in Mathematics, **352**, Springer-Verlag, 1973.
- [64] K. S. Narain, “New Heterotic String Theories in Uncompactified Dimensions < 10 ”, Phys. Lett. **B169** (1986), 41; K. S. Narain, M. H. Samadi and E. Witten, “A Note on the Toroidal Compactification of Heterotic String Theory”, Nucl. Phys. **B279** (1987), 369.
- [65] For reviews see A. Giveon, M. Porrati and E. Rabinovici, “Target Space Duality in String Theory”, Phys. Rep. **244** (1994), 77.
E. Álvarez, L. Álvarez-Gaumé and Y. Lozano, “An Introduction to T-Duality in String Theory”, **hep-th/9410237**.
- [66] P. Griffiths and J. Harris, “*Principles of Algebraic Geometry*”, Wiley-Interscience, 1978.
- [67] P. A. Aspinwall, “ $K3$ Surfaces and String Duality”, **hep-th/9611137**.
- [68] U. Persson, Lecture Notes in Mathematics, 1124, Springer-Verlag.
- [69] N. Seiberg, “Observations on the Moduli Space of Superconformal Field Theories”, Nucl. Phys. **B303** (1988), 286.
- [70] B. Greene and M. R. Plesser, “Duality in Calabi-Yau Moduli Space”, Nucl Phys. **B338** (1990), 15.
P. Candelas, M. Lynker and R. S. Schimmrigk, “Calabi-Yau Manifolds in Weighted $P(4)$ ”, Nucl. Phys. **B341** (1990), 383.
P. Candelas, X. de la Ossa, P. Green and L. Parkes, “A Pair of Calabi-Yau Manifolds as an Exactly Soluble Superconformal Theory”, Nucl. Phys. **B359** (1991), 21.
V. Batyrev, “Dual Polyhedra and Mirror Symmetry for Calabi-Yau Hypersurfaces in Toric Varieties”, **alg-geom/9410003**.
For a series of references, see S. -T. Yau ed., “Essays on Mirror Manifolds”, International Press, Hong-Kong, 1992.
- [71] I. V. Dolgachev, “Mirror Symmetry for Lattice Polarized $K3$ -Surfaces”, **alg-geom/9502005**.

- [72] J. Polchinski and Y. Cai, “Consistency of Open Superstring Theories”, Nucl. Phys. **B296** (1988), 91.
M. Green, “Space-Time Duality and Dirichlet String Theory”, Phys. Lett. **B266** (1991), 325.
J. Polchinski, “Dirichlet Branes and Ramond-Ramond Charges”, Phys. Rev. Lett. **75** (1995), 4724.
J. Polchinski, S. Chaudhuri and C. V. Johnson, “Notes on D-Branes”, **hep-th/9602052**, and “TASI Lectures on D-Branes”, **hep-th/9611050**.
- [73] J. C. Paton and H. M. Chan, “Generalized Veneziano Model with Isospin”, Nucl. Phys. **B10** (1969), 516.
- [74] N. J. Hitchin, A. Karlhede, U. Lindström, M. Rocek, “Hyperkähler Metrics and Supersymmetry”, Comm. Math. Phys. **108** (1987), 535.
- [75] J. Cardy and E. Rabonovici, “Phase Structure of $Z(P)$ Models in the Presence of a Theta Parameter”, Nucl. Phys. **B205** (1982), 1.
- [76] A. Font, L. Ibañez, D. Lüst and F. Quevedo, “Strong-Weak Coupling Duality and Non-Perturbative Effects in String theory”, Phys. Lett. **B249** (1990), 35.
- [77] A. Sen, “Strong-Weak Coupling Duality in Four Dimensional Field theory”, Int. J. Mod. Phys. **A9** (1994), 3707.
- [78] C. Hull and P. Townsend, “Unity of Superstring Dualities”, Nucl. Phys. **B438** (1995), 109.
- [79] M. Duff, “Strong/Weak Coupling Duality from the Dual String”, Nucl. Phys. **B442** (1995), 47.
- [80] P. Townsend, “The Eleven-Dimensional Supermembrane Revisited”, Phys. Lett. **B350** (1995), 184.
- [81] E. Witten, “String Theory Dynamics in Various Dimensions”, Nucl. Phys. **B443** (1995), 85.
- [82] A. Sen, “String-String Duality Conjecture in Six Dimensions and Charged Solitonic Strings”, Nucl. Phys. **B450** (1995), 103.

- [83] J. Harvey and A. Strominger, “The Heterotic String is a Soliton”, Nucl. Phys. **B449** (1995), 535.
- [84] B. Greene, D. Morrison and A. Strominger, “Black Hole Condensation and the Unification of String Vacua”, Nucl. Phys. **B451** (1995), 109.
- [85] C. Vafa and E. Witten, “A One-Loop Test Of String Duality”, Nucl. Phys. **B447** (1995), 261.
- [86] C. Hull and P. Townsend, “Enhanced Gauge Symmetries in Superstring Theories”, Nucl. Phys. **B451** (1995), 525.
- [87] J. Schwarz, “An $Sl(2, \mathbf{Z})$ Multiplet of Type IIB Superstrings”, Phys. Lett. **B360** (1995), 13.
- [88] D. J. Gross, J. A. Harvey E. Martinec and R. Rohm, “Heterotic String”, Phys. Rev. Lett. **54** (1985), 502; “Heterotic String Theory (I). The Free Heterotic String.”, Nucl. Phys. **B256** (1985), 253; “Heterotic String Theory (II). The Interacting Heterotic String.”, Nucl. Phys. **B267** (1986), 75.
- [89] S. Katz, D. R. Morrison and M. R. Plesser, “Enhanced Gauge Symmetry in Type II String Theory”, Nucl. Phys. **477** (1996), 105.
- [90] A. Klemm and P. Mayr, “Strong Coupling Singularities and Non-abelian Gauge Symmetries in $N = 2$ String Theory”, Nucl. Phys. **B469**, 37. (1996),
- [91] P. S. Aspinwall, “Enhanced Gauge Symmetries and $K3$ Surfaces”, Phys. Lett. **B357** (1995), 329..
- [92] P. S. Aspinwall and J. Louis, “On the Ubiquity of $K3$ Fibrations in String Duality”, Phys. Lett. **B369** (1996), 233..
- [93] A. Klemm, W. Lerche and P. Mayr, “ $K3$ -Fibrations and Heterotic-Type II String Duality”, Phys. Lett. **B357** (1995), 313.
- [94] C. Vafa, “Evidence for F-Theory”, Nucl. Phys. **B469** (1996), 403.
- [95] D. R. Morrison and C. Vafa, “Compactifications of F-Theory on Calabi-Yau Three-folds. I.”, Nucl. Phys. **B473** (1996), 74.

- [96] D. R. Morrison and C. Vafa, “Compactifications of F-Theory on Calabi-Yau Threefolds. II.”, Nucl. Phys. **B476** (1996), 437.
- [97] S. Kachru and C. Vafa, “Exact Results for $N = 2$ Compactifications of Heterotic Strings”, Nucl. Phys. **450**, (1995), 69.
- [98] M. J. Duff, P. Howe, T. Inami and K. S. Stelle, “Superstrings in $D = 10$ from Supermembranes in $D = 11$ ”, Phys. Lett. **B191** (1987), 70.
M. J. Duff, R. Minasian and J. T. Liu, “Duality Rotations in Membrane Theory”, Nucl. Phys. **B347** (1990), 394.
M. J. Duff, R. Minasian and J. T. Liu, “Eleven Dimensional Origin of String/String Duality: A One Loop Test”, Nucl. Phys. **B452** (1995), 261.
- [99] E. Witten, “Non-Perturbative Superpotentials in String Theory”, Nucl. Phys. **B474** (1996), 343.
- [100] A. Hanany and E. Witten, “Type IIB Superstrings, BPS Monopoles and Three Dimensional Gauge Dynamics”, Nucl. Phys. **B492** (1997), 152.
- [101] S. Elitzur, A. Giveon and D. Kutasov, “Branes and $N = 1$ Duality in String Theory”, Phys. Lett. **B400** (1997), 269.
- [102] J. de Boer, K. Hori, Y. Oz and Z. Yin, “Branes and Mirror Symmetry in $N = 2$ Supersymmetric Gauge Theories in Three Dimensions”, **hep-th/9702154**.
- [103] E. Witten, “Solutions of Four-Dimensional Fields Theories Via M-Theory”, Nucl. Phys. **B500** (1997), 3.
- [104] N. Evans, C. V. Johnson and A. D. Shapere, “Orientifolds, Branes and Duality of $4D$ Field Theories”, **hep-th/9703210**.
- [105] S. Elitzur, A. Giveon, D. Kutasov, E. Ravinovici and A. Schwimmer, “Brane Dynamics and $N = 1$ Supersymmetric Gauge Theory”, **hep-th/9704104**.
- [106] J. L. F. Barbon, “Rotated Branes and $N = 1$ Duality”, Phys. Lett. **B402** (1997), 59.
- [107] J. Brodie and A. Hanany, “Type IIA Superstrings, Chiral Symmetry, and $N = 1$ $4D$ Gauge Theory Dualities”, **hep-th/9704043**.

- [108] A. Brandhuber, J. Sommenschein, S. Theisen and S. Yanckielowicz, “Brane Configurations and 4D Field Theories”, **hep-th/9704044**.
- [109] O. Aharony and A. Hanany, “Branes, Superpotentials and Superconformal Fixed Points”, **hep-th/9704170**.
- [110] R. Tartar, “Dualities in 4D Theories with Product Gauge Groups from Brane Configurations”, **hep-th/9704198**.
- [111] I. Brunner and A. Karch, “Branes and Six Dimensional Fixed Points”, **hep-th/9705022**.
- [112] B. Kol, “5d Field Theories and M Theory”, **hep-th/9705031**.
- [113] A. Marshakov, M. Martellini and A. Morozov, “Insights and Puzzles from Branes: 4d SUSY Yang-Mills from 6d Models”, **hep-th/9706050**.
- [114] K. Landsteiner, E. López and D. A. Lowe, “ $N = 2$ Supersymmetric Gauge Theories, Branes and Orientifolds”, **hep-th/9705199**.
- [115] A. Brandhuber, J. Sommenschein, S. Theisen and S. Yanckielowicz, **hep-th/9705232**.
- [116] K. Hori, H. Ooguri and Y. Oz, ”Strong Coupling Dynamics of Four Dimensional $N = 1$ Gauge Theory from M theory Fivebrane” **hep-th/9706082**.
- [117] E. Witten, “Branes and the Dynamics of QCD”, **hep-th/9706109**.
- [118] A. Strominger, S. -T. Yau and E. Zaslow, “Mirror Symmetry is T -Duality”, Nucl. Phys. **B479** (1996), 243.
- [119] D. R. Morrison, “The Geometry Underlying Mirror Symmetry”, **alg-geom/9608006**.
- [120] R. Donagi and E. Witten, “Supersymmetric Yang-Mills and Integrable Systems”, Nucl. Phys. **B460** (1996), 299.
- [121] N. Hitchin, “The Self-Duality Equations on a Riemann Surface”, Proc. London Math. Soc. **55** (1987), 59; “Stable Bundles and Integrable Systems”, Duke Math. J. **54** (1987), 91.

- [122] S. Kachru, A. Klemm, W. Lerche P. Mayr and C. Vafa, “Non Perturbative Results on the Point Particle Limit of $N = 2$ Heterotic String Compactifications”, Nucl. Phys. **B459** (1996), 537.
- [123] C. Gómez, R. Hernández and E. López, “ S -Duality and the Calabi-Yau Interpretation of the $N = 4$ to $N = 2$ Flow”, Phys. Lett. **B386** (1996), 115.
- [124] A. Klemm, W. Lerche, P. Mayr, C. Vafa and N. Warner, “Self-Dual Strings and $N = 2$ Supersymmetric Field Theory”, Nucl. Phys. **B477**, (1996), 746.
- [125] C. Gómez, R. Hernández and E. López, “ $K3$ -Fibrations and Softly Broken $N = 4$ Supersymmetric Gauge Theories”, **hep-th/9608104**.
- [126] S. Katz and C. Vafa, “Geometric Engineering of Quantum Field Theories”, Nucl. Phys. **B497** (1997), 173, and “Geometric Engineering of $N = 1$ Quantum Field Theories”, Nucl. Phys. **B497** (1997), 196.
- [127] S. Katz, A. Klemm, C. Vafa, “Geometric Engineering of Quantum Field Theories”, Nucl. Phys. **B497** (1997), 173
- [128] M. Bershadsky, A. Johansen, T. Pantev, V. Sadov and C. Vafa, “F-Theory, Geometric Engineering and $N=1$ Dualities”, **hep-th/9612052**.
- [129] A. Gorskii, I. Krichever, A. Marshakov, A. Mironov and A. Morozov, “Integrability and Exact Seiberg-Witten Solution”, Phys. Lett. **B355** (1995), 466.
- [130] E. Martinec and N. P. Warner, “Integrable Systems and Supersymmetric Gauge Theories”, Nucl. Phys. **B459** (1996), 97.
- [131] C. Gómez, “Elliptic Singularities, θ -Puzzle and Domain Wall”, **hep-th/9711074**.
- [132] G. Dvali and M. Shifman, “Domain Walls in Strongly Coupled Theories”, Phys. Lett. **B396** (1997), 64.
- [133] A. Kovner, M. Shifman and A. Smilga, “Domain Walls in Supersymmetric Yang-Mills Theories”, **hep-th/9706089**.
- [134] A. Smilga and A. Vaselov, “Complex BPS Domain Walls and Phase Transition in Mass in Supersymmetric QCD”, **hep-th/9706217** and “Domain Walls Zoo in Supersymmetric QCD”, **hep-th/9710123**.

- [135] A. Kovner and M. Shifman, “Chirally Symmetric Phase of Supersymmetric Gluodynamics”, **hep-th/9702174**.
- [136] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, “Supersymmetric Instanton Calculus (Gauge Theories with Matter)”, Nucl. Phys. **B260** (1985), 157.
- [137] W. Fulton, “*Intersection Theory*”, Springer-Verlag, 1980.
- [138] A. Sen, “F-Theory and Orientifolds”, Nucl. Phys. B475 (1996), 562.
- [139] E. Witten, “Current Algebra Theorems for the $U(1)$ ‘Goldstone Boson’ ”, Nucl. Phys. B156 (1979), 269.
- [140] G. Veneziano, “ $U(1)$ without Instantons”, Nucl. Phys. B159 (1979), 213.
- [141] G. ‘t Hooft, “Dimensional Reduction in Quantum Gravity”, **gr-qc/9310026**.
- [142] J. D. Bekenstein, “Black Holes and Entropy”, Phys. Rev. **D7** (1973), 2333.
- [143] T. Banks, W. Fischler, S. H. Shenker and L. Susskind, “M-Theory as a Matrix Model: A Conjecture”, Phys. Rev. **D55** (1997), 5112.
- [144] T. Banks, “Matrix Theory”, **hep-th/9710231**.
- [145] W. Taylor, “D-Brane Field Theory on Compact Spaces”, Phys.Lett. **B394** (1997), 283.
O. J. Ganor, S. Ramgoolam, W. Taylor IV, “Branes, Fluxes and Duality in M(atrix)-Theory”, Nucl. Phys. **B492** (1997), 191.
- [146] M. Rozali, “Matrix Theory and U-Duality in Seven Dimensions”, Phys. Lett. **B400** (1997), 260.
- [147] S. Sethi and L. Susskind, “Rotational Invariance in the M(atrix) Formulation of Type IIB Theory”, Phys. Lett. **B400** (1997), 265.
- [148] A. Sen, “D0 Branes on T^n and Matrix Theory”, **hep-th/9709220**.
- [149] N. Seiberg, “Why is the Matrix Model Correct?”, **hep-th/9710009**.